



City Research Online

City St George's, University of London

Citation: Khantoul, B. & Fring, A. (2015). Time-dependent massless Dirac fermions in graphene. Physics Letters A, 379(42), pp. 2704-2706. doi: 10.1016/j.physleta.2015.08.011

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/12485/>

Link to published version: <https://doi.org/10.1016/j.physleta.2015.08.011>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Time-dependent massless Dirac fermions in graphene

Boubakeur Khantoul^{*,°} and Andreas Fring^{*}

^{*} *Department of Mathematics, City University London,
Northampton Square, London EC1V 0HB, UK*

[°] *Department of Physics, University of Jijel, BP 98, Ouled Aissa, 18000 Jijel, Algeria*
E-mail: bobphys@gmail.com, a.fring@city.ac.uk

ABSTRACT: Using the Lewis-Riesenfeld method of invariants we construct explicit analytical solutions for the massless Dirac equation in 2+1 dimensions describing quasi-particles in graphene. The Hamiltonian of the system considered contains some explicit time-dependence in addition to one resulting from being minimally coupled to a time-dependent vector potential. The eigenvalue equations for the two spinor components of the Lewis-Riesenfeld invariant are found to decouple into a pair of supersymmetric invariants in a similar fashion as the known decoupling for the time-independent Dirac Hamiltonians.

The two dimensional massless Dirac equation has recently attracted a lot of renewed attention because it describes quasi-particles in graphene [1, 2, 3], which is well known to possess a large amount of remarkable properties. Especially the Dirac equation in the presence of a magnetic field is of great interest as, unlike electrostatic potentials, such a configuration allows in principle to confine the Dirac fermions [4, 5, 6]. Many exact solutions have been provided for a variety of time-independent Hamiltonians and magnetic field configurations [7, 8, 9], including some for complex magnetic fields leading to pseudo/quasi-Hermitian interactions [10, 11]. While some solutions for the time-dependent Dirac equation in 1+1 dimensions have been constructed [12, 13, 14], little is known about the time-dependent setting with a magnetic field in 2+1 dimensions and no exact solutions have been reported. The aim of this manuscript is to commence filling that apparent gap. We shall demonstrate that the Lewis-Riesenfeld method of invariants [15] is a technique which can be employed successfully to solve this problem.

We consider here the time-dependent massless Dirac equation in two spacial dimensions in the form

$$H\Psi = i\partial_t\Psi, \quad \text{with } H(x, y, t) = \boldsymbol{\sigma} \cdot \mathbf{p}, \quad (1)$$

and the two component wave function $\Psi = (\psi_1, \psi_2)$. The effective Hamiltonian includes $\boldsymbol{\sigma} = (\sigma_x, \sigma_y)$ comprised of the standard Pauli matrices and $\mathbf{p}$ being the two-dimensional

momentum vector minimally coupled to a vector potential $A = (A_x, A_y)$ in the standard fashion $\mathbf{p} = (a(t)p_x + A_x, b(t)p_y + A_y)$. Besides the time-dependence entering through the vector potential, we allow here also for explicit time-dependent factors in front of the momenta, $a(t)$ and $b(t)$. One possibility to think of these factors is that they result from a time-dependent background, as discussed in [16] or alternatively as time-dependent velocities. The velocity of light, the reduced Planck constant and the charge are all set to one in our discussion, i.e., $c = \hbar = e = 1$.

We make now some specially simply choices for the vector potential by taking $A_x = 0$ and $A_y(x, t) = g(t)x$. In addition, we assume that the wave function separates with a free plane wave moving in the y -direction

$$\Psi(x, y, t) = e^{i(ky - \omega t)} \Phi(x, t), \quad (2)$$

with wave number k and frequency ω , such that we are left with the task to solve

$$\hat{H}\Phi = i\partial_t\Phi, \quad \text{with } \hat{H}(x, t) = a(t)\sigma_x p_x + [kb(t) + g(t)x]\sigma_y - \omega\mathbb{I}, \quad (3)$$

for the two component wave function $\Phi = (\phi_1, \phi_2)$. The first observation we make here is that the form of the explicit time-dependence does not allow for the standard decoupling of the systems into a pair of Hamiltonians related to each other by intertwining operators as common in the time-independent Dirac equation in analogy to standard supersymmetric quantum mechanics [17, 18].

We will attempt to solve equation (3) by using the Lewis-Riesenfeld method [15] originally designed to solve the time-dependent Schrödinger equation. The first step in this approach consists of solving the evolution equation

$$\frac{dI(t)}{dt} = \partial_t I(t) + \frac{1}{i}[I(t), \hat{H}(t)] = 0, \quad (4)$$

for the Hermitian time-dependent invariant $I(t)$. As usual in this context we take the invariant to be of the same order and form in the canonical variables as the Hamiltonian

$$I(t) = \alpha(t)p_x + \beta(t)x + \gamma(t), \quad (5)$$

where $\alpha(t)$, $\beta(t)$ and $\gamma(t)$ are now unknown time-dependent matrices. Substituting our Ansatz (5) into the evolution equation (4) yields six constraining equations for the three coefficient matrices

$$[\alpha, \sigma_x] = 0, \quad [\beta, \sigma_y] = 0, \quad g[\alpha, \sigma_y] + a[\beta, \sigma_x] = 0, \quad (6)$$

$$-i\dot{\alpha} = kb[\alpha, \sigma_y] + a[\gamma, \sigma_x], \quad (7)$$

$$-i\dot{\beta} = g[\gamma, \sigma_y], \quad (8)$$

$$-i\dot{\gamma} = kb[\gamma, \sigma_y] + ia\sigma_x\beta - ig\alpha\sigma_y. \quad (9)$$

Expanding the matrices in the $su(2)$ -basis, $\alpha(t) = \alpha_1(t)\mathbb{I} + \alpha_2(t)\sigma_x + \alpha_3(t)\sigma_y + \alpha_4(t)\sigma_z$ with $\alpha_i(t) \in \mathbb{R}$ for $i = 1, 2, 3, 4$ and similarly for $\beta(t)$, $\gamma(t)$, these equations are straightforward

to solve. Starting with (6), the first two equations immediately imply that $\alpha_3 = \alpha_4 = 0$ and $\beta_2 = \beta_4 = 0$. The last equation in (6) then yields $\beta_3 = \alpha_2 g(t)/a(t)$. Proceeding in this way for (7)-(9), we find the following form for the time-dependent invariant

$$I(t) = (\alpha_1 p_x + \gamma_1) \mathbb{I} + \alpha_2 p_x \sigma_x + [\beta_3 x + \gamma_3(t)] \sigma_y, \quad (10)$$

with constants $\alpha_1, \gamma_1, \alpha_2, \beta_3$. The time-dependence of $I(t)$ is entirely contained in the function $\gamma_3(t)$ which is constrained by

$$\dot{\gamma}_3(t) = \alpha_1 g(t), \quad \text{and} \quad \gamma_3(t) = k \alpha_2 \frac{b(t)}{a(t)} = k \beta_3 \frac{b(t)}{g(t)}. \quad (11)$$

In addition we found that $a(t) = \mu g(t)$ with $\mu = \alpha_2/\beta_3$ being constant has to be satisfied. The equations (11) are most conveniently solved in terms of $b(t)$

$$\gamma_3(t) = \left(2k\alpha_1\beta_3 \int^t b(s) ds \right)^{1/2}, \quad g(t) = \frac{k\beta_3}{2\alpha_1} b(s) \left(\int^t b(s) ds \right)^{-1/2}. \quad (12)$$

The next step in the Lewis Riesenfeld approach consists of solving the eigenvalue equation for the time-dependent invariant, i.e., we need to solve

$$I(t)\chi(t) = \lambda\chi(t), \quad (13)$$

for the time-dependent eigenfunction $\chi(t) = (\chi_+(t), \chi_-(t))$ and time-independent eigenvalues λ . For this purpose we note at first that we can write (13) as

$$\begin{pmatrix} 0 & L_- \\ L_+ & 0 \end{pmatrix} \begin{pmatrix} \chi_+ \\ \chi_- \end{pmatrix} = \frac{1}{\alpha_2} (\lambda - \alpha_1 - \gamma_1) \begin{pmatrix} \chi_+ \\ \chi_- \end{pmatrix}. \quad (14)$$

Thus we notice that unlike as the time-dependent Hamiltonian the invariant equation can be decoupled easily and acquires the form of a supersymmetric pair. Acting again with the off-diagonal invariant operator on (14) we obtain the two decoupled equations

$$I_{\pm} \chi_{\pm} = [p_x^2 + W^2 \pm W'] \chi_{\pm} = \frac{1}{\alpha_2^2} (\lambda - \alpha_1 - \gamma_1)^2 \chi_{\pm}, \quad (15)$$

for the two operators $I_{\pm} := L_{\mp} L_{\pm}$, where $W(x, t) = [\gamma_3(t) + \beta_3 x]/\alpha_2$ is the analogue to the superpotential in standard supersymmetric time-independent quantum mechanics. We observe that the potential is still time-dependent, but now a simple re-definition of our variables will move this dependence entirely into $\chi_{\pm}$. Defining the new time-dependent variable $\xi(t) = [\gamma_3(t) + \beta_3 x]/\alpha_2$ converts (15) into two eigenvalue equations for the time-independent quantum harmonic oscillator

$$\left(-\frac{1}{2} \frac{d^2}{d\xi^2} + \frac{\mu^2}{2} \xi^2 \right) \chi_{\pm} = \tilde{\lambda}^{\pm} \chi_{\pm}, \quad (16)$$

with $2\tilde{\lambda}_{\pm} = (\lambda - \alpha_1 - \gamma_1)^2/\beta_3^2 \mp \mu$. Demanding $\chi_{\pm}(\xi)$ to be a square integrable function $L^2(\mathbb{R}, d\xi)$ the solution is of course

$$\chi_{\pm, n}(\xi) = \frac{1}{\sqrt{2^n n!}} \left(\frac{\mu}{\pi} \right)^{1/4} e^{-\mu/2 \xi^2} H_n(\sqrt{\mu} \xi), \quad \tilde{\lambda}_n^{\pm} = \mu \left(n + \frac{1}{2} \right), \quad (17)$$

with H_n denoting the n -th Hermite polynomial. This means the two eigenvalues for the time-dependent spinor components of the invariant in (13) quantize to

$$\lambda_{n,s}^{\pm} = \alpha_1 + \gamma_1 + \beta_3 s \sqrt{\mu(2n+1 \pm 1)}, \quad (18)$$

with $s = \pm 1$ being two possible signs of the square root. As expected these eigenvalues are indeed time-independent. Due to the supersymmetric structure we have the standard shift in the eigenvalues, that is $\lambda_{n+1}^- = \lambda_n^+$.

As argued by Lewis and Riesenfeld [15] the eigenfunction of the Hamiltonian and the invariant just differ by a phase

$$|\Phi_n\rangle = e^{i\delta(t)} |\chi_n\rangle \quad (19)$$

where the real function $\delta(t)$ in (19) must obey

$$\frac{d\delta(t)}{dt} = \langle \chi_n | i\partial_t - \hat{H}(t) | \chi_n \rangle. \quad (20)$$

The right hand side of (20) can be computed directly with our known eigenfunctions. We obtain

$$\langle \chi_n | i\partial_t - \hat{H}(t) | \chi_n \rangle = \langle \chi_{+,n} | i\partial_t + \omega | \chi_{+,n} \rangle + \langle \chi_{+,n} | -ap_x + i(kb + gx) | \chi_{-,n} \rangle \quad (21)$$

$$+ \langle \chi_{-,n} | -ap_x - i(kb + gx) | \chi_{+,n} \rangle + \langle \chi_{-,n} | i\partial_t + \omega | \chi_{-,n} \rangle \\ = 2\omega, \quad (22)$$

where we used the orthogonality of the eigenfunctions, $\langle \chi_{\pm,n} | \partial_t | \chi_{\pm,n} \rangle = 0$, $\langle \chi_{+,n} | p_x | \chi_{-,n} \rangle = -\langle \chi_{-,n} | p_x | \chi_{+,n} \rangle$ and $\langle \chi_{+,n} | x | \chi_{-,n} \rangle = \langle \chi_{-,n} | x | \chi_{+,n} \rangle$. The phase therefore simply becomes

$$\delta(t) = 2\omega t. \quad (23)$$

Thus assembling the results from equations (2), (17), (19) and (23) provides an exact solution to the Dirac equation (1).

We have demonstrated that the Lewis-Riesenfeld method can be applied to construct solutions to the 2+1 dimensional time-dependent Dirac equation. The time-dependence resulted from a background and a magnetic field. Exact solutions for this type of scenario have not been known previously. Clearly there are plenty of open problems and challenges left. For instance, just as in the time-independent scenario one would like to know exact solutions for more complicated vector field configurations, different background scenarios and possibly different assumptions about the motion in the y -direction. These tasks are left for future work, where this note can be taken as encouragement as it demonstrates the successful applications of a method to tackle these kind of problems.

Acknowledgments: BK would like to thank Jijel University for financial support and City University London for kind hospitality. AF would like to thank Alessandro de Martino for useful discussions.

References

- [1] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, Two-dimensional gas of massless Dirac fermions in graphene, *Nature* **438**, 197–200 (2005).
- [2] A. K. Geim and K. S. Novoselov, The rise of graphene, *Nature Material* **6**, 183–191 (2007).
- [3] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, The electronic properties of graphene, *Rev. Mod. Phys.* **81**, 109–162 (2009).
- [4] A. De Martino, L. Dell’Anna, and R. Egger, Magnetic Confinement of Massless Dirac Fermions in Graphene, *Phys. Rev. Lett.* **98**, 066802 (2007).
- [5] L. Dell’Anna and A. De Martino, Multiple magnetic barriers in graphene, *Phys. Rev. B* **79**, 045420 (2009).
- [6] G. Giavaras, P. A. Maksym, and M. Roy, Magnetic field induced confinement-deconfinement transition in graphene quantum dots, *Journal of Physics: Condensed Matter* **21**(10), 102201 (2009).
- [7] V. Lukose, R. Shankar, and G. Baskaran, Novel Electric Field Effects on Landau Levels in Graphene, *Phys. Rev. Lett.* **98**, 116802 (2007).
- [8] T. K. Ghosh, Exact solutions for a Dirac electron in an exponentially decaying magnetic field, *Journal of Physics: Condensed Matter* **21**(4), 045505 (2009).
- [9] S. Kuru, J. Negro, and L. M. Nieto, Exact analytic solutions for a Dirac electron moving in graphene under magnetic fields, *Journal of Physics: Condensed Matter* **21**(45), 455305 (2009).
- [10] O. Panella and P. Roy, Pseudo Hermitian Interactions in the Dirac Equation, *Symmetry* **6**, 103–110 (2014).
- [11] C.-L. Ho and P. Roy, Dirac equation with complex potentials, *Modern Physics Letters A* **29**(40), 1450210 (2014).
- [12] R. R. Landim and I. Guedes, Wave functions for a Dirac particle in a time-dependent potential, *Phys. Rev. A* **61**, 054101 (Mar 2000).
- [13] A. S. de Castro and A. de Souza Dutra, Classes of exact wave functions for general time-dependent Dirac Hamiltonians in $1 + 1$ dimensions, *Phys. Rev. A* **67**, 054101 (2003).
- [14] M. Maamache and H. Lakehal, Solution of the generalized Dirac equation in a time-dependent linear potential: Relativistic geometric amplitude factor, *EPL (Europhysics Letters)* **67**(5), 695 (2004).
- [15] H. Lewis and W. Riesenfeld, An Exact quantum theory of the time dependent harmonic oscillator and of a charged particle time dependent electromagnetic field, *J. Math. Phys.* **10**, 1458–1473 (1969).
- [16] S. Dey and A. Fring, Noncommutative quantum mechanics in a time-dependent background, *Phys. Rev. D* **90**, 084005 (2014).
- [17] E. Witten, Dynamical breaking of supersymmetry, *Nucl. Phys.* **B188**, 513 (1981).
- [18] F. Cooper, A. Khare, and U. Sukhatme, Supersymmetry and quantum mechanics, *Phys. Rept.* **251**, 267–385 (1995).

- [19] A. Mostafazadeh, Pseudo-supersymmetric quantum mechanics and isospectral pseudo-Hermitian Hamiltonians Nucl. Phys. **B640**, 419–434 (2002).