



City Research Online

City St George's, University of London

Citation: Stefanski, B. (1999). WZ Couplings of D-branes and O-planes. Paper presented at the NATO Advanced Study Institute on Progress in String Theory and M-Theory, 24-05-1999 - 05-06-1999, Cargese, France.

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/12562/>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

WZ COUPLINGS OF D-BRANES AND O-PLANES

B. Stefański, jr.

*Department of Applied Mathematics and Theoretical Physics
Cambridge University, Silver Street, Cambridge CB3 9EW, UK
B.Stefanski@damtp.cam.ac.uk*

Abstract In this short note we review the construction and role of Wess-Zumino couplings of Dirichlet branes and Orientifold planes, and show how these combine to give the Green-Schwarz anomaly cancelling terms.

The low energy effective theory of a D-brane has two types of couplings, the Dirac-Born-Infeld (DBI) and Wess-Zumino (WZ) terms. The former describes the coupling of the vector potential and scalars coupling to the NS $\otimes$ NS fields, while the latter gives the coupling of the vector potential and the pullback of the curvature fields to the R $\otimes$ R potentials. The first D-brane WZ term [1] simply states that a D p -brane is charged under $C^{(p+1)}$, the $p + 1$ -form R $\otimes$ R potential

$$S_{WZ}^D = \mu_p \int_{\mathcal{B}_p} C^{(p+1)}, \quad (1.1)$$

where μ_p is the charge density of a D-brane and $\mathcal{B}_p$ its worldvolume. It was then shown that D-branes couple to lower dimensional R $\otimes$ R potentials [2] and that a ‘brane within brane’ picture emerged

$$S_{WZ}^D = \mu_p \int_{\mathcal{B}_p} C \wedge \text{tr} \exp(i\mathcal{F}/2\pi). \quad (1.2)$$

In the above C represents a formal sum of R $\otimes$ R potentials, $\mathcal{F} = F - B$, F is the worldvolume gauge field and B the NS $\otimes$ NS two form.¹

An R^4 term found in [3] required, as a consequence of duality, the presence of a gravitational WZ coupling [4]. Later the entire WZ action

¹ $\mathcal{F}$, rather than F , is present in the action as the latter quantity is not gauge invariant.

was found using the anomaly inflow mechanism [5, 6] and reads

$$S_{WZ}^D = \mu_p \int_{\mathcal{B}_p} C \wedge \text{tr} \exp(i\mathcal{F}/2\pi) \wedge \sqrt{\frac{\hat{A}(R_T)}{\hat{A}(R_N)}}, \quad (1.3)$$

where $\hat{A}$ is the Dirac or A-roof genus and R_T , R_N are the pull-backs of the tangent and normal bundle curvatures to the D-brane world-volume, respectively.

This coupling has a natural interpretation within K-theory [7]. In K-theory there is a natural bilinear pairing of bundles given by the index of the Dirac operator on the tensor product of the two bundles. For E a bundle over a manifold X , with TX the tangent space of X , the map

$$E \rightarrow \text{ch}(E) \sqrt{\hat{A}(TX)}, \quad (1.4)$$

is an isometry with respect to this pairing and the DeRham pairing in $H^*(X)$, the cohomology of X . In fact, following [13], it was shown [14] that D-brane charges are indeed classified by K-theory.

The D-brane WZ couplings were confirmed by string amplitude calculations. In [9] the four-form couplings were computed at tree level. A one-loop amplitude in [10] and tree-level amplitude in [11] confirmed the presence of all the couplings, while in [12] extra, non-anomalous couplings, as well as the normal bundle contributions were determined.

In [8], it was first observed that O-planes too carry gravitational WZ couplings. For consider Type I theory, in which 16 D9-branes, their images and an O9-plane, fill the spacetime. The WZ coupling of the theory is the Green-Schwarz (GS) coupling [15]. This differs from the WZ couplings of 32 D9-branes, indicating that O9-planes too have WZ couplings. By studying one-loop [10], and tree-level [11] scattering amplitudes it was found that the WZ coupling of Op-planes is

$$S_{WZ}^O = -2^{p-4} \mu_p \int_{\mathcal{B}_p} C \wedge \sqrt{\frac{L(R_T/4)}{L(R_N/4)}}, \quad (1.5)$$

where L is the Hirzebruch polynomial. To see that the O9-plane and D9-brane WZ couplings match the GS term consider the following. The massless chiral fields of Type I string theory are a neutral gravitino, a neutral fermion of opposite chirality and $\text{SO}(32)$ fermions. The total anomaly for this theory follows by descent from²

$$I = \frac{1}{2} 2\pi \left(\hat{A}(R) (\text{tr} e^{iR/2\pi} - 2) + \hat{A}(R) \text{tr} e^{iF/2\pi} \right) \Big|_{12\text{-form}}$$

²Note that all of these are real fields.

$$= \pi \frac{1}{(4\pi)^2} (\text{tr} R^2 - \text{tr} F^2) \wedge X_8, \quad (1.6)$$

where

$$X_8 = \frac{1}{(4\pi)^4} \left(\frac{2}{3} \text{tr} F^4 + \frac{1}{12} \text{tr} R^4 + \frac{1}{48} (\text{tr} R^2)^2 - \frac{1}{12} \text{tr} R^2 \text{tr} F^2 \right). \quad (1.7)$$

In the units of [16] the action extracted from the string theory amplitudes [10, 11] is

$$\begin{aligned} S = & -\frac{1}{4\kappa_{10}^2} \int dC^{(2)} \wedge * dC^{(2)} \\ & + \mu_9 \int \left(\frac{2}{(4\pi)^2} C^{(6)} \wedge (\text{tr} R^2 - \text{tr} F^2) + C^{(2)} \wedge X_8 \right). \end{aligned} \quad (1.8)$$

Since $H = dC^{(2)} + \dots$ is gauge invariant and $dC^{(6)} = *dC^{(2)}$ the gauge transformation for $C^{(2)}$ is

$$\delta C^{(2)} = 4\mu_9 \kappa_{10}^2 (\omega_{2,Y}^1 - \omega_{2,L}^1), \quad (1.9)$$

hence (1.8) has an anomalous variation which follows by descent from

$$I_{WZ} = 4\mu_9^2 \kappa_{10}^2 (\text{tr} F^2 - \text{tr} R^2) \wedge X_8. \quad (1.10)$$

The Type I charge density satisfies

$$(\mu_9 \kappa_{10})^2 = \frac{\pi}{2}, \quad (1.11)$$

and hence $I_{WZ} = I$ as required.

O-planes cannot couple to gauge fields so their WZ couplings have to be purely gravitational. Hence it is a consistency check for the D9-brane and GS gauge and mixed couplings to agree.

Acknowledgments

The author would like to thank the conference organisers for financial support and a stimulating programme.

References

- [1] J. Polchinski, *Dirichlet-Branes and Ramond-Ramond Charges*, Phys. Rev. Lett. **75** (1995) 4724, hep-th/9510017;
- [2] M. Li, *Boundary States of D-branes and Dy-Strings*, Nucl. Phys. **B460** (1996) 351, hep-th/9510161; C. Schmidhuber, *D-Brane Actions*, Nucl. Phys. **B467** (1996) 146, hep-th/9601003; M.R. Douglas, *Branes within Branes*, hep-th/9512077;

- [3] C. Vafa and E. Witten, *A One Loop Test of String Dualities*, Nucl. Phys. **B447** (1995) 261, hep-th/9505053;
- [4] M. Bershadsky, C. Vafa, V. Sadov, *D-Branes and Topological Field Theories*, Nucl. Phys. **B463** (1996) 420, hep-th/9511222;
- [5] M.B. Green, J.A. Harvey and G. Moore, *I-Brane Inflow and Anomalous Couplings on D-branes*, Class. Quant. Grav. **14** (1997) 47, hep-th/9605033;
- [6] Y.-K.E. Cheung and Z. Yin, *Anomalies, Branes, and Currents*, Nucl. Phys. **B517** (1998) 69, hep-th/9710206;
- [7] R. Minasian and G. Moore, *K-theory and Ramond-Ramond Charge*, JHEP **9711** (1997) 002, hep-th/9710230;
- [8] K. Dasgupta, D.P. Jatkar, S. Mukhi, *Gravitational Couplings and Z_2 Orientifolds*, Nucl. Phys. **B523** (1998) 465, hep-th/9707224;
- [9] B. Craps and F. Roose, *Anomalous D-brane and Orientifold Couplings from the Boundary State*, Phys. Lett. **445B** (1998) 150, hep-th/9808074;
- [10] J.F. Morales, C.A. Scrucca, M. Serone, *Anomalous Couplings for D-branes and O-planes*, Nucl. Phys. **B552** (1999) 291, hep-th/9812071;
- [11] B. Stefański, jr., *Gravitational Couplings of D-branes and O-planes*, Nucl. Phys. **B548** (1999) 275, hep-th/9812088;
- [12] B. Craps and F. Roose, *(Non-)Anomalous D-brane and O-plane couplings: the Normal Bundle*, Phys. Lett. **450B** (1999) 358, hep-th/9812149;
- [13] For a review see A. Sen, *Non-BPS States and Branes in String Theory*, hep-th/9904207;
- [14] E. Witten, *D-Branes And K-Theory*, JHEP **9812** (1998) 025, hep-th/9810188;
- [15] M.B. Green and J.H. Schwarz, *Anomaly Cancellations in Supersymmetric $D=10$ Gauge Theory and Superstring Theory*, Phys. Lett. **149B** (1984) 117;
- [16] J. Polchinski, *String Theory, Volume 2, Superstring Theory and Beyond*, CUP 1998;