



City Research Online

City St George's, University of London

Citation: Valmaggia, L. R., Day, F. & Rus-Calafell, M. (2016). Using virtual reality to investigate psychological processes and mechanisms associated with the onset and maintenance of psychosis: a systematic review. *Social Psychiatry and Psychiatric Epidemiology*, 51(7), pp. 921-936. doi: 10.1007/s00127-016-1245-0

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/14876/>

Link to published version: <https://doi.org/10.1007/s00127-016-1245-0>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Using virtual reality to investigate psychological processes and mechanisms associated with the onset and maintenance of psychosis: a systematic review

Lucia R. Valmaggia^{1,*},²

Email Lucia.Valmaggia@kcl.ac.uk

Fern Day^{1,3}

Mar Rus-Calafell^{1,2}

¹ Department of Psychology (PO 77), Institute of Psychiatry, Psychology and Neuroscience, King's College London, De Crespigny Park, London, SE5 8AF UK

² South London and Maudsley NHS Trust, London, UK

³ City University London, London, UK

Abstract

Purpose

In the last decade researchers have embraced virtual reality to explore the psychological processes and mechanisms that are involved in the onset and maintenance of psychosis. A systematic review was conducted to synthesise the evidence of using virtual reality to investigate these mechanisms.

AQ1

Methods

Web of Science, PsycINFO, Embase, and Medline were searched.. Reference lists of collected papers were also visually inspected to locate any relevant cited journal articles. A total of 6001 articles were potentially eligible for inclusion; of these, 16 studies were included in the review.

Results

The review identified studies investigating the effect of interpersonal sensitivity, childhood bullying victimisation, physical assault, perceived ethnic discrimination, social defeat, population density and ethnic density on the real-time appraisal of VR social situations. Further studies demonstrated the potential of VR to investigate paranoid ideation, anomalous experiences, self-confidence, self-comparison, physiological activation and behavioural response.

Conclusions

The reviewed studies suggest that VR can be used to investigate psychological processes and mechanisms associated with psychosis. Implications for further experimental research, as well as for assessment and clinical practise are discussed. The present review has been registered in the PROSPERO register: CRD42016038085.

Keywords

Virtual reality

Adverse life events

Daily stressors

Stress sensitivity, psychosis

Introduction

Cognitive models of psychosis have investigated the psychological processes and mechanisms that play a role in psychosis and suggest that stressful life events can influence the way we perceive ourselves, others, and the world around us and can contribute to the onset and maintenance of psychosis [1–5]. These models have been supported by research which suggest that stress is a key feature in the onset and maintenance of mental health symptoms and disorders [6, 7]. Stress can be defined as a physiological reaction to a threat that involves the activation of the hypothalamic–pituitary–adrenal (HPA) axis [8–10]. This reaction is similar across all mammalian species. In humans the psychological conceptualisation of stress proposes that the emotional response and physiological activation that occur in a situation are dependent on the interpretation of a threat and on whether we think we will be able to cope with the event or not [11]. Stress is a dominant feature of many aetiological models of psychosis [9, 12, 13], however, stress is an elusive concept that is defined

differently in different contexts. Reported stressful events can range from mild feelings of everyday stress and minor hassles (for example, living in a densely populated city; travelling in crowded public transport), through significant adverse life events (for example, bully victimisation; ethnic discrimination), to traumatic events (for example, physical assault; sexual abuse).

A number of social factors that can be viewed as stressors have themselves been linked to an increased risk of psychosis. Urbanicity has been associated with higher levels of stress [14] and higher rates of psychosis [15, 16]. Higher psychosis rates have ~~also~~ been found among migrants [17, 18]. ~~Higher incidence rates of psychosis and~~ have been linked to the perception of ethnic density [19, 20]. ~~and~~ Furthermore, associations between perceived ethnic discrimination and psychosis in ethnic minority and immigrant groups have been reported in previous research [21–23]. Childhood bully victimisation has also been associated with an increase in psychosis and other mental health problems in adulthood [24, 25].

Also worth considering in this context is the evidence showing that individuals with schizophrenia have an increased sensitivity to daily hassles and report greater subjective stress [12, 26, 27]. These findings have stimulated interest in stress reactivity as an element of vulnerability [28] and an interest in how individuals with different levels of vulnerability for psychosis respond to daily stressors [29–31]. Interpersonal sensitivity is described as “an undue and excessive awareness of, and sensitivity to, the behaviour and feelings of others... particularly to perceived or actual situations of criticism or rejection...” (p. 342 [32]). Higher interpersonal sensitivity is linked to problems in interpersonal relationships, low self-esteem, as well as negative self-concepts, and it has been found to be higher in individuals at ultra high risk for psychosis [31, 33]. Previous studies have also found that a negative self-concept [34, 35]) and low or unstable self-esteem [34] are linked to higher levels of paranoia in clinical and non-clinical samples.

AQ2

When thinking about possible underlying mechanisms in the onset of psychosis is important to consider that prolonged exposure to social adversity and social exclusion have been linked to high prevalence rates of anomalous experiences (such as a propensity to hallucinations, high endorsement of psychic items or higher trait paranoia) in the general population and have been found to be associated with the same risk factors present in people with psychosis [36–40]. Stressful life events have also been linked to paranoid ideation [41]. Paranoid ideation has been defined as “the unfounded fear that others intend to cause you harm” [42, 43] and is the most common type of delusional belief recorded by

individuals with psychosis [44] and it is a common delusional belief in the general population [45].

AQ3

While there is a long tradition of research in these processes and mechanisms they are usually assessed with retrospective self-report questionnaires or interviews. Limited evidence is available researching underlying psychological processes and mechanisms in real-life or in ecologically valid environments [46, 47]. In the last decade, researchers in the field of psychosis have embraced Immersive Virtual Reality (hereafter shortened to VR) as a method to explore the psychological mechanisms associates with the onset and maintenance of psychosis. VR environments are displayed in colour and in 3D using a head-mounted-display. The user is fully immersed in the environment and can interact with the environment either by head movements, full body turning or with a joystick. The sounds of the environment are heard through headphones [48] (please see <https://www.youtube.com/watch?v=svG6kXC2PQA> for an example). VR environments elicit sense of presence [49] and trigger real time cognitive, emotional, behavioural, and physiological responses to real time situations [50]. This sense of ‘being there’ as if it were a real life situation has lead researcher to describe VR environments as ecologically valid [50, 51]. Aside from allowing to simulate “real life” social situation and bring them into a laboratory setting, the added value of VR is that it allows the researcher to measure real time responses, and at the same time it enables to control and manipulate each aspect of an environment [52]. VR enables thus researchers to expose participants to exactly the same social environment and to measure the individual cognitive, emotional, behavioural and physiological response to the environment. It also enables to investigate the association between an individual prior exposure to social and environmental factors (e.g., history of migration, or bully victimisation) and their real-time response to a social environment. The VR environment itself can also be manipulated, for example, the number of virtual characters in an environment can be changed, as can be their ethnicity, their facial expression, their behaviour or what they say. Sounds, like for example indistinct background conversations, crowd murmuring or laughter or other noises like background music or sound of traffic can also be manipulated.

The aim of this manuscript was to review experimental studies that used VR to investigate the underlying psychological processes and mechanisms associated with the onset and maintenance of psychosis.

Method

Design

We conducted a systematic review of the literature and described the results in a qualitative synthesis. The review was registered on the PROSPERO register CRD42016038085.

Selection procedure

The databases used to conduct the search were Web of Science, PsycINFO, Embase, and Medline. Reference lists of collected papers were also visually inspected to locate any relevant cited journal articles. Studies were included in the review if they were: written in English; contained original empirical findings, and published in a peer-reviewed journal. Studies were excluded from the review if they were treatment studies; case studies; reviews; not available in English; theses; or book chapters. Only studies of immersive VR were included.

Search criteria

Studies for review were identified following a keyword search for the terms ‘virtual reality’ OR ‘VR’ in conjunction with ‘social stress’ OR ‘adverse life events’ OR ‘urban’, OR ‘migration’, OR ‘bullying’, OR ‘discrimination’, OR ‘social defeat’, OR ‘social entrapment’, OR ‘stress’, AND ‘psychological processes’, OR ‘mechanisms’, OR ‘mental health’, OR ‘paranoia’, OR ‘anxiety’, OR ‘depression’, OR ‘trauma’, OR ‘psychosis’ OR ‘schizophrenia’. Appropriate truncations and wild cards were used to identify permutation of the terms searched, e.g., stress* to search for: stress, stressors, stressful; paranoi* to search for: paranoia, paranoid.

Quality assessment

Quality of the reviewed studies was assessed independently by LV and MRC using the Evaluation of Public Health Practise Project (EPHPP) Quality Assessment Tool for Quantitative Studies [53]. A global rating for the paper is described as follows: Strong = no weak ratings; Moderate = one weak rating; Weak = two or more weak ratings on the subscales (selection bias; study design; confounders; blinding; data collection methods; withdrawals and dropouts) The EPHPP is available online (<http://ephpp.ca/tools.html>). Rating disagreements were resolved by consensus.

Results


Information extraction

The search resulted in 5999 articles (final search conducted 4 March, 2016); two more manuscripts were identified by publication alerts email from journals.

From these 6001 studies we identified 16 experimental studies which used full

immersive VR to research adverse life events, daily stressors and psychological processes associated with psychosis (see Fig. 1).

Fig. 1
PRISMA 2009 flow diagram



Qualitative synthesis

The reviewed studies were organised in two categories. The first category describes studies which used VR to investigate the association between baseline characteristics and the real-time response to a neutral social VR environment. The second category includes studies that manipulated one or more aspects of the environment to measure how the manipulation would influence the real-time response to VR environment. Finally, we reviewed overarching findings from both categories regarding whether VR can be used to investigate real-time response to social situations.

Studies exploring the association between baseline characteristics and the real-time response to a neutral social VR environment

The ten studies summarised in Table 1 used VR to explore the association between existing characteristics of participants and their response to a VR social situation. In two studies the environment was a library, the remaining studies used an underground train environment. Both the library environment and the underground were designed to be perceived as neutral social environments. Virtual characters or avatars showed neutral behaviours typical of such social situations, but that could be interpreted as ambiguous like for example, looking at the participant, smiling, or avatars talking to each other. This was done on purpose as the main aim of the studies was to explore ‘unfounded’ real-time paranoid ideation in VR rather than a response to a genuinely hostile environment [54].

Table 1

Studies exploring the association between baseline characteristics of participants and the cognitive response to a neutral social VR environment to ($n = 10$)

Study (country)	Area studied	Design	Participants	Mean age (SD)	VR equipment
Freeman et al. [61] (United Kingdom)	Persecutory ideation	Within subjects design	24 healthy participants (12 male and 12 females)	26 (6)	CAVE Immersive projection system and Crystal Eyes shutter-glasses
Freeman et al. [62] (United Kingdom)	Persecutory ideation	Within subjects design	30 healthy participants (15 males and 15 females; 70 % White ethnicity)	22 (5)	CAVE Immersive projection system and Crystal Eyes shutter-glasses

Freeman et al. [70] (United Kingdom)	Paranoid ideation and social anxiety	Within subjects design	200 participants from general population (100 males and 100 females; 63 % White ethnicity; 47 % graduate degree or higher; 17 % unemployed)	37.5 (13.3)	HMD: VR1280
Freeman et al. [64] (United Kingdom)	Persecutory ideation and associated factors	Within subjects design	Group 1 ($n = 30$): low non-clinical paranoia (83 % White ethnicity) Group 2 ($n = 30$): high non-clinical paranoia (77 % White ethnicity) Group 3 ($n = 30$): Clinical Persecutory delusions (53 % White ethnicity) Each group had 18 males and 12 females	Group 1: 44.2 (11.2) Group 2: 36 (11.7) Group 3: 44.2 (11.7)	HMD: VR1280
Freeman et al. [57] (United Kingdom)	Paranoia and PTSD following an assault	Within subject comparison	106 participants recruited in an A&E department after having experienced a distressing physical assault (79 males and 27 females; 52 % White ethnicity; 31 % unemployed)	34.4 (11.6)	HMD: NVIS SX111

						1
Freeman et al. [56] (United Kingdom)	Paranoia in VR as predictor of paranoia and PTSD 6 months after experiencing an assault	Within subject comparison	As above	As above	As above	2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214

Valmaggia et al. [55] (United Kingdom)	Childhood bullying victimisation and paranoid ideation	Cross sectional between participants independent design	64 Ultra High Risk individuals (UHR) (38 males 26 females; 36 % White British; 56 % Unemployed; 13 % Higher education 43 Healthy Controls (20 males, 23 females; 37 % White British; 7 % Unemployed; 33 % Higher education)	UHR 22.5 (4.0) HC 24 (4.0)	HMD: VR1280
Valmaggia et al. [59] (United Kingdom)	Social defeat and paranoid ideation	Cross sectional between participants independent design	As above	As above	HMD: VR1280

HMD Head Mounted Display, *GPTS-B* Green Paranoia Thoughts Scale-B, *VAS* Visual Anxious State Paranoia Scale, *GSR* Galvanic skin response, *HR* Heart rate, *STAI* Spielberg Trait Anxiety Scale, *BSI* Brief Symptom Inventory, *LSHS* Launay Slade Hallucination Scale, *DASS* Depression Measure, *PSCS* Private Self-Consciousness Scale, *SADS* Social Avoidance and distress scale, *PSWQ* Penn State Worry questionnaire, *BCSS* Brief core schema scale, *CAPS* Cardiff Anxiety Scale, *SSQ* Social support questionnaire, *SELSA* Social and emotional loneliness scale for adults, *NSS* Negative Symptom Scale, *CPQ* Cognitive processing questionnaire, *MDS* Mental defeat Qualities questionnaire, *RIQ* Response to intrusion questionnaire, *SBQ* Safety behaviour Crisis support scale, *PSRS* Psychotic Symptoms rating Scale, *PEDQ* Perceived Ethnic Discrimination Interaction Anxiety Scale, *DACOBS* Davos Assessment of Cognitive Biases Scale, *RSES* Revised Social Skills Scale

Childhood bullying victimisation, physical assault, perceived ethnic discrimination, and social defeat

Five studies investigated whether having experienced adverse life events was associated with higher levels of paranoid ideations in the neutral social VR environments. All five studies used the VR underground train environment. Childhood bullying victimisation [55]; being the victim of a physical assault [56, 57]; perceived ethnic discrimination [58] and social defeat [59] were found to be associated with real-time paranoid ideation in the VR social

environment.

Interpersonal sensitivity

The correlation between interpersonal sensitivity and real-time paranoid ideation during VR was investigated in five studies [60–64]. All studies found that heightened interpersonal sensitivity was associated with higher levels of paranoia in the VR library or in the underground train.

Studies investigating the effect of a manipulation on the response to a VR environment

The six studies listed in Table 2 have used a manipulation to investigate the real-time response to the VR environment. The studies have used different environments: a bar, a student flat, and the underground train described earlier.

Table 2


Studies exploring the cognitive, emotional, behavioural, and physiological response to a mar

Study (country)	Area studied	Design	Participants	Mean age (SD)	VR equipment
Atherton et al. [67] (United Kingdom)	Self-confidence and paranoid ideation	Within subjects design	26 males (from a general population sample who scored >17 on GPTS-Paranoia)	43.4 (16.3)	HMDs: NVIS SX111 or VR1280
Brinkman et al. [69] (The Netherlands)	Physiological response to changes in population density and ethnic density	Two by two within-subject design	2 Males (1 with schizoaffective disorder) 24 Healthy Controls (20 male and 4 female; all	29 (9.2)	HMD: Emagin Z800 3DVisor

			White Dutch)		
Freeman et al. [68] (United Kingdom)	Persecutory ideation and social comparison after height was altered	Two condition, two period crossover trail	60 females from the general population (with paranoid thinking in the last month; 78 % White ethnicity; 55 % graduate degree or higher)	31.5 (13)	HMD: NVIS SX111
Fornells-Ambrojo et al. [60] (United Kingdom)	Perception of trustworthiness and interpersonal distance	Between group comparison	61 healthy male (77 %: White ethnicity; 90 % students)	25.3 (7.3)	CAVE Immersive projection system and Crystal Eyes shutter-glasses
Veling et al. [65] (The Netherlands)	Paranoid thoughts and associated social anxiety, cognitive bias, self esteem	Cross sectional between participants independent design	17 patients with a first episode of psychosis (FEP) (14 males, 3 females; 35 % White Dutch ethnicity) 24 healthy controls (20 males, 4 females; 100 % White Dutch ethnicity)	FEP 27.3 (5.5) HC 29 (9.2)	Emagin Z800 3D Visor
			Group 1: 20 UHR (7 males, 13 females 75 % White		

Veling et al. [66] (The Netherlands)	Paranoid thoughts and stress sensitivity	Cross sectional between participants independent design	Dutch ethnicity) Group 2: 55 Individuals with psychosis (42 males, 13 females; 53 % White Dutch ethnicity) Group 3: 42 siblings of people with psychosis (23 males, 19 females; 75 % White Dutch ethnicity) Group 4: 53 controls with negative first degree family history of psychosis (25 males, 28 females; 70 % White Dutch ethnicity)	UHR 24 (4.5) Psychosis 26 (4.7) Siblings 26.4 (4.8) Controls 24.6 (4.4)	Sony HMZ-T1
--------------------------------------	--	---	---	--	----------------

HMD Head Mounted Display, *GPTS-B* Green Paranoia Thoughts Scale-B, *VAS* Visual An State Paranoia Scale, *GSR* Galvanic skin response, *HR* Heart rate, *STAI* Spielberg Trait A Scale, *BSI* Brief Symptom Inventory, *LSHS* Launay Slade Hallucination Scale, *DASS* Depression Measure, *PSCS* Private Self-Consciousness Scale, *SADS* Social Avoidance and distress scale, *Penn State* Worry questionnaire, *BCSS* Brief core schema scale, *CAPS* Cardiff Anomalous Social support questionnaire, *SELSA* Social and emotional loneliness scale for adults, *SC* Symptom Scale, *CPQ* Cognitive processing questionnaire, *MDS* Mental defeat Scale, *TM* questionnaire, *RIQ* Response to intrusion questionnaire, *SBQ* Safety behaviour questionnaire support scale, *PSRS* Psychotic Symptoms rating Scale, *PEDQ* Perceived Ethnic Discrimination Interaction Anxiety Scale, *DACOB*S Davos Assessment of Cognitive Biases Scale, *RSES*



Population density and ethnic density

Two studies investigated how manipulating population density and ethnic density effected paranoid ideation in a VR bar environment [65 , 66]. The findings suggested that increased population density, hostility of the avatars, and a different ethnicity were associated with higher levels of paranoia in the VR bar.

Self-confidence and social comparison

Two experiments indicated that VR can be useful to measure self-confidence and social comparison. In the first study, participants underwent a manipulation aimed to induce low or high self-confidence. In the high self confidence induction they were instructed to think about a time in their life when they felt self-confident and they were provided with a list of confidence statements. In the low-confidence induction, participants underwent a similar procedure but regarding their least self-confident experience. Low self-confidence was

associated with higher paranoid ideation in VR underground train [67].

The second experiment manipulated the height of participants in the VR environment and found that a lowered height was associated with negative social-comparison [68].

Physiological activation and behavioural response

Manipulation of the VR environment triggered not only the cognitive and emotional response to a social situation, but also the physiological changes and behavioural response. Galvanic skin response, heart rate and behavioural response (distance kept from the avatars) was found to be triggered by changes in population density and ethnic density in a VR bar environment [69]. The same environment was used to investigate galvanic skin response in a sample of individuals with psychosis [65]. Distance kept from the avatar was also measured in a study which manipulated the contingency behaviour of avatars while participants were exploring a VR student flat [60].

VR to investigate real time responses to social situations

Paranoid ideations

As summarised in Tables 1 and 2, the large majority of studies investigated paranoid ideation and the associated correlates. VR was found to be a valid research method, not only to elicit and measure paranoid ideation in participants, but also to generate participant's attributions about positive, neutral and negative mental states to virtual characters as if they were real people [55, 58, 59, 61–63, 65, 66]. Furthermore, as illustrated in Tables 1 and 2 VR was found to be effective in generating paranoid ideation in people from general population samples [56, 57, 60–62, 68, 70] as well as from individuals with an ultra high risk for psychosis [55, 58, 59, 63, 66, 71] and those already experiencing a psychotic disorder [65, 66].

Anomalous experiences

Individuals experiencing anomalous experiences (such as a propensity to hallucinations, high endorsement of psychic items or higher trait paranoia) reported higher levels of paranoid ideation when exposed to a VR environment [60, 62–64, 66, 70]. These results confirm the validity of VR to assess real-time paranoia in response to a social situation, as it would be expected that an individual who reports trait paranoia would also experience paranoid ideation in an ecologically valid social VR environment.

Quality assessment

Several studies reached a strong rating on the EPHPP global rating (see Tables 1, 2, last column). No studies were excluded based on their quality. However, it is worth mentioning that a strong global rating on the EPHPP allows a moderate score on one of more subscales. Table 3 illustrates the subscale scores for each study: the majority of studies ~~reach-moderate~~reached moderate scores on selection bias, indicating that the participants recruited for the study were not very likely to be representative of the target population. Indeed quite a few studies were conducted with healthy controls and student samples. The majority of studies had a within subject design, ~~they~~ did not always control~~led~~ for confounders, and ~~often~~ the outcome assessor was not ~~always~~ blind to the experimental condition. Most studies had a sound data collection method and few reported withdrawals or drop-outs.

Table 3

Immersive VR studies EPHPP quality assessment rating

Study	A	B	C	D	E	F
	Selection bias	Study design	Confounders	Blinding	Data collection method	Withdraw and drop out
Atherton et al. [67]	Moderate	Moderate	Strong	Moderate	Strong	Strong
Brinkman et al. [69]	Weak	Moderate	Weak	Weak	Strong	Strong
Fornells-Ambrojo et al. [60]	Weak	Strong	Moderate	Moderate	Strong	Strong
Freeman et al. [61]	Moderate	Moderate	Weak	Moderate	Moderate	Strong
Freeman et al. [62]	Strong	Moderate	Weak	Moderate	Strong	Strong
Freeman et al. [70]	Moderate	Moderate	Strong	Moderate	Strong	Strong
Freeman et al. [64]	Strong	Moderate	Strong	Moderate	Strong	Strong
Freeman et al. [68]	Strong	Moderate	Strong	Moderate	Strong	Strong
Freeman et al. [56]	Strong	Moderate	Strong	Moderate	Strong	Strong
Freeman et al. [57]	Moderate	Moderate	Strong	Moderate	Strong	Strong
Shaikh et al. [58]	Moderate	Moderate	Strong	Moderate	Strong	Moderate

Valmaggia et al. [63]	Moderate	Moderate	Moderate	Moderate	Strong	Moderate
Valmaggia et al. [55]	Moderate	Moderate	Moderate	Moderate	Strong	Moderate
Valmaggia et al. [59]	Moderate	Moderate	Strong	Moderate	Strong	Moderate
Veling et al. [65]	Moderate	Moderate	Moderate	Moderate	Moderate	Moderate
Veling et al. [66]	Moderate	Moderate	Moderate	Moderate	Moderate	Moderate

AQ4

Discussion

We sought to review experimental studies that used VR to investigate the underlying psychological processes and mechanisms associated with the onset and maintenance of psychosis. The term stress was considered to encompass a broad range of events and experiences encountered in the psychosocial domain that provoke feelings of distress, mental strain, or physiological symptoms of stress. These experiences included traumatic events, social adversity, stressful life events, and subjective feelings of stress and inability to cope.

Childhood bullying victimisation [55]; being the victim of a physical assault [56, 57]; perceived ethnic discrimination [58] and social defeat [59] were associated with paranoid ideation in a VR social environment. The ability to use VR to study the effect of adverse life events on real-time response to social situations in a controlled environment offers an exciting new way to research what contributes to stress sensitivity. Measuring the real-time physiological activation in social situations offers a number of possibilities not only for research purposes but also for assessment and treatment [10, 47]. Interestingly, VR has been shown to be an effective way to overcome excessive self-criticism in people with depression [72, 73], suggesting that VR can be not just a method to assess components of interpersonal sensitivity, but that it also has the potential to be used in the treatment of decrease interpersonal sensitivity.

It has generally been assumed that stress exposure leads to the development of psychosis and psychotic-like-experiences, however, reverse causality is also possible. Symptoms, personality traits and unusual behaviour may predispose individuals to stressful experiences [74]: for example, early symptoms may predispose an individual to bullying victimisation [25, 75, 76]. VR could be

embedded in longitudinal research to allow the repeated measurements of current stress response alongside repeated measure of life stressors and symptoms. This type of research could also help in defining resilience factors in people who experience adverse life events, but do not go on to develop psychosis [77, 78]. Previous studies have recognised the similarity between anomalous experiences and intrusive symptoms of post-traumatic stress disorder (PTSD) [79] as well as co-morbid PTSD in people with psychosis (e.g., [80]). While not the topic of the current review, is also worth mentioning that the search revealed a rich body of literature demonstrating the use of VR-assisted therapy for mental health problems (for recent reviews see [71, 81, 82]), and the use of VR in the research, assessment and treatment of psychosis [83, 84] and PTSD [48, 85–87] has previously been the topic of extensive reviews. Lessons can, therefore, be learned from those studies to inform future experimental studies into the psychological mechanisms of psychosis.

Further evidence for the fact that VR is an ecologically valid method to assess real-time responses to social environments was found by studies looking at the effect of manipulating population density and the ethnic density of avatars on the levels of paranoid ideation and physiological activation [65, 66, 69]. Of interest in the context of urbanicity is the finding that changes in the architectural characteristics of a room in VR can be used to change the physiological reaction to closed spaces: compared to participants in a VR room without windows, participants in the room with windows and a nice calming view had lower cortisol reactivity to stress and a quicker recovery after a stress induction test [88].

Using VR to investigate anomalous experiences opens up engaging avenues for further research in the association between paranoid ideation and the mechanisms which underlie the propensity to experience hallucinations, such as self-monitoring, attributional biases, and mental imagery (e.g., [89, 90]). VR studies can also contribute to research the differences between clinical and non-clinical samples of people who report anomalous experiences [91, 92].

Limitations

There were limitations regarding the ~~research conducted to date~~ reviewed studies.. The design of all studies was cross-sectional and life events were assessed retrospectively. A number of studies had small sample sizes and healthy control participants in some of the studies were exclusively from a student population, thus differences in social class, employment status, and level of education between people with psychosis and student samples may contribute to some of the differences found between groups. A final possible limitation is also

the fact that the reviewed studies were conducted by a relatively small group of researchers who collaborate with each other and who have a strong research interest in paranoid ideation. The focus of research on paranoid ideation reflects on one hand, the long tradition and large body of research of the role of psychological mechanisms in the aetiology of paranoid symptoms [93–96]. On the other hand, this strong focus on paranoid ideation also reflects the interest of the clinical academics who conducted the studies, with the large majority of reviewed studies conducted by Freeman and colleagues, or by researchers who collaborated with him. Future VR research should ensure that other areas of stress research are not neglected.

Conclusion

The reviewed studies suggest that VR can be used to investigate psychological processes and mechanisms associated with psychosis. The ability to assess real-time, cognitive, emotional, behavioural and physiological responses in a controlled but ecologically valid environment has enormous potential for future research in the mechanisms involved in the onset and maintenance of psychosis.

Acknowledgments

We acknowledge the NIHR Biomedical Research Centre for Mental Health at the South London and Maudsley NHS Foundation Trust and Institute of Psychiatry King's College London for their support.

References

1. Garety PA, Kuipers E, Fowler D, Freeman D, Bebbington PE (2001) A cognitive model of the positive symptoms of psychosis. *Psychol Med* 31(2):189–195
2. Garety PA, Bebbington P, Fowler D, Freeman D, Kuipers E (2007) Implications for neurobiological research of cognitive models of psychosis: a theoretical paper. *Psychol Med* 37(10):1377–1391. doi:10.1017/S003329170700013X
3. Freeman D, Garety PA, Kuipers E, Fowler D, Bebbington PE (2002) A cognitive model of persecutory delusions. *Br J Clin Psychol/Br Psychol Soc* 41(Pt 4):331–347
4. Bentall RP, Fernyhough C, Morrison AP, Lewis S, Corcoran R (2007) Prospects for a cognitive-developmental account of psychotic experiences. *Br*

J Clin Psychol/Br Psychol Soc 46(Pt 2):155–173.
doi:10.1348/014466506X123011

5. Morrison AP, Wells A (2003) A comparison of metacognitions in patients with hallucinations, delusions, panic disorder, and non-patient controls. *Behav Res Ther* 41(2):251–256
6. Van Dam DS, Van Nierop M, Viechtbauer W, Velthorst E, Van Winkel R, Bruggeman R, Cahn W, De Haan L, Kahn RS, Meijer CJ, Myin-Germeys I, Van Os J, Wiersma D (2015) Childhood abuse and neglect in relation to the presence and persistence of psychotic and depressive symptomatology. *Psychol Med* 45(7):1363–1377. doi:10.1017/S0033291714001561
7. Pariante CM (2015) Psychoneuroimmunology or immunopsychiatry? *Lancet Psychiatry* 2(3):197–199. doi:10.1016/S2215-0366(15)00042-5
8. Aiello G, Horowitz M, Hepgul N, Pariante CM, Mondelli V (2012) Stress abnormalities in individuals at risk for psychosis: a review of studies in subjects with familial risk or with “at risk” mental state. *Psychoneuroendocrinology* 37(10):1600–1613. doi:10.1016/j.psyneuen.2012.05.003
9. Walker E, Mittal V, Tessner K (2008) Stress and the hypothalamic pituitary adrenal axis in the developmental course of schizophrenia. *Annu Rev Clin Psychol* 4:189–216. doi:10.1146/annurev.clinpsy.4.022007.141248
10. Day FL, Valmaggia LR, Mondelli V, Papadopoulos A, Papadopoulos I, Pariante CM, McGuire P (2014) Blunted cortisol awakening response in people at ultra high risk of developing psychosis. *Schizophr Res* 158(1–3):25–31. doi:10.1016/j.schres.2014.06.041
11. Lazarus RS (1991) Progress on a cognitive-motivational-relational theory of emotion. *Am Psychol* 46(8):819–834
12. Corcoran C, Walker E, Huot R, Mittal V, Tessner K, Kestler L, Malaspina D (2003) The stress cascade and schizophrenia: etiology and onset. *Schizophr Bull* 29(4):671–692
13. Zubin J, Spring B (1977) Vulnerability—a new view of schizophrenia. *J Abnorm Psychol* 86(2):103–126
14. Lederbogen F, Kirsch P, Haddad L, Streit F, Tost H, Schuch P, Wust S,

Pruessner JC, Rietschel M, Deuschle M, Meyer-Lindenberg A (2011) City living and urban upbringing affect neural social stress processing in humans. *Nature* 474(7352):498–501.

<http://nature.com/nature/journal/v474/n7352/abs/nature10190-f1.2.html>
(supplementary-information)

15. Allardyce J, Gilmour H, Atkinson J, Rapson T, Bishop J, McCreadie RG (2005) Social fragmentation, deprivation and urbanicity: relation to first-admission rates for psychoses. *Br J Psychiatry* 187:401–4016

16. Vassos E, Pedersen CB, Murray RM, Collier DA, Lewis CM (2012) Meta-analysis of the association of urbanicity with schizophrenia. *Schizophr Bull* 38(6):1118–1123. doi:10.1093/schbul/sbs096

17. Morgan C, Hutchinson G (2009) The social determinants of psychosis in migrant and ethnic minority populations: a public health tragedy. *Psychol Med* 1:1–5. doi:10.1017/S0033291709005546

AQ5

18. Seeman MV (2011) Canada: psychosis in the immigrant Caribbean population. *Int J Soc Psychiatry* 57(5):462–470.
doi:10.1177/0020764010365979

19. Boydell J, van Os J, McKenzie K, Allardyce J, Goel R, McCreadie RG, Murray RM (2001) Incidence of schizophrenia in ethnic minorities in London: ecological study into interactions with environment. *BMJ* 323(7325):1336–1338

20. Das-Munshi J, Becares L, Boydell JE, Dewey ME, Morgan C, Stansfeld SA, Prince MJ (2012) Ethnic density as a buffer for psychotic experiences: findings from a national survey (EMPIRIC). *Br J Psychiatry* 201(4):282–290. doi:10.1192/bjp.bp.111.102376

21. Karlsen S, Nazroo JY, McKenzie K, Bhui K, Weich S (2005) Racism, psychosis and common mental disorder among ethnic minority groups in England. *Psychol Med* 35(12):1795–1803. doi:10.1017/S0033291705005830

22. Veling W, Selten JP, Susser E, Laan W, Mackenbach JP, Hoek HW (2007) Discrimination and the incidence of psychotic disorders among ethnic minorities in The Netherlands. *Int J Epidemiol* 36(4):761–768.
doi:10.1093/ije/dym085

23. Stowkowy J, Liu L, Cadenhead KS, Cannon TD, Cornblatt BA, McGlashan TH, Perkins DO, Seidman LJ, Tsuang MT, Walker EF, Woods SW, Bearden CE, Mathalon DH, Addington J (2016) Early traumatic experiences, perceived discrimination and conversion to psychosis in those at clinical high risk for psychosis. *Soc Psychiatry Psychiatric Epidemiol* 51:1–7. doi:10.1007/s00127-016-1182-y

AQ6

24. Lereya ST, Copeland WE, Costello EJ, Wolke D (2015) Adult mental health consequences of peer bullying and maltreatment in childhood: two cohorts in two countries. *Lancet Psychiatry* 2(6):524–531. doi:10.1016/S2215-0366(15)00165-0

25. Arseneault L, Cannon M, Fisher HL, Polanczyk G, Moffitt TE, Caspi A (2011) Childhood trauma and children's emerging psychotic symptoms: a genetically sensitive longitudinal cohort study. *Am J Psychiatry* 168(1):65–72. doi:10.1176/appi.ajp.2010.10040567

26. Lukoff D, Snyder K, Ventura J, Nuechterlein KH (1984) Life events, familial stress, and coping in the developmental course of schizophrenia. *Schizophr Bull* 10(2):258–292

27. Norman RMG, Malla AK (1993) Stressful life events and schizophrenia. II: Conceptual and methodological issues. *Br J Psychiatry* 162 (FEB.):166–174

28. Myin-Germeys I, Krabbendam L, Delespaul PAEG, Van Os J (2003) Do life events have their effect on psychosis by influencing the emotional reactivity to daily life stress? *Psychol Med* 33(2):327–333

29. Lee SY, Kim KR, Park JY, Park JS, Kim B, Kang JI, Lee E, An SK, Kwon JS (2011) Coping strategies and their relationship to psychopathologies in people at ultra high-risk for psychosis and with schizophrenia. *J Nerv Ment Dis* 199(2):106–110. doi:10.1097/NMD.0b013e3182083b96

30. Jalbrzikowski M, Sugar CA, Zinberg J, Bachman P, Cannon TD, Bearden CE (2014) Coping styles of individuals at clinical high risk for developing psychosis. *Early intervention in psychiatry* 8(1):68–76. doi:10.1111/eip.12005

31. Masillo A, Day F, Laing J, Howes O, Fusar-Poli P, Byrne M, Bhattacharyya S, Fiori Nastro P, Girardi P, McGuire PK, Valmaggia LR (2012) Interpersonal sensitivity in the at-risk mental state for psychosis.

Psychol Med 42(9):1835–1845. doi:10.1017/S0033291711002996

32. Boyce P, Parker G (1989) Development of a scale to measure interpersonal sensitivity. *Australasian Psychiatry* 23(3):341–351

33. Masillo A, Valmaggia LR, Saba R, Brandizzi M, Lindau JF, Solfanelli A, Curto M, Narilli F, Telesforo L, Kotzalidis GD, Di Pietro D, D'Alema M, Girardi P, Fiori Nastro P (2016) Interpersonal sensitivity and functioning impairment in youth at ultra-high risk for psychosis. *Eur Child Adolesc Psychiatry* 25(1):7–16. doi:10.1007/s00787-015-0692-6

34. Tiernan B, Tracey R, Shannon C (2014) Paranoia and self-concepts in psychosis: a systematic review of the literature. *Psychiatry Res* 216(3):303–313. doi:10.1016/j.psychres.2014.02.003

35. Lincoln TM, Mehl S, Ziegler M, Kesting ML, Exner C, Rief W (2010) Is fear of others linked to an uncertain sense of self? The relevance of self-worth, interpersonal self-concepts, and dysfunctional beliefs to paranoia. *Behav Ther* 41(2):187–197. doi:10.1016/j.beth.2009.02.004

36. van Os J, Linscott RJ, Myin-Germeys I, Delespaul P, Krabbendam L (2009) A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness-persistence-impairment model of psychotic disorder. *Psychol Med* 39(2):179–195. doi:10.1017/S0033291708003814

37. Kelleher I, Devlin N, Wigman JT, Kehoe A, Murtagh A, Fitzpatrick C, Cannon M (2014) Psychotic experiences in a mental health clinic sample: implications for suicidality, multimorbidity and functioning. *Psychol Med* 44(8):1615–1624. doi:10.1017/S0033291713002122

38. Kelleher I, Wigman JT, Harley M, O'Hanlon E, Coughlan H, Rawdon C, Murphy J, Power E, Higgins NM, Cannon M (2015) Psychotic experiences in the population: association with functioning and mental distress. *Schizophr Res* 165(1):9–14. doi:10.1016/j.schres.2015.03.020

39. Wigman JT, Devlin N, Kelleher I, Murtagh A, Harley M, Kehoe A, Fitzpatrick C, Cannon M (2014) Psychotic symptoms, functioning and coping in adolescents with mental illness. *BMC Psychiatry* 14:97. doi:10.1186/1471-244X-14-97

40. Fusar-Poli P, Borgwardt S, Bechdolf A, Addington J, Riecher-Rossler A,

Schultze-Lutter F, Keshavan M, Wood S, Ruhrmann S, Seidman LJ, Valmaggia L, Cannon T, Velthorst E, De Haan L, Cornblatt B, Bonoldi I, Birchwood M, McGlashan T, Carpenter W, McGorry P, Klosterkötter J, McGuire P, Yung A (2013) The psychosis high-risk state: a comprehensive state-of-the-art review. *JAMA Psychiatry* 70(1):107–120. doi:10.1001/jamapsychiatry.2013.269

41. Bentall RP, de Sousa P, Varese F, Wickham S, Sitko K, Haarmans M, Read J (2014) From adversity to psychosis: pathways and mechanisms from specific adversities to specific symptoms. *Soc Psychiatry Psychiatr Epidemiol* 49(7):1011–1022. doi:10.1007/s00127-014-0914-0

42. Freeman D, Garety PA (2000) Comments on the content of persecutory delusions: does the definition need clarification? *Br J Clin Psychol/Br Psychol Soc* 39(Pt 4):407–414

43. Freeman D, Pugh K, Antley A, Slater M, Bebbington P, Gittins M, Dunn G, Kuipers E, Fowler D, Garety P (2008) Virtual reality study of paranoid thinking in the general population. *Br J Psychiatry: J Ment Sci* 192(4):258–263. doi:10.1192/bjp.bp.107.044677

44. Garety PA, Everitt BS, Hemsley DR (1988) The characteristics of delusions: a cluster analysis of deluded subjects. *Eur Arch Psychiatry Neurol Sci* 237(2):112–114

45. Freeman D (2007) Suspicious minds: the psychology of persecutory delusions. *Clin Psychol Rev* 27(4):425–457. doi:10.1016/j.cpr.2006.10.004

46. Myin-Germeys I, Oorschot M, Collip D, Lataster J, Delespaul P, van Os J (2009) Experience sampling research in psychopathology: opening the black box of daily life. *Psychol Med* 39(9):1533–1547. doi:10.1017/S0033291708004947

47. Reininghaus U, Depp CA, Myin-Germeys I (2016) Ecological interventionist causal models in psychosis: targeting psychological mechanisms in daily life. *Schizophr Bull* 42(2):264–269. doi:10.1093/schbul/sbv193

48. Rizzo A, Buckwalter G, Forbell E, Reist C, Difede J, Rothbaum BO, Lange B, Koenig S, Talbot S (2013) Virtual reality applications to address the wounds of war. *Psychiatric Annals* 43(3):123–138. doi:10.3928/00485713-20130306-08

49. Slater M (2004) Presence and emotions. *Cyberpsychol Behav* 7 (1):121. doi:10.1089/109493104322820200 (**author reply 123**)
50. Parsons TD, Courtney CG, Arizmendi B, Dawson M (2011) Virtual reality stroop task for neurocognitive assessment. *Stud Health Technol Inform* 163:433–439
51. Seidel RJ, Chatelier PR (2013) Virtual reality, training's future?: perspectives on virtual reality and related emerging technologies, vol 6. Springer Science & Business Media
52. Rizzo A, Kim GJ (2005) A SWOT analysis of the field of virtual reality rehabilitation and therapy. *Presence, Teleoperators Virtual Environ* 14:119–146
53. Thomas BH, Ciliska D, Dobbins M, Micucci SA (2004) A Process for Systematically Reviewing the Literature: providing the Research Evidence for Public Health Nursing Interventions. *Worldviews Evid Based Nurs* 1(3):176–184
54. Freeman D (2008) Studying and treating schizophrenia using virtual reality: a new paradigm. *Schizophr Bull* 34(4):605–610
55. Valmaggia LR, Day FL, Kroll J, Laing J, Byrne M, Fusar-Poli P, McGuire P (2015) Bullying victimisation and paranoid ideation in people at ultra high risk for psychosis. *Schizophr Res* 168(1–2):68–73
56. Freeman D, Antley A, Ehlers A, Dunn G, Thompson C, Vorontsova N, Garety P, Kuipers E, Glucksman E, Slater M (2014) The use of immersive Virtual Reality (VR) to predict the occurrence 6 months later of paranoid thinking and posttraumatic stress symptoms assessed by self-report and interviewer methods: a study of individuals who have been physically assaulted. *Psychol Assess* 26(3):841–847
57. Freeman D, Thompson C, Vorontsova N, Dunn G, Carter LA, Garety P, Kuipers E, Slater M, Antley A, Glucksman E, Ehlers A (2013) Paranoia and post-traumatic stress disorder in the months after a physical assault: a longitudinal study examining shared and differential predictors. *Psychol Med* 43(12):2673–2684
58. Shaikh M, Ellett L, Dutt A, Day F, Laing K, Kroll J, Pterella S, McGuire P, Valmaggia L (2016) Perceived ethnic discrimination and persecutory paranoia in individuals at ultra-high risk for psychosis. *Psychiatry Res*

59. Valmaggia LR, Day F, Garety P, Freeman D, Antley A, Slater M, Swapp D, Myin-Germeys I, McGuire P (2015) Social defeat predicts paranoid appraisals in people at high risk for psychosis. *Schizophr Res* 168(1–2):16–22
60. Fornells-Ambrojo M, Elenbaas M, Barker C, Swapp D, Navarro X, Rovira A, Sanahuja JMT, Slater M (2016) Hypersensitivity to contingent behavior in paranoia: a new virtual reality paradigm. *J Nerv Ment Dis* 204(2):148–152
61. Freeman D, Slater M, Bebbington PE, Garety PA, Kuipers E, Fowler D, Met A, Read CM, Jordan J, Vinayagamoorthy V (2003) Can virtual reality be used to investigate persecutory ideation? *J Nerv Ment Dis* 191(8):509–514
62. Freeman D, Garety PA, Bebbington P, Slater M, Kuipers E, Fowler D, Green C, Jordan J, Ray K, Dunn G (2005) The psychology of persecutory ideation II: a virtual reality experimental study. *J Nerv Ment Dis* 193(5):309–315
63. Valmaggia LR, Freeman D, Green C, Garety P, Swapp D, Antley A, Prescott C, Fowler D, Kuipers E, Bebbington P, Slater M, Broome M, McGuire PK (2007) Virtual reality and paranoid ideations in people with an ‘at-risk mental state’ for psychosis. *Br J Psychiatry* 191(SUPPL. 51):s63–s68
64. Freeman D, Pugh K, Vorontsova N, Antley A, Slater M (2010) Testing the continuum of delusional beliefs: an experimental study using virtual reality. *J Abnorm Psychol* 119(1):83–92
65. Veling W, Brinkman WP, Dorrestijn E, van der Gaag M (2014) Virtual reality experiments linking social environment and psychosis: a pilot study. *Cyberpsychol Behav Soc Netw* 17(3):191–195
66. Veling W, Pot-Kolder R, Counotte J, van Os J, van der Gaag M (2016) Environmental Social Stress. A Virtual Reality Study. *Schizophr Bull*, Paranoia and Psychosis Liability. doi:10.1093/schbul/sbw031
67. Atherton S, Antley A, Evans N, Cernis E, Lister R, Dunn G, Slater M, Freeman D (2016) Self-confidence and paranoia: an experimental study using an immersive virtual reality social situation. *Behav Cogn Psychother* 44(1):56–64

68. Freeman D, Evans N, Lister R, Antley A, Dunn G, Slater M (2014) Height, social comparison, and paranoia: an immersive virtual reality experimental study. *Psychiatry Res* 218(3):348–352
69. Brinkman WP, Veling W, Dorrestijn E, Sandino G, Vakili V, van der Gaag M (2011) Virtual reality to study responses to social environmental stressors in individuals with and without psychosis. *Stud Health Technol Inf* 167:86–91
70. Freeman D, Gittins M, Pugh K, Antley A, Slater M, Dunn G (2008) What makes one person paranoid and another person anxious? The differential prediction of social anxiety and persecutory ideation in an experimental situation. *Psychol Med* 38(8):1121–1132
71. Goncalves R, Pedrozo AL, Coutinho ESF, Figueira I, Ventura P (2012) Efficacy of Virtual Reality Exposure Therapy in the Treatment of PTSD: A Systematic Review. *PLoS One* 7(12) (no pagination) **e48469**. [doi:10.1371/journal.pone.0048469](https://doi.org/10.1371/journal.pone.0048469)
72. Falconer CJ, Slater M, Rovira A, King JA, Gilbert P, Antley A, Brewin CR (2014) Embodying compassion: A virtual reality paradigm for overcoming excessive self-criticism. *PLoS One* 9(11) (no pagination) **e111933**. [doi:10.1371/journal.pone.0111933](https://doi.org/10.1371/journal.pone.0111933)
73. Falconer CJ, Rovira A, King JA, Gilbert P, Antley A, Fearon P, Ralph N, Slater M, Brewin CR (2016) Embodying self-compassion within virtual reality and its effects on patients with depression. *Br J Psychiatry Open* 2(1):74–80
74. Bebbington PE, Bhugra D, Brugha T, Singleton N, Farrell M, Jenkins R, Lewis G, Meltzer H (2004) Psychosis, victimisation and childhood disadvantage: evidence from the second British National Survey of Psychiatric Morbidity. *Br J Psychiatry* 185:220–226. [doi:10.1192/bjp.185.3.220](https://doi.org/10.1192/bjp.185.3.220)
75. Shakoor S, McGuire P, Cardno AG, Freeman D, Plomin R, Ronald A (2015) A shared genetic propensity underlies experiences of bullying victimization in late childhood and self-rated paranoid thinking in adolescence. *Schizophr Bull* 41(3):754–763. [doi:10.1093/schbul/sbu142](https://doi.org/10.1093/schbul/sbu142)
76. Goldman-Mellor S, Caspi A, Arseneault L, Ajala N, Ambler A, Danese A, Fisher H, Hucker A, Odgers C, Williams T, Wong C, Moffitt TE (2016) Committed to work but vulnerable: self-perceptions and mental health in

NEET 18-year olds from a contemporary British cohort. *J Child Psychol Psychiatry* 57(2):196–203. doi:10.1111/jcpp.12459

77. Brett CMC, Peters ER, McGuire PK (2015) Which psychotic experiences are associated with a need for clinical care? *Eur Psychiatry* 30(5):648–654. doi:10.1016/j.eurpsy.2014.12.005

78. Underwood R, Kumari V, Peters E (2016) Appraisals of psychotic experiences: an experimental investigation of symptomatic, remitted and non-need-for-care individuals. *Psychol Med* 46(6):1249–1263. doi:10.1017/S0033291715002780

79. Hardy A, Fowler D, Freeman D, Smith B, Steel C, Evans J, Garety P, Kuipers E, Bebbington P, Dunn G (2005) Trauma and hallucinatory experience in psychosis. *J Nerv Ment Dis* 193(8):501–507

80. Varese F, Smeets F, Drukker M, Lieveise R, Lataster T, Viechtbauer W, Read J, van Os J, Bentall RP (2012) Childhood adversities increase the risk of psychosis: a meta-analysis of patient-control, prospective- and cross-sectional cohort studies. *Schizophr Bull* 38(4):661–671. doi:10.1093/schbul/sbs050

81. Eichenberg C, Wolters C (2012) Virtual realities in the treatment of mental disorders: a review of the current state of research. In: *InTech*. doi:10.5772/50094

AQ8

82. Valmaggia LR, Latif L, Kempton MJ, Rus-Calafell M (2016) Virtual reality in the psychological treatment for mental health problems: a systematic review of recent evidence. *Psychiatry Res* 236:189–195

83. Veling W, Moritz S, van der Gaag M (2014) Brave new worlds-review and update on virtual reality assessment and treatment in psychosis. *Schizophr Bull* 40(6):1194–1197. doi:10.1093/schbul/sbu125

84. Macedo M, Marques A, Queiros C (2015) Virtual reality in assessment and treatment of schizophrenia: a systematic review. *Jornal Brasileiro de Psiquiatria* 64(1):70–81

85. Jiandani N, Nair SR, Shukla H (2014) Efficacy of virtual reality exposure therapy in the management of symptoms associated with post traumatic stress disorder. *Value in Health* 17(7):A572

86. Wiederhold BK, Wiederhold MD (2014) Virtual reality for posttraumatic stress disorder. *Advances in virtual reality and anxiety disorders* 211–233

AQ9

87. Rizzo A, Parsons TD, Lange B, Kenny P, Buckwalter JG, Rothbaum B, Difede J, Frazier J, Newman B, Williams J, Reger G (2011) Virtual reality goes to war: a brief review of the future of military behavioral healthcare. *J Clin Psychol Med Settings* 18(2):176–187

88. Fich LB, Jonsson P, Kirkegaard PH, Wallergard M, Garde AH, Hansen T (2014) Can architectural design alter the physiological reaction to psychosocial stress? A virtual TSST experiment. *Physiol Behav* 135:91–97

89. Johns LC, Kompus K, Connell M, Humpston C, Lincoln TM, Longden E, Preti A, Alderson-Day B, Badcock JC, Cella M, Fernyhough C, McCarthy-Jones S, Peters E, Raballo A, Scott J, Siddi S, Sommer IE, Laroi F (2014) Auditory verbal hallucinations in persons with and without a need for care. *Schizophr Bull* 40(SUPPL. 4):S255–S264. doi:10.1093/schbul/sbu005

90. Stinson K, Valmaggia LR, Antley A, Slater M, Freeman D (2010) Cognitive triggers of auditory hallucinations: an experimental investigation. *J Behav Ther Exp Psychiatry* 41(3):179–184. doi:10.1016/j.jbtep.2009.12.003

91. Brett C, Heriot-Maitland C, McGuire P, Peters E (2014) Predictors of distress associated with psychotic-like anomalous experiences in clinical and non-clinical populations. *Br J Clin Psychol* 53(2):213–227. doi:10.1111/bjc.12036

92. Ward TA, Gaynor KJ, Hunter MD, Woodruff PWR, Garety PA, Peters ER (2014) Appraisals and responses to experimental symptom analogues in clinical and nonclinical individuals with psychotic experiences. *Schizophr Bull* 40(4):845–855. doi:10.1093/schbul/sbt094

93. Freeman D, Garety P (2014) Advances in understanding and treating persecutory delusions: a review. *Soc Psychiatry Psychiatr Epidemiol* 49(8):1179–1189. doi:10.1007/s00127-014-0928-7

94. Howes OD, Murray RM (2014) Schizophrenia: an integrated sociodevelopmental-cognitive model. *Lancet* 383(9929):1677–1687. doi:10.1016/S0140-6736(13)62036-X

95. Myin-Germeys I, Delespaul P, van Os J (2005) Behavioural sensitization

to daily life stress in psychosis. *Psychol Med* 35(5):733–741

96. Reininghaus U, Kempton MJ, Valmaggia L, Craig TK, Garety P, Onyejiaka A, Gayer-Anderson C, So SH, Hubbard K, Beards S, Dazzan P, Pariante C, Mondelli V, Fisher HL, Mills JG, Viechtbauer W, McGuire P, van Os J, Murray RM, Wykes T, Myin-Germeys I, Morgan C (2016) Stress sensitivity, aberrant salience, and threat anticipation in early psychosis: an experience sampling study. *Schizophr Bull* 42(3):712–722.
doi:10.1093/schbul/sbv190