



City Research Online

City St George's, University of London

Citation: Baron, J., Hirani, S. P. & Newman, S. P. (2016). Challenges in Patient Recruitment, Implementation, and Fidelity in a Mobile Telehealth Study. *Telemedicine and e-Health*, 22(5), pp. 400-409. doi: 10.1089/tmj.2015.0095

This is the supplemental version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/16645/>

Link to published version: <https://doi.org/10.1089/tmj.2015.0095>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Table 3. Contacts recorded on the web server between intervention participants (n=40) and the technical support team

Technical / device-related operations [†]	Total	Mean, SD	Minimum-maximum	Completed/Expected (%)
Outgoing phone calls Technical assistance	10 ^Δ	0.25±0.87	0-7	-
Non-transmission	19	0.48±0.72	0-2	-
End of trial	32	NA*	NA	32/40 (80%)
Incoming phone calls	5	0.12± 0.33	0-1	
Text messages	29	0.73± 0.60	0-2	-

[†]This does not include contacts with the engineer, attempted but unsuccessful phone calls, and voicemails.

^ΔA further 33 outgoing phone calls were recorded on a separate log sheet provided by the technical support team

* NA: not applicable; participants could only receive 1 end of trial call.

Note: Incoming phone calls refer to instances when MTH participants initiated contact. In the last column, % of expected contacts are only provided for intervention components of a fixed nature, i.e. for which specific completion rates could be expected.

Table 4. Contacts between intervention participants (n=40) and the mobile telehealth (MTH) nurse

Contact type (n=40)	Total	Mean, SD	Minimum-maximum	Completed/Expected (%)
Clinical feedback	23	0.58± 0.78	0-3	-
Patient education and assistance				
Outgoing phone calls Introductory	37	NA*	NA	37/40 (92.5%)
6 weekly education	32	0.80 ± 0.93	0-3	32/240 (13.3%)
Incoming phone calls	7	0.18± 0.38	0-1	-
Text messages [‡]	164	4.10 ± 2.56	0-9	

* NA: not applicable; participants could only receive 1 introductory call.

Note: These data do not include attempted but unsuccessful phone calls, or voicemails. Clinical feedback calls refer to outgoing calls made by a MTH nurse that were not logged on the web server as educational or introductory calls. Data in the last column 'completed/expected' are only provided for intervention components that were to be delivered at a known frequency, i.e. for which specific completion rates could be expected based on the protocol.

[‡]A minority (5.7%) of these messages were considered to be characteristic of clinical feedback and advice (see text).