



City Research Online

City St George's, University of London

Citation: Kerr, O. (2014). Comment on 'Nonlinear eigenvalue problems'. Journal of Physics A: Mathematical and Theoretical, 47(36), 368001. doi: 10.1088/1751-8113/47/36/368001

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/17470/>

Link to published version: <https://doi.org/10.1088/1751-8113/47/36/368001>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Solutions to $y'(x) = \cos [\pi x^p y(x)^q]$

Oliver S. Kerr

*Department of Mathematics, City University London,
Northampton Square, London, EC1V 0HB, U.K.*

Abstract

The asymptotic behaviour of solutions to $y'(x) = \cos[\pi x^p y(x)^q]$ is considered. This is a generalisation of the problem of the behaviour of $y'(x) = \cos[\pi xy(x)]$ that was investigated by Bender, Fring and Komijani [1]. We present a derivation of the asymptotic results that follows the approach used in Kerr [2].

1 Introduction

In Bender, Fring and Komijani [1] a detailed asymptotic analysis of the nonlinear initial-value problem

$$y'(x) = \cos[\pi xy(x)], \quad y(0) = a \quad (1)$$

was presented which focused on the solutions for $x \geq 0$. They showed that for $a > 0$ solutions could be split into classes depending on the initial conditions such that solution with $a_{n-1} < a < a_n$ displayed an oscillatory region with n maxima before decaying monotonically to zero. They then found the result that as $n \rightarrow \infty$, $a_n \sim 2^{5/6} \sqrt{n}$. This result was subsequently derived using a different approach by Kerr [2]. Here we use this alternative derivation to obtain equivalent results to the generalisation of the original problem where we look for asymptotic solutions to

$$y'(x) = \cos [\pi x^p y(x)^q], \quad y(0) = a \quad (2)$$

where p and q are positive integers¹

Much of this derivation is essentially the same as that of Kerr [2]

2 Outline

The typical behaviour of solutions to (2) is essentially the same as the original problem and is shown by the solid lines in figure 1 (with $p = q = 1$). There is an initial oscillatory phase where the frequency increases and the amplitude decreases as the initial value, $y(0)$, increases. These oscillatory solutions drift downwards until they undergo a transition to monotonic decay towards the horizontal axis.

Some of the basic behaviour of the solutions of (2) can be understood by considering the lines in the x - y plane where $x^p y^q$ is constant. The situation is shown schematically




Fig. 1: Plots of solutions to (2) with $y(0) = 2, 4, 6, 8$ and $p = q = 1$. The dotted lines in $x/p > y/q$ show the curves $x^p y^q = C$ to which these converge asymptotically as $x \rightarrow \infty$. The dashed lines in $x/p < y/q$ give the estimate of the mean path of the oscillatory part of these curves.

in figure 2. The arguments here are essentially the same as those in section 3 of Kerr, except the region is now divided by the line $x/p = y/q$ and the lines under consideration are lines of the form $x^p y^q = c$.

If we consider lines where $x^p y^q = 2n$ then solutions will have gradient 1 where they intersect these lines, similarly when $x^p y^q = 2n+1$ they will intersect with gradient -1 , and when $x^p y^q = 2n \pm 1/2$ they will intersect with gradient 0. The gradients of the solutions will have gradients with magnitude at most 1, while the lines $x^p y^q = c$ for constants c have gradients greater than 1 in magnitude for $x/p < y/q$, and less than 1 for $x/p > y/q$. In the region $x/p < y/q$ solutions must cross the lines $x^p y^q = c$ from left to right, with a maximum each time it crosses a line $x^p y^q = 2n + \frac{1}{2}$, $n = 1, 2, 3, \dots$. In the region $x/p > y/q$ this restriction no longer holds. This results in the solutions having intrinsically different behaviour above and below the line $x/p = y/q$.

3 Solutions in the region $x/p > y/q$

This is essentially the same argument as previously in Kerr [2].

Any solution that enters a region $2n + \frac{1}{2} \leq x^p y^q \leq 2n+1$ is trapped in this region as x increases as the gradient of a solution on the lower boundary is 0, and on the upper boundary is -1 . Indeed, in such a region any solution that is initially above a

¹ Strictly speaking this requirement to be positive integers can be relaxed.




Fig. 2: Schematic plot for solutions in the region $x/p > y/q$ and the influences of the lines of form $x^p y^q = c$. The dotted lines show the trajectories of various solutions, while the dashed line shows the path of the separatrix dividing solutions that converge to $x^p y^q = 2n + 1/2$ from those that converge to $x^p y^q = 2n + 5/2$.

line $x^p y^q = 2n + \frac{1}{2} + \epsilon$ will have a negative gradient of magnitude greater than $\sin \pi \epsilon$ and so must eventually pass below $x^p y^q = 2n + \frac{1}{2} + \epsilon$, whose gradient tends to zero as $x \rightarrow \infty$. Thus all solutions in this region asymptote to the line $x^p y^q = 2n + \frac{1}{2}$.

All solutions in the region $2n - \frac{1}{2} < x^p y^q < 2n + \frac{1}{2}$ will have positive gradients and so will pass into the region $2n + \frac{1}{2} \leq x^p y^q \leq 2n + 1$ from below, and will have one maximum in the region $x/p > y/q$.

There is one solution in the region $2n + 1 < x^p y^q < 2n + \frac{3}{2}$ that stays in this region. Solutions initially below this curve will pass into the region $2n + \frac{1}{2} \leq x^p y^q \leq 2n + 1$ and remain there, while those above it will end up in the region $2n + \frac{5}{2} \leq x^p y^q \leq 2n + 3$. This curve is indicated by the dashed line in figure 2. By a similar argument to that given previously it can be shown that these seperatrices tend towards their asymptotes $x^p y^q = 2n + 3/2$ from below. We will denote the point where the separatrix crosses the line $x/p = y/q$ as $x/p = y/q = b_n$, and hence $2n + 1 < p^p q^q b_n^{p+q} < 2n + 3/2$, or $(2n + 1)^{1/(p+q)} / (p^p q^q)^{1/(p+q)} < b_n < (2n + 3/2)^{1/(p+q)} / (p^p q^q)^{1/(p+q)}$.

Clearly, any solution that ends up just above the curve $x^p y^q = 2n + \frac{1}{2}$ will have crossed n lines given by $x^p y^q = \frac{1}{2}, \frac{5}{2}, \frac{9}{2}, \dots$ and so will have n maxima. Hence any solution that crosses the line $x/p = y/q$ with $b_{n-1} < x/p = y/q < b_n$ will have n maxima.

4 Solutions in the region $x/p < y/q$

As before, for large values of $y(0)$ the solution $y(x)$ will tend to oscillate quickly with small amplitude. The previous arguments hold here.

If lines of constant $x^p y^q = c$ are given locally by the lines $x + \alpha y = C$ then the mean path of the oscillatory solution is given by

$$\frac{dy}{dx} = \frac{\sqrt{1 - \alpha^2} - 1}{\alpha}. \quad (3)$$

The value of α is determined by the curves $x^p y^q = c$. On such curves

$$\frac{dy}{dx} = -\frac{pc^{1/q}}{qx^{(p+q)/q}} = -\frac{py}{qx}. \quad (4)$$

Since the gradient of the lines $x + \alpha y = C$ is $-1/\alpha$, we find $\alpha = qx/py$ and so the equation for the slope of the average curve is given by

$$\frac{dy}{dx} = \sqrt{\left(\frac{py}{qx}\right)^2 - 1} - \frac{py}{qx}. \quad (5)$$

This has solutions

$$\left(\sqrt{p^2 y^2 - q^2 x^2} + py\right)^p \left((p+q)y - \sqrt{p^2 y^2 - q^2 x^2}\right)^{p+q} = 2^p p^p q^{p+q} y(0)^{2p+q}. \quad (6)$$

The solution curves meet the line $x/p = y/q$ at the point $x/p = y/q = \beta$ when

$$y(0) = \frac{(p+q)^{\frac{p+q}{2p+q}}}{2^{\frac{p}{2p+q}}} \beta \quad (7)$$

If the a_n are the values of $y(0)$ which correspond to the b_n , and so are solutions with n maxima then we would have here

$$a_n \approx \frac{(p+q)^{\frac{p+q}{2p+q}}}{2^{\frac{p}{2p+q}}} b_n \approx \frac{2^{\frac{1}{p+q}} (p+q)^{\frac{p+q}{2p+q}} n^{\frac{1}{p+q}}}{2^{\frac{p}{2p+q}} p^{\frac{p}{p+q}} q^{\frac{q}{p+q}}}. \quad (8)$$

5 Conclusions

We have used alternative derivation of Kerr [2] to analyse the generalisation the problem considered by Bender, Fring and Komijani [1]. With luck I've got the coefficients right.

References

- [1] Carl M. Bender, Andreas Fring, and Javad Komijani. Nonlinear eigenvalue problems. Accepted for publication in *J. Phys. A*, 2014.
- [2] Oliver S. Kerr. On “nonlinear eigenvalue problem”. Submitted for publication in *J. Phys. A*, 2014.