



City Research Online

City St George's, University of London

Citation: Walsh, B. & Laudicella, M. (2017). Disparities In Cancer Care And Costs At The End Of Life: Evidence From England's National Health Service. *Health Affairs*, 36(7), pp. 1218-1226. doi: 10.1377/hlthaff.2017.0167

This is the supplemental version of the paper.


This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://openaccess.city.ac.uk/id/eprint/17775/>

Link to published version: <https://doi.org/10.1377/hlthaff.2017.0167>


Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Exhibit A1: Average Number of Elective and Emergency Bed Days in Last Six Months: Partitioned by Deprivation Quintiles.




Source: (Authors' analysis)

Exhibit A2: Predicted Cost by Number of Comorbidities (2, 3-7, 8+) and SES Quintile



Source: (Authors' analysis)

Exhibit A3: Probability of an Elective, Emergency, and Outpatient Visit in Last Month of Life



Source: (Authors' analysis)

Exhibit A4: Top 5 Healthcare Resource Group (HRGs) in the Last 6 Months of Life

	Lowest SES Quintile	Quintile 2	Quintile 3	Quintile 4	Highest SES Quintile
Colorectal					
	Elective				
Same day Chemotherapy admission	28.07%	30.73%	31.31%	33.14%	30.72%
Malignant Large Intestinal Disorders LOS 0 days	12.50%	13.33%	12.62%	12.87%	14.75%
Single Plasma Exchange, Leucopheresis or Red Cell Exchange W LOS 2 days or less	5.99%	5.68%	6.03%	5.92%	6.15%
Vascular Access except for Renal Replacement Therapy W/O CC	4.37%	4.40%	5.23%	5.08%	5.42%
Malignant Large Intestinal Disorders W LOS 1 day or more W/O Major CC	N/A	3.43%	3.61%	2.95%	3.35%
	Emergency				
Malignant Large Intestinal Disorders W LOS 1 day or more W/O Major CC	7.95%	8.18%	8.10%	8.37%	8.39%
Malignant Large Intestinal Disorders W LOS 0 days	6.79%	6.65%	6.54%	6.28%	6.12%
Malignant Large Intestinal Disorders W LOS 1 day or more W Major CC	6.41%	6.85%	7.10%	7.07%	6.76%
Non-Malignant General Abdominal Disorders W LOS 0 days	3.30%	2.94%	3.12%	3.29%	3.13%
Malignant General Abdominal Disorders W LOS 1 day or more W/O Major CC	2.33%	2.42%	2.50%	2.38%	2.67%
Breast					
	Elective				
Same day Chemotherapy admission	40.29%	40.50%	40.88%	42.33%	40.14%
Malignant Breast Disorders W Intermediate CC	9.54%	9.10%	9.21%	10.28%	9.19%
Malignant Breast Disorders W/O CC	9.50%	9.05%	8.80%	7.57%	10.94%
Single Plasma Exchange, Leucopheresis or Red Cell Exchange W LOS 2 days or less and	5.46%	5.34%	6.07%	5.84%	5.04%

Chronic Kidney Disease W LOS 1 day or less associated W Renal Dialysis	5.57%	4.48%	N/A	N/A	N/A
	Emergency				
Malignant Breast Disorders W Major CC	9.58%	9.74%	9.11%	9.39%	9.21%
Malignant Breast Disorders W Intermediate CC	5.82%	5.66%	5.67%	5.43%	5.85%
Lobar, Atypical or Viral Pneumonia W Major CC	3.35%	3.26%	3.42%	4.09%	4.10%
Pleural Effusion W Major CC	3.63%	3.24%	3.11%	2.88%	2.71%
Brain Tumours or Cerebral Cysts W CC	2.61%	2.71%	2.49%	2.57%	2.90%
Prostate					
	Elective				
Same day Chemotherapy admission	13.33%	17.88%	16.88%	19.03%	20.65%
Single Plasma Exchange, Leucopheresis or Red Cell Exchange W LOS 2 days or less and	10.20%	13.97%	15.19%	15.93%	15.85%
Kidney, Urinary Tract and Prostate Neoplasms W LOS 1 day or less	14.77%	14.08%	13.61%	12.94%	15.73%
Chronic Kidney Disease W LOS 1 day or less associated W Renal Dialysis	19.85%	6.86%	13.00%	10.40%	3.62%
Bladder Intermediate Endoscopic Procedure	2.91%	3.44%	2.96%	N/A	N/A
	Emergency				
Kidney, Urinary Tract and Prostate Neoplasms W LOS 2 days or more W Intermediate CC	5.30%	5.30%	5.85%	5.76%	5.86%
Kidney, Urinary Tract and Prostate Neoplasms W LOS 1 day or less	5.16%	5.02%	5.06%	4.83%	4.65%
Kidney, Urinary Tract and Prostate Neoplasms W LOS 2 days or more W Major CC	3.77%	4.36%	4.34%	4.52%	4.68%
Lobar, Atypical or Viral Pneumonia W Major CC	4.39%	3.98%	4.15%	3.62%	3.54%
Non interventional acquired cardiac conditions	2.79%	2.79%	2.60%	2.31%	2.74%
Lung					
	Elective				
Same day Chemotherapy admission	34.09%	34.81%	35.81%	35.37%	34.51%
Fibre optic Bronchoscopy	11.81%	11.00%	10.78%	10.42%	10.26%
Respiratory Neoplasms W/O CC	8.88%	9.19%	9.03%	9.56%	9.79%
Respiratory Neoplasms W Major CC	6.20%	6.75%	6.42%	6.54%	7.55%
Single Plasma Exchange, Leucopheresis or Red Cell Exchange W LOS 2 days or less and	5.98%	6.01%	6.13%	6.62%	6.28%
	Emergency				

Respiratory Neoplasms W Major CC	20.50%	21.35%	21.40%	21.22%	21.28%
Respiratory Neoplasms W CC	8.00%	8.08%	8.32%	8.17%	8.18%
Lobar, Atypical or Viral Pneumonia W Major CC	5.70%	5.40%	5.11%	4.95%	4.50%
Respiratory Neoplasms W/O CC	2.65%	2.94%	3.28%	3.04%	3.79%
Non interventional acquired cardiac conditions	3.10%	2.91%	2.86%	2.72%	N/A

Source: (Authors' analysis)

Exhibit A5: Top 5 Healthcare Resource Group (HRGs) in the Last Month of Life

	Lowest SES Quintile	Quintile 2	Quintile 3	Quintile 4	Highest SES Quintile
Colorectal					
	Elective				
Same day Chemotherapy admission/attendance	16.55%	17.28%	18.08%	20.66%	18.38%
Malignant Large Intestinal Disorders W LOS 0 days	9.50%	9.00%	9.93%	10.13%	10.87%
Single Plasma Exchange, Leucophoresis or Red Cell Exchange W LOS 2 days or less	6.34%	7.08%	8.02%	6.82%	8.05%
Malignant Large Intestinal Disorders W LOS 1 day or more W/O Major CC	5.56%	6.68%	6.52%	5.42%	6.25%
Planned Procedures not carried out	N/A	3.91%	3.70%	N/A	3.28%
	Emergency				
Malignant Large Intestinal Disorders W LOS 1 day or more W Major CC	9.24%	9.36%	9.81%	9.96%	9.27%
Malignant Large Intestinal Disorders W LOS 1 day or more W/O Major CC	8.92%	9.41%	9.49%	9.78%	9.69%
Malignant Large Intestinal Disorders W LOS 0 days	8.73%	8.16%	8.39%	7.64%	7.56%
Non-Malignant General Abdominal Disorders W LOS 0 days	3.05%	2.57%	2.54%	2.80%	2.74%
Lobar, Atypical or Viral Pneumonia W Major CC	2.77%	2.81%	2.63%	2.41%	2.47%
Breast					
	Elective				
Same day Chemotherapy admission	29.11%	28.73%	26.75%	30.04%	28.67%
Malignant Breast Disorders W Intermediate CC	11.10%	9.82%	10.81%	10.36%	10.39%
Single Plasma Exchange, Leucophoresis or Red Cell Exchange W LOS 2 days or less	6.34%	7.53%	7.18%	6.98%	6.61%
Malignant Breast Disorders W/O CC	6.48%	7.33%	6.19%	7.58%	9.30%
Malignant Breast Disorders W Major CC	N/A	4.65%	6.31%	4.91%	4.81%
	Emergency				
Malignant Breast Disorders W Major CC	12.45%	12.57%	11.48%	11.83%	11.35%

Malignant Breast Disorders W Intermediate CC	6.57%	6.69%	6.75%	6.36%	6.52%
Lobar, Atypical or Viral Pneumonia W Major CC	5.24%	4.45%	4.44%	4.14%	3.90%
Pleural Effusion W Major CC	3.75%	3.14%	3.71%	4.20%	4.14%
Brain Tumours or Cerebral Cysts W CC	2.42%	2.56%	2.54%	2.93%	2.93%
Prostate					
	Elective				
Single Plasma Exchange, Leucophoresis or Red Cell Exchange W LOS 2 days or less	10.61%	15.88%	18.52%	19.81%	17.63%
Kidney, Urinary Tract and Prostate Neoplasms W LOS 1 day or less	13.57%	11.52%	9.74%	10.36%	13.15%
Same day Chemotherapy admission	9.41%	10.75%	9.59%	11.20%	10.56%
Chronic Kidney Disease W LOS 1 day or less associated W Renal Dialysis	18.27%	7.55%	8.71%	8.41%	N/A
Kidney, Urinary Tract and Prostate Neoplasms W LOS 1 day or less	4.60%	4.07%	4.80%	4.53%	4.94%
	Emergency				
Kidney, Urinary Tract and Prostate Neoplasms W LOS 1 day or less	5.48%	5.94%	6.28%	6.10%	6.79%
Lobar, Atypical or Viral Pneumonia W Major CC	6.43%	5.78%	6.11%	5.53%	5.34%
Kidney, Urinary Tract and Prostate Neoplasms W LOS 1 day or less	6.29%	6.16%	6.04%	5.75%	5.34%
Kidney, Urinary Tract and Prostate Neoplasms W LOS 2 days or more W Major CC	5.05%	6.02%	5.67%	6.18%	6.51%
Non-Transient Stroke or Cerebrovascular Accident, Nervous system infections or Encephalopathy W CC	N/A	N/A	2.22%	2.24%	2.32%
Lung					
	Elective				
Same day Chemotherapy admission	23.44%	22.49%	23.58%	22.76%	23.13%
Respiratory Neoplasms W Major CC	11.39%	12.73%	12.55%	12.38%	13.63%
Fibre optic Bronchoscopy	12.19%	12.20%	11.71%	10.25%	10.42%
Respiratory Neoplasms W/O CC	7.84%	9.28%	9.10%	9.55%	8.87%
Single Plasma Exchange, Leucophoresis or Red Cell Exchange W LOS 2 days or less	N/A	5.94%	5.95%	7.29%	6.60%
	Emergency				
Respiratory Neoplasms W Major CC	26.80%	28.15%	27.77%	27.40%	28.08%

Respiratory Neoplasms W CC	8.87%	8.86%	9.27%	9.04%	9.06%
Lobar, Atypical or Viral Pneumonia W Major CC	7.07%	6.75%	6.55%	6.32%	5.43%
Respiratory Neoplasms W/O CC	2.89%	3.26%	3.70%	3.26%	3.88%
Pleural Effusion W Major CC	N/A	N/A	2.58%	2.76%	2.80%

Source: (Authors' analysis)

Regressions from Exhibit 2

Colorectal

Variable	Cost of End of Life Care	
	Marginal Effect	Standard Error
Socioeconomic Group		
Lowest SES Quantile	Reference	Reference
2	-£507***	150
3	-£293*	151
4	-£486***	151
Highest SES Quantile	-£456***	156
Age at Diagnosis	£18	30
Age at Diagnosis Squared	-£1***	0
Days from Diagnosis to Death	-£5***	0
Days from Diagnosis to Death Squared	£0***	0
Sex		
Male	Reference	Reference
Female	£216**	93
Year of Diagnosis		
2001	Reference	Reference
2002	£795	2414
2003	£839	2393
2004	£645	2389
2005	£828	2389
2006	£1,885	2390
2007	£1,763	2391
2008	£1,648	2392
2009	£1,859	2393
2010	£1,584	2394
Weighted Charlson Score		
Score = 2	Reference	Reference
Score = 3	£5,887***	206
Score = 4	£7,425***	337
Score = 5	£9,100***	555
Score = 6	£3,282***	191
Score = 7	£3,704***	338
Score = 8	£3,845***	111
Score = 9	£4,630***	175
Score = 10	£6,298***	331

Score = 11	£7,469***	571
Score = 12	£8,106***	1084
Score = 13	£6,866***	2133
Score = 14	£6,953*	3749
Score = 15	£3,324	3933
Region		
London	Reference	Reference
North East	-£1,357***	233
North West	-£1,510***	187
Yorkshire & Humber	-£745***	201
East Midlands	-£1,412***	209
West Midlands	-£906***	197
East of England	-£2,475***	195
South East	-£2,303***	210
South Central	-£1,860***	225
South West	-£1,854***	199
Log Likelihood	-672,785	
Observations	66,061	

Source: (Authors' analysis)

Breast

Variable	Cost of End of Life Care	
	Marginal Effect	Standard Error
Socioeconomic Group		
Lowest SES Quantile	Reference	Reference
2	£41	191
3	-£40**2	187
4	-£512***	188
Highest SES Quantile	-£526***	194
Age at Diagnosis	£10	29
Age at Diagnosis Squared	£0	0
Days from Diagnosis to Death	-£1***	0
Days from Diagnosis to Death Squared	£0***	0
Year of Diagnosis		
2001	Reference	Reference
2002	£126	1528
2003	£249	1510
2004	£905	1510
2005	£1,006	1510
2006	£1,672	1512
2007	£1,601	1516
2008	£1,577	1520
2009	£1,488	1525
2010	£1,696	1538
Weighted Charlson Score		
Score = 2	Reference	Reference
Score = 3	£5,714***	275
Score = 4	£5,968***	402
Score = 5	£7,098***	686
Score = 6	£4,720***	227
Score = 7	£5,538***	427
Score = 8	£5,803***	162
Score = 9	£6,464***	287
Score = 10	£7,359***	542
Score = 11	£9,184***	945
Score = 12	£4,887***	1309
Score = 13	£4,666**	2126
Score = 14	£3,136	5918
Score = 15	£3,715	6344

Region		
London	Reference	Reference
North East	-£1,368	302
North West	-£1,105	233
Yorkshire & Humber	-£1,355	248
East Midlands	-£1,599	261
West Midlands	-£721	246
East of England	-£2,393	242
South East	-£2,194	260
South Central	-£1,977	272
South West	-£1,104	245
Log Likelihood	-365,877	
Observations	36,698	

Source: (Authors' analysis)

Prostate

Variable	Cost of End of Life Care	
	Marginal Effect	Standard Error
Socioeconomic Group		
Lowest SES Quantile	Reference	Reference
2	-£320	197
3	-£208	196
4	-£373*	195
Highest SES Quantile	-£564***	201
Age at Diagnosis	-£171**	73
Age at Diagnosis Squared	£1**	0
Days from Diagnosis to Death	-£2***	0
Days from Diagnosis to Death Squared	£0***	0
Year of Diagnosis		
2001	Reference	Reference
2002	£1,276	2193
2003	£1,597	2177
2004	£1,657	2175
2005	£1,821	2175
2006	£2,237	2176
2007	£2,341	2179
2008	£2,592	2182
2009	£2,376	2185
2010	£2,020	2193
Weighted Charlson Score		
Score = 2	Reference	Reference
Score = 3	£6,256***	223
Score = 4	£7,724***	269
Score = 5	£7,986***	350
Score = 6	£6,742***	365
Score = 7	£6,741***	590
Score = 8	£5,754***	173
Score = 9	£6,506***	216
Score = 10	£7,880***	272
Score = 11	£8,641***	386
Score = 12	£8,693***	656
Score = 13	£7,577***	1187
Score = 14	£9,106***	2861
Score = 15	£10,313**	4320

Region		
London	Reference	Reference
North East	-£1,238***	312
North West	-£1,120***	244
Yorkshire & Humber	-£1,169***	258
East Midlands	-£1,386***	264
West Midlands	-£821***	256
East of England	-£2,131***	245
South East	-£2,299***	271
South Central	-£1,745***	281
South West	-£1,179***	247
Log Likelihood	-393,424	
Observations	39,329	

Source: (Authors' analysis)

Lung

Variable	Cost of End of Life Care	
	Marginal Effect	Standard Error
Socioeconomic Group		
Lowest SES Quantile	Reference	Reference
2	£13	72
3	£87	76
4	-£81	79
Highest SES Quantile	£60	86
Age at Diagnosis	-£181***	23
Age at Diagnosis Squared	£1***	0
Days from Diagnosis to Death	-£3***	0
Days from Diagnosis to Death Squared	£0***	0
Sex		
Male	Reference	Reference
Female	£34	50
Year of Diagnosis		
2001-2005	Reference	Reference
2006	£1,022***	131
2007	£758***	129
2008	£772***	131
2009	£1,130***	135
2010	£910***	140
Weighted Charlson Score		
Score = 2	Reference	Reference
Score = 3	£3,624***	79
Score = 4	£4,707***	140
Score = 5	£6,047***	262
Score = 6	£4,107***	165
Score = 7	£4,559***	254
Score = 8	£4,302***	66
Score = 9	£4,952***	90
Score = 10	£5,571***	165
Score = 11	£6,481***	313
Score = 12	£7,437***	585
Score = 13	£5,347***	971
Score = 14	£10,055***	2615
Score = 15	£5,448	4899
Region		

London	Reference	Reference
North East	-£983***	116
North West	-£974***	99
Yorkshire & Humber	-£1,124***	103
East Midlands	-£1,036***	113
West Midlands	-£896***	109
East of England	-£1,675***	109
South East	-£2,334***	119
South Central	-£1,149***	127
South West	-£1,249***	113
Log Likelihood	-1,163,832	
Observations	116,749	

Source: (Authors' analysis)

Regressions from Exhibit 3

Colorectal

	Probability of Elective		Cost of Elective Care	
Variable	Marginal Effect	Standard Error	Marginal Effect	Standard Error
Socioeconomic Group				
Lowest SES Quantile	Reference	Reference	Reference	Reference
2	0.031***	0.006	-£280	189
3	0.050***	0.006	£72	192
4	0.061***	0.006	£111	192
Highest SES Quantile	0.072***	0.006	£182	200
Age at Diagnosis	0.016***	0.001	£135***	37
Age at Diagnosis Squared	0.000***	0.000	-£1***	0
Days from Diagnosis to Death	0.000***	0.000	£0	1
Days from Diagnosis to Death Squared	0.000**	0.000	£0**	0
Sex				
Male	Reference	Reference	Reference	Reference
Female	-0.044***	0.004	-£145	119
Year of Diagnosis				
2001	Reference	Reference	Reference	Reference
2002	0.025	0.118	£2,232	3294
2003	0.041	0.117	£1,958	3236
2004	0.045	0.117	£1,750	3227
2005	0.020	0.117	£1,744	3225
2006	0.044	0.117	£2,386	3225
2007	0.047	0.117	£2,461	3226
2008	0.047	0.117	£2,058	3227
2009	0.054	0.117	£2,157	3228
2010	0.040	0.117	£1,987	3229
Weighted Charlson Score				
Score = 2	Reference	Reference	Reference	Reference
Score = 3	0.139***	0.007	£2,685***	295
Score = 4	0.145***	0.010	£4,305***	487
Score = 5	0.143***	0.015	£5,893***	819
Score = 6	0.166***	0.008	£190	243
Score = 7	0.095***	0.013	£358	435
Score = 8	0.250***	0.005	-£56	150
Score = 9	0.191***	0.007	£131	199
Score = 10	0.175***	0.011	£1,389***	355
Score = 11	0.170***	0.017	£2,803***	641
Score = 12	0.175***	0.030	£3,566***	1280
Score = 13	0.231***	0.064	£4,468*	2689
Score = 14	0.245**	0.113	£1,501	3396
Score = 15	0.277*	0.154	-£1,155	3208

Region				
London	Reference	Reference	Reference	Reference
North East	-0.003	0.009	- £1,096***	293
North West	-0.052***	0.007	-£687***	240
Yorkshire & Humber	0.003	0.008	-£491*	253
East Midlands	-0.019**	0.008	- £1,375***	264
West Midlands	-0.002	0.008	-£379	247
East of England	-0.041***	0.008	- £1,043***	250
South East	-0.027***	0.008	- £1,007***	268
South Central	-0.048***	0.009	- £1,146***	290
South West	-0.035***	0.008	-£984***	255
Pseudo R2	0.113			
Prob > LR Chi2	0.000			
Observations	66,061		27,564	

Source: (Authors' analysis)

Breast

	Probability of Elective		Cost of Elective Care	
Variable	Marginal Effect	Standard Error	Marginal Effect	Standard Error
Socioeconomic Group				
Lowest SES Quantile	Reference	Reference	Reference	Reference
2	0.017***	0.007	£468*	282
3	0.027***	0.007	-£251	268
4	0.046***	0.007	£29	271
Highest SES Quantile	0.029***	0.007	£55	283
Age at Diagnosis	0.009***	0.001	-£43	42
Age at Diagnosis Squared	0.000***	0.000	£0	0
Days from Diagnosis to Death	0.000	0.000	£0	1
Days from Diagnosis to Death Squared	0.000	0.000	£0	0
Year of Diagnosis				
2001	Reference	Reference	Reference	Reference
2002	-0.023	0.067	-£886	2632
2003	-0.024	0.066	-£627	2612
2004	-0.010	0.066	-£50	2613
2005	-0.018	0.066	£134	2613
2006	0.003	0.066	£169	2616
2007	-0.017	0.066	£142	2622
2008	0.000	0.066	£329	2627
2009	0.007	0.066	£197	2634
2010	0.003	0.067	£821	2660
Weighted Charlson Score				
Score = 2	Reference	Reference	Reference	Reference
Score = 3	0.111***	0.010	£2,425***	514
Score = 4	0.098***	0.014	£1,516**	664
Score = 5	0.092***	0.021	£7,031***	1795
Score = 6	0.219***	0.009	£566*	289
Score = 7	0.144***	0.014	£339	486
Score = 8	0.309***	0.007	£903***	222
Score = 9	0.259***	0.010	£1,415***	330
Score = 10	0.219***	0.016	£2,037***	599
Score = 11	0.197***	0.024	£6,703	1418
Score = 12	0.150***	0.047	-£911	1227
Score = 13	0.079	0.074	-£604	2316

Score = 14	0.208	0.309	£3,404	12866
Score = 15	0.522**	0.223	-£3,952**	1872
Region				
London	Reference	Reference	Reference	Reference
North East	-0.027**	0.011	-£932**	432
North West	-0.075***	0.009	£415	343
Yorkshire & Humber	-0.029***	0.009	- £1,710***	354
East Midlands	-0.039***	0.010	- £1,878***	373
West Midlands	0.002	0.009	-£655*	346
East of England	-0.055***	0.009	- £1,29***6	351
South East	-0.041***	0.010	- £1,257***	381
South Central	-0.049***	0.010	- £1,328***	391
South West	-0.010	0.009	-£132	344
Pseudo R2	0.197			
Prob > LR Chi2	0.000			
Observations	36,698		27,564	

Source: (Authors' analysis)

Prostate

	Probability of Elective		Cost of Elective Care	
Variable	Marginal Effect	Standard Error	Marginal Effect	Standard Error
Socioeconomic Group				
Lowest SES Quantile	Reference	Reference	Reference	Reference
2	0.018**	0.008	-£549*	308
3	0.040***	0.008	-£142	302
4	0.044***	0.008	£1	300
Highest SES Quantile	0.051***	0.008	-£5	307
Age at Diagnosis	0.005*	0.003	-£46	96
Age at Diagnosis Squared	0.000***	0.000	£0	1
Days from Diagnosis to Death	0.000***	0.000	£0	1
Days from Diagnosis to Death Squared	0.000***	0.000	£0	0
Year of Diagnosis				
2001	Reference	Reference	Reference	Reference
2002	0.083	0.097	£3,973	4207
2003	0.095	0.096	£4,124	4182
2004	0.107	0.096	£3,722	4177
2005	0.119	0.096	£4,088	4176
2006	0.130	0.096	£4,469	4179
2007	0.135	0.096	£4,358	4182
2008	0.133	0.096	£4,584	4185
2009	0.132	0.096	£3,890	4188
2010	0.105	0.096	£3,143	4198
Weighted Charlson Score				
Score = 2	Reference	Reference	Reference	Reference
Score = 3	0.100***	0.008	£1,229***	361
Score = 4	0.099***	0.010	£3,077***	441
Score = 5	0.095***	0.014	£2,771***	585
Score = 6	0.128***	0.014	£1,417**	547
Score = 7	0.103***	0.023	£273	898
Score = 8	0.214***	0.006	-£427*	251
Score = 9	0.153***	0.008	-£333	312
Score = 10	0.167***	0.010	£1,238***	379
Score = 11	0.134***	0.014	£2,559***	565
Score = 12	0.114***	0.025	£1,331	1006
Score = 13	0.135***	0.045	-£1,637	1712

Score = 14	0.197*	0.106	£1,301	3609
Score = 15	0.034	0.167	-£8,623	7203
Region				
London	Reference	Reference	Reference	Reference
North East	0.000	0.012	-£696	472
North West	0.015	0.010	-£375	369
Yorkshire & Humber	-0.026**	0.010	- £1,345***	404
East Midlands	0.000	0.010	- £1,262***	403
West Midlands	0.026***	0.010	£235	387
East of England	-0.011	0.010	-£665*	376
South East	-0.019*	0.011	-£1,031**	421
South Central	-0.015	0.011	-£394	431
South West	0.023**	0.010	-£153	372
Pseudo R2	0.081			
Prob > LR Chi2	0.000			
Observations	39,329		13,615	

Source: (Authors' analysis)

Lung

	Probability of Elective		Cost of Elective Care	
Variable	Marginal Effect	Standard Error	Marginal Effect	Standard Error
Socioeconomic Group				
Lowest SES Quantile	Reference	Reference	Reference	Reference
2	0.025***	0.004	£256***	97
3	0.041***	0.004	£377***	103
4	0.054***	0.004	£196*	104
Highest SES Quantile	0.066***	0.005	£292**	114
Age at Diagnosis	0.022***	0.001	£11	31
Age at Diagnosis Squared	0.000***	0.000	£0*	0
Days from Diagnosis to Death	0.000***	0.000	£5***	0
Days from Diagnosis to Death Squared	0.000***	0.000	£0***	0
Sex				
Male	Reference	Reference	Reference	Reference
Female	-0.043***	0.003	-£166**	68
Year of Diagnosis				
2001-2005	Reference	Reference	Reference	Reference
2006	0.048***	0.009	£792***	195
2007	0.054***	0.009	£778***	193
2008	0.047***	0.009	£764***	194
2009	0.050***	0.009	£955***	198
2010	0.066***	0.009	£628***	203
Weighted Charlson Score				
Score = 2	Reference	Reference	Reference	Reference
Score = 3	0.078***	0.005	£1,187***	111
Score = 4	0.057***	0.007	£1,979***	205
Score = 5	0.040***	0.011	£3,785***	455
Score = 6	0.093***	0.009	£1,586***	221
Score = 7	0.009	0.012	£1,614***	364
Score = 8	0.170***	0.004	£1,241***	82
Score = 9	0.131***	0.005	£1,262***	105
Score = 10	0.107***	0.007	£1,519***	183
Score = 11	0.115***	0.013	£2,799***	394
Score = 12	0.072***	0.022	£5,156***	989
Score = 13	0.066	0.043	£106	825
Score = 14	0.013*	0.079	£15,468*	8,338

Score = 15	-0.047	0.200	£8,791	14,304
Region				
London	Reference	Reference	Reference	Reference
North East	-0.004	0.006	-£960***	157
North West	-0.012**	0.006	£0	134
Yorkshire & Humber	-0.027***	0.006	- £1,085***	142
East Midlands	-0.006	0.006	-£797***	151
West Midlands	0.020***	0.006	-£308**	144
East of England	-0.034***	0.006	-£734***	148
South East	-0.041***	0.007	- £1,448***	164
South Central	-0.009	0.007	-£259	170
South West	-0.006	0.006	-£462***	151
Pseudo R2	0.073			
Prob > LR Chi2	0.000			
Observations	116,749		52,122	

Source: (Authors' analysis)

Regressions from Exhibit 4

Colorectal

	Probability of Emergency		Cost of Emergency Care	
Variable	Marginal Effect	Standard Error	Marginal Effect	Standard Error
Socioeconomic Group				
Lowest SES Quantile	Reference	Reference	Reference	Reference
2	-0.025***	0.005	-£506***	115
3	-0.036***	0.005	-£563***	115
4	-0.049***	0.005	-£754***	116
Highest SES Quantile	-0.061***	0.005	-£693***	121
Age at Diagnosis	-0.014***	0.001	-£63***	23
Age at Diagnosis Squared	0.000***	0.000	£0***	0
Days from Diagnosis to Death	0.000***	0.000	-£3***	0
Days from Diagnosis to Death Squared	0.000***	0.000	£0	0
Sex				
Male	Reference	Reference	Reference	Reference
Female	0.018***	0.003	£618***	72
Year of Diagnosis				
2001	Reference	Reference	Reference	Reference
2002	-0.066	0.110	-£820	2360
2003	-0.040	0.109	-£659	2345
2004	-0.014	0.109	-£1,290	2342
2005	-0.010	0.109	-£1,025	2343
2006	0.008	0.109	-£709	2344
2007	0.008	0.109	-£982	2344
2008	0.020	0.109	-£1,068	2345
2009	0.017	0.109	-£827	2346
2010	0.018	0.109	-£987	2347
Weighted Charlson Score				
Score = 2	Reference	Reference	Reference	Reference
Score = 3	0.290***	0.006	£1,426***	153
Score = 4	0.328***	0.008	£1,914***	221
Score = 5	0.347***	0.011	£2,396***	331
Score = 6	0.314***	0.006	-£852***	138
Score = 7	0.378***	0.008	-£506***	220
Score = 8	0.335***	0.004	-£841***	94

Score = 9	0.380***	0.005	£28	126
Score = 10	0.397***	0.006	£862***	206
Score = 11	0.407***	0.009	£1,185***	327
Score = 12	0.417***	0.014	£1,408	596
Score = 13	0.333***	0.045	-£780	1015
Score = 14	0.290**	0.090	£2,642	2695
Score = 15	0.348**	0.107	£47	2772
Region				
London	Reference	Reference	Reference	Reference
North East	-0.002	0.008	- £1,022***	175
North West	-0.017**	0.006	-£851***	141
Yorkshire & Humber	-0.004	0.007	-£780***	152
East Midlands	0.005	0.007	-£962***	157
West Midlands	-0.026***	0.007	-£912***	150
East of England	-0.050***	0.007	- £1,304***	150
South East	-0.020***	0.007	- £1,589***	161
South Central	-0.015**	0.008	- £1,171***	173
South West	-0.038***	0.007	- £1,385***	152

Source: (Authors' analysis)

Breast

	Probability of Emergency		Cost of Emergency Care	
Variable	Marginal Effect	Standard Error	Marginal Effect	Standard Error
Socioeconomic Group				
Lowest SES Quantile	Reference	Reference	Reference	Reference
2	-0.020***	0.006	-£161	139
3	-0.025***	0.006	-£468***	137
4	-0.040***	0.006	-£501***	138
Highest SES Quantile	-0.034***	0.007	-£726***	141
Age at Diagnosis	-0.002**	0.001	£56***	22
Age at Diagnosis Squared	0.000***	0.000	£0	0
Days from Diagnosis to Death	0.000**	0.000	-£1*	0
Days from Diagnosis to Death Squared	0.000***	0.000	£0**	0
Year of Diagnosis				
2001	Reference	Reference	Reference	Reference
2002	-0.025	0.060	£1,307	1108
2003	-0.012	0.059	£1,172	1087
2004	0.002	0.059	£1,391	1085
2005	0.022	0.059	£1,180	1084
2006	0.031	0.060	£1,477	1085
2007	0.026	0.060	£1,697	1088
2008	0.024	0.060	£1,476	1090
2009	0.020	0.060	£1,373	1093
2010	0.011	0.060	£1,596	1103
Weighted Charlson Score				
Score = 2	Reference	Reference	Reference	Reference
Score = 3	0.449***	0.007	£1,140***	172
Score = 4	0.483***	0.007	£1,532***	244
Score = 5	0.481***	0.010	£1,595***	371
Score = 6	0.411***	0.007	£70	150
Score = 7	0.470***	0.008	£1,121***	259
Score = 8	0.424***	0.006	£304**	118
Score = 9	0.464***	0.006	£871***	168
Score = 10	0.483***	0.008	£1,753***	299
Score = 11	0.478***	0.011	£1,333***	424
Score = 12	0.484***	0.020	£680	760
Score = 13	0.486***	0.032	£1,489	1401

Score = 14	0.449***	0.007	-£1,984	2554
Score = 15	0.483***	0.007	-£151	3612
Region				
London	Reference	Reference	Reference	Reference
North East	-0.013	0.011	-£776***	217
North West	-0.009	0.008	-£705***	167
Yorkshire & Humber	-0.008	0.009	-£790***	178
East Midlands	0.001	0.009	-£783***	187
West Midlands	-0.018**	0.009	-£701***	178
East of England	-0.046***	0.008	- £1,200***	176
South East	-0.032***	0.009	- £1,314***	191
South Central	-0.033***	0.009	-£939***	200
South West	-0.054***	0.008	- £1,039***	179
Pseudo R2	0.216			
Prob > LR Chi2	0.000			
Observations	36,698		13,067	

Source: (Authors' analysis)

Prostate

	Probability of Emergency		Cost of Emergency Care	
Variable	Marginal Effect	Standard Error	Marginal Effect	Standard Error
Socioeconomic Group				
Lowest SES Quantile	Reference	Reference	Reference	Reference
2	-0.010	0.006	-£291**	141
3	-0.021***	0.006	-£318**	141
4	-0.027***	0.006	-£569***	141
Highest SES Quantile	-0.032***	0.006	-£701***	145
Age at Diagnosis	-0.011***	0.002	£47	54
Age at Diagnosis Squared	0.000***	0.000	£0	0
Days from Diagnosis to Death	0.000***	0.000	-£1	0
Days from Diagnosis to Death Squared	0.000***	0.000	£0	0
Year of Diagnosis				
2001	Reference	Reference	Reference	Reference
2002	0.038	0.060	-£617	1643
2003	0.037	0.060	-£355	1631
2004	0.056	0.060	-£251	1629
2005	0.046	0.060	-£240	1629
2006	0.049	0.060	-£151	1629
2007	0.053	0.060	-£55	1632
2008	0.052	0.060	£194	1634
2009	0.050	0.060	£207	1636
2010	0.039	0.061	£309	1641
Weighted Charlson Score				
Score = 2	Reference	Reference	Reference	Reference
Score = 3	0.323***	0.007	£1,325***	155
Score = 4	0.367***	0.010	£2,129***	181
Score = 5	0.383***	0.015	£2,455***	233
Score = 6	0.313***	0.012	£1,458***	249
Score = 7	0.331***	0.021	£1,924***	398
Score = 8	0.292***	0.004	£587***	127
Score = 9	0.377***	0.007	£1,407***	150
Score = 10	0.405***	0.011	£2,141***	182
Score = 11	0.413***	0.017	£2,680***	255
Score = 12	0.491***	0.044	£3,083***	429
Score = 13	0.490***	0.080	£2,772***	775

Score = 14	-	-	£3,019	1857
Score = 15	-	-	£6,125**	2803
Region				
London	Reference	Reference	Reference	Reference
North East	-0.011	0.010	- £1,191***	223
North West	-0.011	0.008	- £1,080***	176
Yorkshire & Humber	0.006	0.008	- £1,037***	186
East Midlands	-0.007	0.008	- £1,295***	190
West Midlands	-0.017**	0.008	- £1,291***	186
East of England	-0.042***	0.007	- £1,648***	178
South East	-0.023***	0.008	- £1,785***	197
South Central	-0.021**	0.008	- £1,521***	205
South West	-0.038***	0.007	- £1,815***	179
Pseudo R2	0.266			
Prob > LR Chi2	0.000		-309,771	
Observations	39,329		31,192	

Source: (Authors' analysis)

Lung

	Probability of	Cost of Emergency
--	----------------	-------------------

	Emergency		Care	
Variable	Marginal Effect	Standard Error	Marginal Effect	Standard Error
Socioeconomic Group				
Lowest SES Quantile	Reference	Reference	Reference	Reference
2	-0.013***	0.003	-£178***	55
3	-0.017***	0.003	-£259***	58
4	-0.024***	0.003	-£434***	59
Highest SES Quantile	-0.033***	0.004	-£333***	65
Age at Diagnosis	-0.011***	0.001	-£142***	17
Age at Diagnosis Squared	0.000***	0.000	£1***	0
Days from Diagnosis to Death	0.000***	0.000	-£2***	0
Days from Diagnosis to Death Squared	0.000***	0.000	£0***	0
Sex				
Male	Reference	Reference	Reference	Reference
Female	0.001	0.002	£359***	38
Year of Diagnosis				
2001-2005	Reference	Reference	Reference	Reference
2006	0.005	0.005	£306***	112
2007	-0.002	0.005	-£89	110
2008	0.005	0.005	-£120	111
2009	0.006	0.005	£138	114
2010	-0.002	0.006	£29	118
Weighted Charlson Score				
Score = 2	Reference	Reference	Reference	Reference
Score = 3	0.338***	0.003	£841***	59
Score = 4	0.358***	0.004	£1,586***	95
Score = 5	0.369***	0.005	£2,187***	164
Score = 6	0.302***	0.006	£1,235***	115
Score = 7	0.364***	0.006	£1,622***	169
Score = 8	0.338***	0.003	£755***	50
Score = 9	0.370***	0.003	£1,452***	63
Score = 10	0.386***	0.004	£1,924***	106
Score = 11	0.388***	0.005	£2,039***	183
Score = 12	0.386***	0.008	£2,364***	327
Score = 13	0.373***	0.019	£2,554***	667
Score = 14	-	-	£2,369**	1188
Score = 15	-	-	-£1,905	1467
Region				

London	Reference	Reference	Reference	Reference
North East	-0.018***	0.005	-£400	86
North West	-0.037***	0.004	-£797	74
Yorkshire & Humber	-0.011**	0.004	-£511	77
East Midlands	-0.025***	0.005	-£509	84
West Midlands	-0.027***	0.004	-£756	81
East of England	-0.045***	0.004	-£793	82
South East	-0.019***	0.005	-£1,356	89
South Central	-0.025***	0.005	-£892	96
South West	-0.049***	0.005	-£931	85
Psuedo R2	0.204			
Prob > LR Chi2	0.000			
Observations	116,749		95,427	

Source: (Authors' analysis)