



City Research Online

City St George's, University of London

Citation: Pereira, L., Owen-Smith, N. & Moleon, M. (2013). Facultative predation and scavenging by mammalian carnivores: seasonal, regional and intra-guild comparisons. *Mammal review*, 44(1), pp. 44-55. doi: 10.1111/mam.12005

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/19463/>

Link to published version: <https://doi.org/10.1111/mam.12005>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

1 **Facultative predation and scavenging by mammalian**

2 **carnivores: seasonal, regional and intra-guild**

3 **comparisons**

4

5 **ABSTRACT**

6 **1.** The extent to which vertebrate carnivores shift facultatively between predation and
7 scavenging has recently been emphasized. In this article we propose as a null
8 foundation that all carnivores have to do is wait until animals eventually succumb to
9 the debilitating effects of advancing age. However, this may be insufficient,
10 recognizing the intense competition among other scavengers and decomposers for
11 the rich food. Moreover, the availability of carcasses of animals dying from causes
12 besides predation varies seasonally, so carnivores must be adapted to exploit other
13 sources of food in order to survive through the seasonal cycle.

14 **2.** We explore how mammalian carnivores cope with seasonality in carrion supply
15 and prey vulnerability to predation. We focus mainly on large carnivores and
16 ungulates, and we compare ecological communities living in northern temperate and
17 African savanna ecosystems.

18 **3.** When carrion is scarce, carnivores can (i) take advantage of temporarily
19 vulnerable segments of prey populations, such as newborn young, heavily pregnant
20 females and males distracted or debilitated by reproductive activities, (ii) switch to
21 carcass remains left by or stolen from other carnivores, or (iii) exploit small animals
22 and non-animal food sources.

23 **4.** Relationships between carnivores tending towards predation or scavenging can be
24 both competitive and facilitatory. For instance, top-carnivores can provide a quite
25 constant supply of carcasses throughout the year, which crucially subsidizes
26 scavengers in the seasons when carrion availability by other causes is low.

Alterations of seasonal patterns due to human-caused environmental change may enhance the role of top-carnivores as buffers of anthropogenic perturbations of natural processes.

5. Megaherbivores, which are not normally regarded as prey but can indeed provide huge carrion subsidies, may strongly influence the future interspecific interactions between carnivores and the proportion of food flowing towards scavenging relative to predation.

Keywords: African savannas, carrion supply, competition, facilitation, global warming, megaherbivore, northern temperate ecosystems, prey vulnerability, seasonality

Running head: Predation versus scavenging in carnivores

Word count: 8636

Please cite as: Pereira, L.M., Owen-Smith, N. and Moleon, M. (2013) Facultative predation and scavenging by mammalian carnivores: seasonal, regional and intra-guild comparisons. *Mammal Review*, DOI: 10.1111/mam.12005.

INTRODUCTION

All animals die eventually, and their carcasses provide a food source for scavengers. By definition, scavengers feed on the remains (i.e. carrion) of animals that have died, while predators kill animals in order to feed on the resulting carcasses. Nevertheless, vertebrates traditionally labelled predators usually do not pass by opportunities to consume carcasses when encountered (DeVault et al. 2003, Selva et al. 2005, Wilson & Wolkovich 2011); indeed, what we find in nature is a quite continuous gradient from the most genuine predators that rarely eat dead animals (e.g. cheetah *Acinonyx jubatus*) to those mostly consuming carrion (e.g. brown hyaena *Hyaena brunnea*; e.g. Skinner & Chimimba 2005). Therefore, the conventional distinction between predators and scavengers may be misleading with regard to both the inter-specific interactions and the energy fluxes taking place within food webs.

Schaller (1972) distinguished four ways whereby carnivores secure their food: (1) Additive predation (i.e. killing healthy animals); (2) Compensatory predation (i.e. preying on young, old or sick animals vulnerable to mortality from other causes, thereby promoting the survival of the remainder of the prey population); (3) Kleptoparasitism (i.e. displacing other carnivores from their kills); (4) Scavenging (i.e. feeding solely on animals that have died from disease, malnutrition or other causes, or on the remains of predator kills). Individual carnivores can exhibit several or even all of these feeding patterns, which shows that there may be a continuum between predatory versus scavenging strategies not only at the inter- but also at the intra-specific level.

In this article we propose as a null foundation that all carnivores have to do to feed is wait until animals eventually succumb to the debilitating effects of advancing age. In practice this is untenable, recognizing the intense competition among other scavengers and decomposers for the rich food constituted by animal body tissues (Wilson & Wolkovich 2011). Hence, helping animals die a little sooner can gain prior access to the meat. However, the energy costs associated with hunting are high,

74 especially for large mammals; for instance, transport costs for the largest carnivores
75 (lions –*Panthera leo*– and polar bears –*Ursus maritimus*–) are 2-3 times higher than
76 expected for their size (Carbone et al. 2007). The challenge that carnivores face is
77 how to secure a regular supply of food when the susceptibility of animals to die or be
78 killed varies seasonally and spatially. As a result, most large mammalian carnivores
79 shift facultatively between hunting and scavenging depending on seasonal and
80 regional variation in the susceptibility of their prey to mortality.

81 Seasonal variation in the availability of food is a central theme in the ecology
82 of herbivores (Owen-Smith 2002). However, its role in carnivory has not been
83 adequately addressed by previous reviews (Cortés-Avizanda et al. 2009, DeVault et
84 al. 2011, Wilson & Wolkovich 2011), and there are important consequences for intra-
85 guild relationships among carnivores as well as for biomass pathways through food
86 webs. Seasonal variation in high northern latitudes is governed by temperature
87 variation and snow accumulation, whereas in more tropical latitudes rainfall and
88 hence surface water availability is the dominant influence. Seasonal variation in the
89 vulnerability of their prey to being killed has implications for the conservation of
90 carnivores in human-modified ecosystems. Those perceived as predators are largely
91 eliminated in order to protect domestic animals (e.g. Gittleman et al. 2001), and
92 human hunting can substantially modify carrion subsidies (e.g. Wilmers et al. 2003).

93 Recent reviews addressing overlaps between scavenging and hunting have
94 been orientated towards northern ecosystems, which retain somewhat impoverished
95 remnants of their former carnivore complement (Jędrzejewski et al. 1993, DeVault et
96 al. 2003, Cortés-Avizanda et al. 2009). Those addressing African ecosystems with
97 largely intact carnivore assemblages have emphasised predation and its
98 consequences, rather than the scavenging that also takes place (Sinclair et al. 2003,
99 Hayward 2006). We aim to balance this oversight by considering how large
100 mammalian carnivores cope with seasonal variation in food availability through
101 flexibility in their feeding strategies from strict predation to mainly scavenging. In

particular, we address the following questions: (1) How do carnivores cope with the seasonal period when prey are least vulnerable to mortality?; (2) How does facultative scavenging or predation affect competitive or facilitative relationships among carnivores?; (3) How do these patterns differ between depauperate northern communities and intact African ones?; (4) How do human activities alter trophic interactions and fluxes through predation versus scavenging?

SEASONAL VARIATION IN PREY MORTALITY AND VULNERABILITY TO PREDATION: CARNIVORE RESPONSES

Seasonal variation in food availability for carnivores may be generated in various ways. In the absence of predators, almost all deaths among adult ungulates in high northern latitudes occur during the winter months, when food shortages may lead directly to death or compromise the ability of these animals to withstand thermal stress (Clutton-Brock & Albon 1982, Coulson et al. 2001). The ability of ungulates to escape predation is also hampered by deep winter snow, benefitting especially cursorial predators like grey wolves (Post et al. 1999, Jędrzejewski et al. 2002, Hebblewhite 2005) and coyotes (*C. latrans*; Bekoff & Wells 1981, Huegel & Rongstad 1985). In the Białowieża Forest in Poland, 25-75% of discovered carcasses of the various ungulate species had apparently died from causes other than predation, despite the rich assemblage of predators there (Jędrzejewski et al. 1993). Although the frozen meat may present an obstacle for some carnivores (Haynes 1982, Selva et al. 2003), wolverines (*Gulo gulo*) take advantage of the refrigeration provided by high northern winters by amassing caches of carrion to augment opportunistically found carcasses (Inman et al. 2012a). Frozen carcasses may become accessible to other carnivores in a narrow pulse during the spring thaw. Northern ungulates are generally in worst condition during the transition from winter into spring, when adult elk can make up the majority of the diet of grey wolves (Metz et al. 2012).

In African savannas, deaths of herbivores attributed to causes other than predation are concentrated towards the end of the dry season when food is limiting (Mduma et al. 1999, Owen-Smith 2008). In Tanzania's Serengeti National Park, 64% of deaths of wildebeest (*Connochaetes taurinus*) occurred during the course of the dry season extending through July-December when the shortage of green grass became accentuated (Mduma et al. 1999). Body size influences the relative susceptibility of ungulates to starvation rather than predation, but with regional variation evident. In Serengeti, 75-90% of deaths of ungulates in the size range 170-250 kg was ascribed to predation, while only 20% of the mortality among buffalo (*Syncerus caffer*) and giraffe (*Giraffa camelopardis*) was due to predation (Sinclair et al. 2003). In contrast, in South Africa's Kruger National Park, 94% of found carcasses of all species weighing less than 1000 kg were attributed to predator kills, and uneaten carcasses were recorded only during disease outbreaks or severe droughts (Owen-Smith & Mills 2008). A proportion of the carcasses generated during disease outbreaks and droughts or extreme winter weather may remain uneaten, apparently because they become desiccated and lose the water content that most carnivores depend on from their food, especially under tropical African conditions (Gasaway et al. 1991, Cooper et al. 1999).

Migratory movements also affect the patterns of both ungulate mortality and vulnerability to predation. In Serengeti, deaths not mediated by predation are amplified by the huge local concentrations of migrants, saturating the capacity of resident predators to exploit them. Spotted hyenas (*Crocuta crocuta*) commute long distances to reach places where these ungulates are concentrated (Hofer & East 1993). Hence, migratory wildebeest, zebra (*Equus quagga*) and Thompson's gazelle (*Eudorcas thomsonii*) form the majority of spotted hyena's diet in the Serengeti plains despite only being available there for 26% of the year (Hofer & East 1993). When these migratory ungulates reach the Kenia's Masai Mara region during the dry season, they greatly increase the supply of carrion there. The desiccated carcasses

help support carnivores after the migrants have left (Cooper et al. 1999). In Africa, buffalo become most vulnerable to predation on adults when at their weakest during the transition period around the start of the rains (Owen-Smith 2008).

Prey vulnerability, dependent on age and body condition, is especially important for cursorial predators like wolves (*Canis lupus*), spotted hyenas, wild dogs (*Lycaon pictus*) and cheetahs, which capture their prey through pursuits limited in the duration that they can be sustained (FitzGibbon & Fanshawe 1989, Huggard 1994, Pole et al. 2004). On the other hand, ambush hunters like most felids, which rely on stalking to within attack distance, are less affected by seasonal variation in the vulnerability of their prey. For example, cougars (*Puma concolor*) kill mule deer (*Odocoileus hemionus*) adults in better condition than those captured by grey wolves, although both species are selective towards calves and old individuals when hunting larger prey like elk (*Cervus canadensis*; Husseman et al. 2003). African lions also more successfully catch weakened prey (Sinclair & Arcese 1995).

In spring or the early wet season, vulnerable newborn animals are generally produced, and become a food source for more omnivorous carnivores. Brown (*U. arctos*) and black bears (*U. americanus*) switch to hunting newly born caribou (*Rangifer tarandus*), elk and moose calves, albeit for only a few weeks (Gasaway et al. 1992, Mahoney & Schaefer 2002, Swenson et al. 2007, Barber-Meyer et al. 2008). Red foxes (*Vulpes vulpes*) cause much mortality among newborn ungulates from roe deer (*Capreolus capreolus*; Jarnemo et al. 2004) to reindeer, and wolverines also become predators on elk in this vulnerable life history stage (Tveraa et al. 2003, Gustine et al. 2006). Grey wolves concentrate their hunting on elk calves during summer and autumn and kill newborn bison (*Bos bison*) in early spring (Metz et al. 2012). Likewise, African carnivores such as lions and even jackals (*C. mesomelas*) take advantage of opportunities to run down juvenile ungulates that are more easily caught than the adult ungulates that they normally seek (Mills & Shenk 1992, Owen-Smith & Mason 2005, Owen-Smith 2008).

Male and female ungulates also show seasonal variation in their relative susceptibility to predation (Owen-Smith 2008). Females that are heavily pregnant are less capable of evading predation, and especially vulnerable during parturition (Molinari-Jobin et al. 2004). Male ungulates compete for mating rights while in prime condition in autumn after the end of the growing season, thereby exposing themselves to heightened risks of predation both through injuries sustained in fights and by isolating themselves from the security of the herd (FitzGibbon 1990a). Furthermore, the energy invested in rutting behavior leaves these males in a weakened state after the end of the mating period and hence more vulnerable to predators (FitzGibbon 1990b). Over the annual cycle, male ungulates invariably incur higher mortality rates than adult females (Pole et al. 2004, Owen-Smith et al. 2005, Owen-Smith 2008, Grange et al. 2012).

In summary, for predators the lean time of the year is towards the end of the summer or the rains, after calves have become robust and food remains plentiful (Fig. 1). Then carnivores have to turn their attention to alternative prey that, despite their lower meat yield, are more readily available. During summer, coyotes and foxes commonly hunt hares or mice (Bekoff & Wells 1981, Nellis & Keith 1976, Hamlin et al. 1984, Patterson et al. 1998), while wolves shift from moose towards beavers (*Caster canadensis*) on Isle Royale (Peterson & Page 1988), and towards white-tailed (*O. virginianus*) or mule deer in place of elk in Yellowstone National Park and elsewhere (Metz et al. 2012). Bears concentrate on pulses of fruit, small mammals or fish (Watts & Jonkel 1988), and even jackals may include a considerable proportion of fruits in their diet (Atkinson et al. 2002). During the wet season, lions favour zebras and wildebeest more than other prey species (Owen-Smith 2008), aided by concealing tall grass. In general, smaller prey species tend to be targeted by predators during the season of lesser vulnerability for herbivores.

Seasonality thus plays an important role in the feeding strategies of carnivores because it determines both the availability of carrion that can be exploited

as well as the vulnerability of prey species to being killed. This in turn has implications for intra-guild dynamics among carnivores, which is the topic of the next section.

COMPETITION AND FACILITATION AMONGST CARNIVORES IN SEASONAL ENVIRONMENTS

Predators both facilitate scavengers by making the remains of their prey carcasses available, and compete with them by killing and consuming animals that would eventually have died and nourished the scavengers. Furthermore, carnivores towards both ends of the predator–scavenger continuum compete aggressively by killing other carnivores when opportunities present themselves (Palomares & Caro 1999, Caro & Stoner 2003), and by stealing prey where possible through superiority in size or numbers (Cooper 1990, Honer et al. 2002). Creel et al. (2001) distinguish five forms of competition between carnivores, including active avoidance, avoidance leading to habitat shifts, dietary overlap, kleptoparasitism and killing. In more complex ecosystems, relationships become more complicated because by depressing the abundance of smaller carnivores, larger carnivores release populations of even smaller carnivores that were previously in competition with the middle sized carnivores (DeVault et al. 2011, Levi & Wilmers 2012). Nevertheless, facilitation remains the least well-documented relationship between carnivores (Steinmetz et al. 2008). What are the consequences for the coexistence of these species in seasonally variable environments?

In temperate ecosystems of North America, grey wolves have displaced coyotes as the primary predators on ungulates through their superior body and group size (Smith et al. 2003). Nevertheless, coyotes benefit from the remains of kills left by wolves, with this carrion particularly important in winters with low snow accumulation when few deer die of starvation (Wilmers & Getz 2004, 2005, Wilmers & Post 2006). North American wolves rarely scavenge (Metz et al. 2012). They leave little meat on

carcasses of the mainly young elk that they kill during summer, but more in winter when their prey selection shifts towards larger animals (Metz et al. 2012). During winter, coyotes benefit from these carcasses (Paquet 1992, Gese et al. 1996, Crabtree & Sheldon 1999, Wilmers & Getz 2004, 2005, Sala 2006). Thus top predators play a facilitative role in providing carcasses for other carnivores to scavenge outside of the season when carrion is abundant (Wilmers & Getz 2005, van Dijk et al. 2008). However, in Europe wolves commonly feed on carrion (e.g. Cuesta et al. 1991, Śmietana & Klimek 1993, Selva et al. 2003, Valdmann et al. 2005), depleting the food available to other scavengers. Nevertheless, in southern Scandinavia wolverines have benefitted from the recolonisation of by wolves through scavenging on kills of ungulates as large as moose (van Dijk et al. 2008). In Białowieża, wolves were the only vertebrate able to open deeply frozen European bison (*Bison bonasus*) carcasses, thus increasing scavenging opportunities for other species (Selva et al. 2003).

Competitive interactions among carnivores are strongly governed by relative body size. By limiting the abundance of coyotes through their size superiority, grey wolves allow foxes to thrive (Levi & Wilmers 2012). A comparable situation exists in Serengeti where locally high densities of lions and spotted hyenas are associated with a low density of cheetahs, but high densities of two jackal species (*C. mesomelas*, *C. aureus*; Durant et al. 2011). In the absence of wolves, coyotes kill more adult deer, benefitting from the higher proportion of aging adults and vulnerable offspring in these populations in these circumstances (Huegel & Rongstad 1985, Gese & Grothe 1995, Crabtree & Sheldon 1999, Prugh 2005). Under these conditions, coyotes outcompete and limit foxes that are likewise dependent mainly on small mammals during summer (Levi & Wilmers 2012).

In Africa, carrion is less readily available during wet season conditions when very few animals die of malnutrition. Furthermore, killing newly born calves provides less food than the carcasses of adult ungulates. At this time of the year, scavengers

depend more strongly on the carcass remains of animals killed by more capable predators (Henschel & Skinner 1990). Pursuit predators that undertake little or no scavenging, like cheetahs and wild dogs, are also disadvantaged during the wet season. Apart from juvenile prey, they must seek those few adults that can be overtaken because of injuries or advancing age (FitzGibbon & Fanshawe 1989). The paucity of this prey base may explain why both cheetahs and wild dogs cover vast home ranges at low density (Mills & Funston 2003). Both of these predators, the least powerful within the African large carnivore guild, also commonly lose animals they have killed to spotted hyenas and lions, and hence avoid regions where these larger species are abundant (Creel & Creel 1996, Durant 1998, Mills & Gorman 1997, Mills et al. 2004). However, large packs of wild dogs may hold one or two hyenas at bay while they rapidly devour meat from the kills they have made (Fanshawe & FitzGibbon 1993). Furthermore, cheetahs and wild dogs usually hunt during the day, thereby restricting kleptoparasitism by other predators that are mainly active nocturnally. Both species also incur direct mortality through being killed by lions and spotted hyenas, and have their distribution and abundance restricted as a result (Creel & Creel 1996, Durant 1998, Mills & Gorman 1997, Mills et al. 2004). Leopards also kill cheetahs sometimes (Palomares & Caro 1999), and commonly eat jackals and other small carnivores (Kruuk 1972, Schaller 1972).

The two African top carnivores (lions and spotted hyenas) interact intensely. Lions often kill hyenas, and hyenas can kill lions, especially as cubs (Palomares & Caro 1999). Lions and spotted hyenas generally overlap quite substantially in the ungulate species they hunt, but with regional variation (Hayward 2006, Owen-Smith 2008, Owen-Smith & Mills 2008). Spotted hyenas hunt both zebra and wildebeest in Ngorongoro Crater where lions frequently scavenge on hyena kills (Kruuk 1972), but seldom do so in the Kruger Park (Owen-Smith & Mills 2008). Lions listen for the cackling sounds of hyenas during the night to locate carcasses of animals killed by hyenas (Houston 1974, Schaller 1972). While it is commonly believed that lions are

social to increase their hunting success, strength in numbers is also important in defending their kills against the packs formed by hyenas, and vice versa (Cooper 1991, Honer et al. 2002). The relative density of lions versus hyenas appears to reflect the relative amount of hunting versus scavenging these competitors undertake, dependent on habitat conditions (Table 1). Spotted hyenas typically obtain one third or more of their food by scavenging in regions where their abundance is 1.5-2X that of lions, but in open grasslands they become mostly hunters and outnumber lions by a factor of four or more. Lions scavenge more where hyenas are more abundant. Scavenging from hyena kills helps lion prides survive in habitats deficient in the cover they need to stalk effectively, while hyenas persist in bushy areas where they cannot easily pursue prey over long distances by stealing carcasses from leopards (*P. pardus*; Pienaar 1969, Kruuk 1972, Bailey 1993). Another crucial adaptation of spotted hyenas is their capacity to subsist on the bones of carcasses left by lions. This ensures that these hyenas will always find some food, no matter how strong the predominance of lions at carcasses.

Distinctions in locomotory adaptations influence the extent to which large mammalian carnivores concentrate on found carcasses versus own kills as a food resource. Assisted by their loping gait, spotted hyenas can commute distances of 40-60 km between their dens and foraging areas, to get to places where prey species happen to be concentrated (Hofer & East 1993, 1995). To locate sufficient carcasses, the brown hyena, perhaps the best example of an obligate mammalian scavenger, wander on average 31 km and up to 54 km nightly, and may obtain over 90% of its food from animals found dead or the remains of predator kills (Mills 1990). They also commonly forage along shorelines where fish or seal carcasses wash up, earning the local name “strandwolf” (or “beach wolf”). In Serengeti, lions walked for on average only two hours per day, and no more than five hours even when actively hunting, thus covering at most 20 km (Schaller 1972). Wolverines are the prime scavengers in high northern latitudes, ranging widely to locate carcasses of animals that have died

in snow (Inman et al. 2012b). In summer they switch to becoming predators on newborn reindeer and caribou, as do foxes (Tveraa et al. 2003).

Mammalian carnivores compete for carcasses also with other vertebrate scavengers, invertebrates and decomposers. The Old World and New World vultures are the only obligate scavengers among vertebrates, exploiting their ability to soar widely, and thereby locate carcasses during times when few animals die, except through the agency of a predator (Ruxton & Houston 2004). The extremely efficient searching strategies of vultures, together with crucial physiological and immunological adaptations to counteract the potential lethal effects of toxins and pathogens (Houston & Cooper 1975), make them supremely adapted to exploit carrion (Houston 1979). Nevertheless, spotted hyenas can locate carcasses of animals that have died during the night before vultures become active, and hyenas and lions watch vultures during the day to find carcasses (Kruuk 1972, Schaller 1972, Houston 1979). Moreover, once they discover a carcass, large carnivores are able to displace vultures from the carcass, unless the vultures are present in very high numbers (Kruuk 1972). Mammalian carnivores that commonly scavenge seem adapted to deal with the toxins and pathogens that build up in carcasses with putrefaction, including the microparasites that might have caused the death of the animal (Cooper et al. 1999). While lions become infected with bovine tuberculosis from feeding on the buffalo they kill in the Kruger Park, and may suffer early death as a result, spotted hyenas feeding on these same carcasses seem unaffected, and appear immune to anthrax (Bengis et al. 2003).

CARRION SUBSIDY FROM MEGAHERBIVORE CARCASSES

Megaherbivores like elephants (*Loxodonta africana*), hippopotami (*Hippopotamus amphibius*) and rhinoceroses (*Ceratotherium simum* and *Diceros bicornis*) are usually excluded from the prey base supporting predators (Sinclair et al. 2003, Owen-Smith & Mills 2008). Adults of these species are too large to be killed even by

lions, although immature animals remain vulnerable (Joubert 2006). Nevertheless, adults eventually die, and their carcasses become available for consumption by various carnivores. Animals of these species coming to the end of their potential lifespan would have been born 40-60 years ago, at a time when their populations were still recovering from the remnants left by human hunters earlier in the 20th century. This situation is now changing in many parks, and as the population structure stabilizes with lowered recruitment, death from old age will make an increasing contribution to mortality. How might this contribution to the carcasses available to carnivores affect the abundance of predators, and the balance between hunting and scavenging strategies?

Given a life expectancy of 60-70 years for elephants, and 40-50 years for hippos and rhinos, 2-3% of the adult segment would be expected to die annually once a stationary age distribution prevails, even if zero mortality occurred prior to the end of the lifespan. Megaherbivores typically constitute 40-70% of the herbivore biomass in intact African ecosystems (Owen-Smith 1988). Hence the food supply for carnivores scavenging opportunistically on these carcasses could be elevated by a third or more, resulting in higher carnivore densities and potentially more predation on smaller ungulates. An increase in predation risk for other herbivores by this proportion could tip their population trend from stable to decreasing.

However, megaherbivores succumbing to malnutrition die mainly during the late dry season and largely near surface water (Conybeare & Haynes 1984). This seasonal and spatial restriction limits the effective contribution of this food to supporting carnivore populations. Moreover, the time needed to consume such large carcasses makes them subject to greater putrefaction, restricting their utilization by predators less adapted to tolerate toxins and pathogens. Whatever the situation, the pulse of such easily available food could perhaps divert predation away from other prey species during this period.

CONCLUSIONS

In this review we have highlighted the seasonal variation that exists in the availability of food for carnivores, affecting competitive versus facilitative relationships along the continuum from predation to scavenging. We have outlined the more complex relationships that develop in the species-rich mammalian assemblages that persist in African savanna regions, compared with the more impoverished communities that remain in North America and Europe. Nevertheless, humans have had a substantial effect, even in Africa, quite recently by suppressing the contribution that megaherbivores formerly made to carcass production through decimating the populations of these beasts. Further effects continue today as humans eliminate the large predators that sometimes kill livestock and restrict the number of animals that die of old age.

The food supply for carnivores depends not only on the seasonal cycle of plentiful and sparse food for herbivores dependent on temperature and precipitation, but also on the local distribution, age structure and body condition of prey populations. From a simplistic perspective, carnivores need not be killers, because herbivores die eventually through reaching the end of the lifespan. One of the major problems, however, is that these deaths are likely to be seasonally pulsed. For carnivores to survive through the annual cycle, they must be able to secure sufficient food year-round. Hence, during wet season or summer conditions, carnivores turn to temporarily vulnerable segments of prey populations, including newly born young, heavily pregnant females, and males distracted or debilitated by breeding activities; and failing this, to small animals and fruits.

Ambush predators retain a greater chance of killing healthy prey opportunistically than cursorial predators, so that felids are more obligate hunters than canids, hyenids, ursids or mustelids. Northern temperate ecosystems currently lack a top ambush predator equivalent to African lions (Schaller 1972) or Asian tigers (Karanth & Sunquist 1995), following the extinction of the saber-tooth "cats"

(*Smilodon* spp) that were once the supreme ambush predators there. Grey wolves are the top predator by default, but by concentrating on smaller mammals during summer they leave little meat uneaten, forcing coyotes to hunt even smaller prey. While polar bears in the far north are primarily carnivorous (Derocher et al. 2002), the brown and grizzly bears syntopic with wolves are largely omnivorous. Wolverines fill the role of a scavenger that hunts opportunistically in the far north of both Eurasia and North America, but are restricted to regions where the snow cover persists well into spring (Inman et al. 2012a).

Relationships among African carnivores are complex, with both lions and spotted hyenas shifting in their role as primarily predators or scavengers depending on their relative numbers. Hyenas remain carnivores through being able to subsist on portions of carcasses inedible to lions, in particular bones, and also through being able to drive other carnivores from their kills. Hence despite being competitive, these carnivores are also interdependent. Although African wild dogs adopt a similar hunting technique to wolves, they remain much less abundant than wolves because lions and hyenas preemptively deplete the vulnerable segment from large ungulate populations, and compete directly for smaller ungulates. The paucity of scavenging by African wild dogs and cheetahs probably reflects their dismal prospects of gaining much thereby, although both of them do consume carcasses in captivity.

The comparison between Holarctic and African carnivore assemblages calls into question why in South America the scavenger niche is filled entirely by birds (Keast 1969). Mammalian scavengers were represented by borhyaenids into the Pliocene (Marshall et al. 1982), but the placental carnivores that replaced them had no equivalent. The numerous equids and gomphotheres that existed in South America until the terminal Pleistocene (Webb 1977) would have provided an abundance of carcasses, albeit only seasonally in the absence of a top predator. Extant jaguar (*P. onca*) and puma subsist primarily on deer and peccaries (Aranda & Sánchez-Cordero 1996). Jaguars occasionally scavenge on cattle carcasses (López-

González & Piña 2002), in the absence of competition from any more obligate mammalian scavenger.

The limiting time for carnivores is when their herbivore prey are well-nourished and hence robust, which is during the summer or wet season months. The success of scavengers may then be crucially dependent on the presence of predators able to kill large ungulates and thereby provide large carcasses even at this time of the year; a situation that now persists solely in Africa and tropical Asia. The indirect cascading effects in food webs initiated by carrion shortages can be magnified by human hunting. On the one hand, intensive game hunting restricts the proportion of wild ungulate populations attaining terminal senescence and thus the carcasses produced by causes other than predation. On the other hand, hunting can lead to a carrion surfeit in areas where disposal of hunting leftovers from the field is a common practice (Wilmers et al. 2003). In Africa, subsidies from megaherbivore carcasses may disrupt pre-existing patterns of hunting versus scavenging, with ramifying consequences for prey populations.

Human impacts can have further, indirect implications for mammalian carnivores. Northern temperate ecosystems have been shown to be highly vulnerable to global climatic change. Global warming has been associated with a shorter period with deep snow during winter in Yellowstone National Park, reducing the number of ungulates dying from starvation, and hence the supply of carcasses to carrion-reliant species during this period (Wilmers & Getz 2005). However, the carcasses contributed through predation by wolves, together with the opening of intact carcasses of large ungulates by these top carnivores, counteract the food bottleneck to some extent. Here, the protection of top carnivores, widely threatened worldwide (e.g. Gittleman et al. 2001), is essential for maintaining the entire ecosystem. Global warming consequences on ungulate mortality patterns in African savannas could be very different. There, an expansion of the dry period might

increase the number of herbivores succumbing to malnutrition and diseases, thus producing more carcasses to be consumed by opportunistic scavengers.

Finally, long-distance herbivore migrations have been largely constrained in Africa over the last two centuries due to human pressure and to the fencing of protected reserves. This could dramatically affect the carrion subsidies available in some areas, thus diminishing scavenging opportunities for carnivores and increasing predation pressure on prey populations. Ultimately protected areas may become too small to conserve viable populations of the larger carnivores, with consequences ramifying through the food web.

Overall, we must recognise that relationships among carnivores based on hunting versus scavenging strategies are flexible and subject to changes in response to prevailing circumstances. This functional complexity must be taken into account when assessing the consequences of global change for ecosystem function.

ACKNOWLEDGEMENTS

The last author was supported by a postdoctoral grant from the Spanish Ministry of Education (Plan Nacional de I+D+i 2008-2011).

REFERENCES

- Aranda M, Sánchez-Cordero V (1996) Prey spectra of Jaguar (*Panthera onca*) and Puma (*Puma concolor*) in tropical forests of Mexico. *Studies of Neotropical Fauna and Environment* 31: 65–67.
- Atkinson RPG, Macdonald DW, Kamizota R (2002) Dietary opportunism in side-striped jackals. *Journal of Zoology* 257: 129–139.
- Bailey TN (1993) *The African Leopard. Ecology and Behaviour of a Solitary Felid*. Columbia University Press, New York.
- Barber-Meyer SM, Mech LD, White PJ (2008) Elk calf survival and mortality following wolf restoration to Yellowstone National Park. *Wildlife Monographs* 169: 1–30.
- Bekoff M, Wells MC (1981) Behavioural budgeting by wild coyotes: the influence of food resources and social organisation. *Animal Behaviour* 29: 794–801.

493 Bengis RG, Grant R, de Vos V (2003) Wildlife diseases and veterinary controls: a savanna ecosystem
 494 perspective. In: du Toit JT, Biggs HC, Rogers KH (eds) *The Kruger experience: ecology and*
 495 *management of savanna heterogeneity*, 349–369. Island Press, Washington D.C.
 496 Bertram B (1979) Serengeti predators and their social systems. In: Sinclair ARE, Norton-Griffiths M
 497 (eds) *Serengeti. Dynamics of an ecosystem*, 221–248. University of Chicago Press, Chicago.
 498 Carbone C, Teacher J, Rowcliffe M (2007) The costs of carnivory. *PLoS Biology* 5: 363–368.
 499 Caro TM, Stoner CJ (2003) The potential for interspecific competition among African carnivores.
 500 *Biological Conservation* 110: 67–75.
 501 Clutton-Brock TH, Albon SD (1982) Winter mortality in red deer. *Journal of Zoology* 198: 515–520.
 502 Conybeare A, Haynes G (1984) Observations on elephant mortality and bones in water holes.
 503 *Quaternary Research* 22: 189–200.
 504 Cooper SM (1990) The hunting behaviour of spotted hyenas in a region containing both sedentary and
 505 migratory prey. *African Journal of Ecology* 28: 131–141.
 506 Cooper SM (1991) Optimal hunting group size: the need for lions to defend their kills against loss to
 507 spotted hyaenas. *African Journal of Ecology* 29: 130–136.
 508 Cooper SM, Holekamp KE, Smale L (1999) A seasonal feast: long-term analysis of feeding behaviour in
 509 the spotted hyena (*Crocuta crocuta*). *African Journal of Ecology* 37: 149–160.
 510 Coulson T, Catchpole EA, Albon SD, Morgan BJT, Pemberton JM, Clutton-Brock TH, Crawley MJ,
 511 Grenfell BT (2001) Age, sex, density, winter weather, and population crashes in Soay sheep.
 512 *Science* 292: 1528–1531.
 513 Cortés-Avizanda A, Selva A, Carrete M, Donázar JA (2009) Effects of carrion resources on herbivore
 514 spatial distribution are mediated by facultative scavengers. *Basic and Applied Ecology* 10: 265–372.
 515 Cuesta L, Bárcena F, Palacios F, Reig S (1991) The trophic ecology of the Iberian Wolf (*Canis lupus*
 516 *signatus* Cabrera, 1907). A new analysis of stomach's data. *Mammalia* 55: 239–254.
 517 Crabtree RL, Sheldon JW (1999) The ecological role of coyotes on Yellowstone's northern range.
 518 *Yellowstone Science* 7: 15–23.
 519 Creel S, Creel NM (1996) Limitation of wild dogs by competition with larger carnivores. *Conservation*
 520 *Biology* 10: 526–538.
 521 Creel S (2001) Four factors modifying the effect of competition on carnivore population dynamics as
 522 illustrated by African wild dogs. *Conservation Biology* 15: 271–274.
 523 Derocher A, Wiig O, Andersen M (2002) Diet composition of polar bears in Svalbard and the western
 524 Barents Sea. *Polar Biology* 25: 448–452.

DeVault TL, Rhodes OE, Shivik JA (2003) Scavenging by vertebrates: behavioural, ecological and evolutionary perspectives on an important energy transfer pathway in terrestrial ecosystems. *Oikos* 102: 225–234.

DeVault T, Olson Z, Beasley JC, Rhodes OE (2011) Mesopredators dominate competition for carrion in an agricultural landscape. *Basic and Applied Ecology* 12: 268–274.

Durant SM (1998) Competition, refuges and coexistence: an example from Serengeti carnivores. *Journal of Animal Ecology* 67: 370–386.

Durant SM, Croft ME, Hilborn R, Bashir S, Hando J, Thomas L (2011) Long-term trends in carnivore abundance using distance sampling in Serengeti National Park, Tanzania. *Journal of Applied Ecology* 48: 1490–1500.

Fanshawe JH, Fitzgibbon CD (1993) Factors influencing the hunting success of an African wild dog pack. *Animal Behaviour* 45: 479–490.

Ferreira SM, Funston PJ (2010) Estimating lion population variables: prey and disease effects in Kruger National Park, South Africa. *Wildlife Research* 37: 194–206.

Fitzgibbon CD, Fanshawe JH (1989) The condition and age of Thomson's gazelles killed by cheetahs and wild dogs. *Journal of Zoology* 218: 99–108.

Fitzgibbon CD (1990a) Why do hunting cheetahs prefer male gazelles? *Animal Behaviour* 40: 837–845.

Fitzgibbon CD (1990b) Anti-predator behaviour strategies of immature Thomson's gazelles: hiding and the prone response. *Animal Behaviour* 40: 846–855.

Gasaway WC, Mossestad KJ, Stander PE (1989) Demography of spotted hyenas in an arid savanna, Etosha National Park, South West Africa, Namibia. *Madoqua* 16: 121–127.

Gasaway WC, Mossestad KT, Stander PE (1991) Food acquisition by spotted hyaenas in Etosha National Park, Namibia: predation versus scavenging. *African Journal of Ecology*, 29, 64–75.

Gasaway WC, Boertje RD, Grangaard DV, Kelleyhouse DG, Peterson RO, Larson DG (1992) Predation limiting moose at low densities in Alaska and Yukon and implications for conservation. *Wildlife monographs* 120: 3–59.

Gese EM, Grothe S (1995) Analysis of coyote predation on deer and elk during winter in Yellowstone National Park, Wyoming. *American Midland Naturalist* 133: 36–43.

Gese EM, Ruff RL, Crabtree RL (1996) Foraging ecology of coyotes (*Canis latrans*): the influence of extrinsic factors and a dominance hierarchy. *Canadian Journal of Zoology* 74: 769–783.

Gittleman JL, Funk SM, Macdonald D, Wayne RK (eds) (2001) *Carnivore conservation*. Cambridge University Press, Cambridge.

557 Grange S, Owen-Smith N, Gaillard J-M, Druce DJ, Moleón M, Mgobozi M. (2012) Changes of population
558 trends and mortality patterns in response to the reintroduction of large predators: The case study of
559 African ungulates. *Acta Oecologica* 42: 16–29.

560 Gustine DD, Parker KL, Lay RJ, Gillingham MP, Heard DC (2006) Calf survival of woodland caribou in a
561 multipredator ecosystem. *Wildlife Monographs* 165: 1–32.

562 Hamlin KL, Riley SJ, Pyrah D, Dod AR, Mackie RJ (1984) Relationships among mule deer fawn
563 mortality, coyotes, and alternative prey species during summer. *Journal of Wildlife Management* 48:
564 489–499.

565 Hanby JP, Bygott JD, Packer C (1995) Ecology, demography, and behavior of lions in two contrasting
566 habitats: Ngorongoro Crater and the Serengeti Plains. In: Sinclair ARE, Arcese P (eds) *Serengeti II*,
567 315–331. University of Chicago Press, Chicago.

568 Haynes G (1982) Utilization and skeletal disturbances of North American prey carcasses. *Arctic* 35:
569 266–281.

570 Hayward MW (2006) Prey preferences of spotted hyena (*Crocuta crocuta*) and degree of dietary overlap
571 with the lion (*Panthera leo*). *Journal of Zoology* 270: 666–614.

572 Hebblewhite M (2005) Predation by wolves interacts with North Pacific Oscillation on a western North
573 American elk population. *Journal of Animal Ecology* 74: 226–233.

574 Henschel JR, Skinner JD (1990) The diet of the spotted hyenas in Kruger National Park. *African Journal*
575 *of Ecology* 28: 69–82.

576 Hofer H, East ML (1993) The commuting system of Serengeti spotted hyaenas: how a predator copes
577 with migratory prey. I. Social organization. *Animal Behaviour* 46: 547–557.

578 Hofer H, East ML (1995) Population dynamics, population size, and the commuting system of Serengeti
579 spotted hyenas. In: Sinclair ARE, Arcese P (eds) *Serengeti II*, 332–363. University of Chicago
580 Press, Chicago.

581 Honer OP, Wachter B, East ML, Hofer H (2002) The response of spotted hyenas to long-term changes
582 in prey populations: functional response and interspecific kleptoparasitism. *Journal of Animal*
583 *Ecology* 71: 236–246.

584 Houston DC (1974) The role of griffon vultures *Gyps africanus* and *Gyps ruppellii* as scavengers.
585 *Journal of Zoology* 172: 35–46.

586 Houston DC (1979) The adaptation of scavengers. In: Sinclair ARE, Norton-Griffiths MN (eds)
587 *Serengeti. Dynamics of an ecosystem*, 263–286. University of Chicago Press, Chicago.

588 Houston DC, Cooper JE (1975) The digestive tract of the White-back Griffon Vulture and its role in
589 disease transmission among wild ungulates. *Journal of Wildlife Diseases* 11: 306–313.

590 Huegel CN, Rongstad OJ (1985) Winter foraging patterns and consumption rates of northern Wisconsin
 591 coyotes. *American Midland Naturalist* 113: 203–207.
 592 Huggard DJ (1994) Prey selectivity of wolves in Banff National Park. II. Age, sex and condition of elk.
 593 *Canadian Journal of Zoology* 71: 140–147.
 594 Husseman JS, Murray DL, Power G, Mack C, Wenger CR, Quigley H (2003) Assessing differential prey
 595 selection between two sympatric large carnivores. *Oikos* 101: 591–601.
 596 Inman RM, Nagoin AJ, Persson J, Mattison J (2012a) The wolverine's niche: linking reproductive
 597 chronology, caching, competition and climate. *Journal of Mammalogy* 93: 634–644.
 598 Inman, RM, Packila ML, Inman KH, McCue AJ, White GC, Persson J, Aber BC, Orme ML, Alt KL, Cain
 599 SL, Fredrick JA, Oakleaf BJ & Sartorius SS (2012b) Spatial ecology of wolverines at the southern
 600 periphery of distribution. *Journal of Wildlife Management* 76: 778–792.
 601 Jarnemo A, Liberg O, Lockewant S, Olsson A, Wahlstrom K (2004) Predation by red fox on European
 602 roe deer fawns in relation to age, sex and birth date. *Canadian Journal of Zoology* 82: 416–422.
 603 Jędrzejewski W, Schmidt K, Milkowski L, Jędrzejewska B, Okarma H (1993) Foraging by lynx and its
 604 role in ungulate mortality: the local (Białowieża Forest) and the Palaearctic viewpoints. *Acta*
 605 *Theriologica* 38: 385–403.
 606 Jędrzejewski W, Schmidt K, Theuerkauf J, Jędrzejewska B, Selva N, Zub K, Szymura L (2002) Kill rates
 607 and predation by wolves on ungulate populations in Białowieża Primeval Forest (Poland). *Ecology*
 608 83: 1341–1356.
 609 Joubert D (2006) Hunting behaviour of lions on elephants in the Chobe National Park, Botswana.
 610 *African Journal of Ecology* 44: 279–281.
 611 Karanth U, Sunquist ME (1995) Prey selection by tiger, leopard and dhole in tropical forests. *Journal of*
 612 *Animal Ecology* 64: 439–450.
 613 Keast A (1969) Comparisons of the contemporary mammalian faunas of the southern continents. *The*
 614 *Quarterly Review of Biology* 44: 121–167.
 615 Kruuk H (1972) *The spotted hyena: A study of predation and social behaviour*. University of Chicago
 616 Press, Chicago.
 617 Levi T, Wilmers CC (2012) Wolves-coyotes-foxes: a cascade among carnivores. *Ecology*, 93, 921–929.
 618 López-González CA, Piña L (2002) Carrion use by jaguars (*Panthera onca*) in Sonora, Mexico.
 619 *Mammalia* 66: 603–605.
 620 Mahoney SP, Schaefer JA (2002) Long-term changes in demography and migration of Newfoundland
 621 caribou. *Journal of Mammalogy* 83: 957–963.
 622 Marshall LG, Webb SD, Sepkowski JJ, Raup DM (1982) Mammalian evolution and the great American
 623 interchange. *Science* 215: 1351–1357.

624 Mattisson J, Persson J, Segerstrom P (2011) Temporal and spatial interactions between an obligate
 625 predator, the Eurasian lynx, and a facultative scavenger, the wolverine. *Canadian Journal of*
 626 *Zoology* 89: 79-89.
 627 Metz MC, Smith DW, Vucetich JA, Stahler DR, Peterson RO (2012) Seasonal patterns of predation for
 628 gray wolves in the multi-prey system of Yellowstone National Park. *Journal of Animal Ecology* 81:
 629 553–563.
 630 Mduma SAR, Sinclair ARE, Hilborn R (1999) Food regulates the Serengeti wildebeest: a 40-year record.
 631 *Journal of Animal Ecology* 68: 1101–1122.
 632 Mills MG (1990) *Kalahari hyaenas: the comparative behavioural ecology of two species*. Allen & Unwin,
 633 London.
 634 Mills MGL, Shenk TM (1992) Predator-prey relationships: The impact of lion predation on wildebeest
 635 and zebra populations. *Journal of Animal Ecology* 61: 693–702.
 636 Mills MGL, Biggs HC (1993) Prey apportionment and related ecological relationships between large
 637 carnivores and prey in the Kruger National Park. *Symposium of the Zoological Society of London*
 638 65: 253–268.
 639 Mills MGL, Gorman ML (1997) Factors affecting the density and distribution of wild dogs in the Kruger
 640 National Park. *Conservation Biology* 11: 1397–1406.
 641 Mills MGL, Juritz JM, Zucchini W (2001) Estimating the size of spotted hyaena populations through
 642 playback recordings allowing for non-response. *Animal Conservation* 4: 335–344.
 643 Mills MGL, Broomhall IS, du Toit JT (2004) Cheetah *Acinonyx jubatus* feeding ecology in the Kruger
 644 National Park and a comparison across African savanna habitats: is the cheetah only a successful
 645 hunter on open grass plains? *Wildlife Biology* 10: 177–186.
 646 Molinari-Jobin A, Molinari P, Loison A, Gaillard J-M, Brestenmoser V (2004) Life cycle period and
 647 activity of prey influence their susceptibility to predators. *Ecography* 27: 323–329.
 648 Nellis CH, Keith LB (1976) Population dynamics of coyotes in Central Alberta, 1964-68. *Journal of*
 649 *Owen-Smith N (1988) Megaherbivores. The influence of very large body size on ecology*. Cambridge
 650 University Press, Cambridge.
 651 Owen-Smith N (2002) *Adaptive herbivore ecology. From resources to populations in variable*
 652 *environments*. Cambridge University Press, Cambridge.
 653 Owen-Smith N (2008) Changing vulnerability to predation related to season and sex in an African
 654 ungulate assemblage: how additive is predation? *Oikos* 117: 602–610.
 655 Owen-Smith N, Mason DR (2005) Comparative changes in adult vs juvenile survival affecting population
 656 trends of African ungulates. *Journal of Animal Ecology* 74: 762–773.

657 Owen-Smith N, Mills MGL (2008) Shifting prey selection generates contrasting herbivore dynamics
658 within a large-mammal predator-prey web. *Ecology* 89: 1120–1133.

659 Owen-Smith N, Mason DR, Ogutu JO (2005) Correlates of survival rates for ten African ungulate
660 populations: density, rainfall and predation. *Journal of Animal Ecology* 74: 774–788.

661 Palomares F, Caro TM (1999) Interspecific killing among mammalian carnivores. *The American*
662 *Naturalist* 153: 492–508.

663 Paquet P (1992) Prey use strategies of sympatric wolves and coyotes in Riding Mountain National Park,
664 Manitoba. *Journal of Mammalogy* 73: 337–343.

665 Patterson BR, Benjamin LK, Messier F (1998) Prey switching and feeding habits of the eastern coyote in
666 relation to white-tailed deer and hare densities. *Canadian Journal of Zoology* 76: 1885–1897.

667 Peterson RO, Page RE (1988) The rise and fall of Isle Royale wolves 1975-1986. *Journal of*
668 *Mammalogy* 69: 89–99.

669 Pienaar U de V (1969) Predator-prey relationships amongst the larger mammals of the Kruger National
670 Park. *Koedoe* 12: 108–176.

671 Pole A, Gordon IJ, Gorman ML, Macaskill M (2004) Prey selection by African wild dogs in southern
672 Zimbabwe. *Journal of Zoology* 262: 207–215.

673 Post E, Peterson RO, Stenseth NC, McLaren BE (1999) Ecosystem consequences of wolf behavioural
674 responses to climate. *Nature* 401: 905–907.

675 Prugh L (2005) Coyote prey selection and community stability during a decline in food supply. *Oikos*
676 110: 253–264.

677 Ruxton GD, Houston DC (2004) Obligate vertebrate scavengers must be large soaring fliers. *Journal of*
678 *Theoretical Biology* 228: 431–436.

679 Schaller GB (1972) *The Serengeti lion: A study of predator prey relations*. University of Chicago Press,
680 Chicago.

681 Sala E (2006) Top predators provide insurance against climate change. *Trends in Ecology and*
682 *Evolution* 21: 479–480.

683 Selva N, Jędrzejewska B, Jędrzejewski W, Wajrak A (2003) Scavenging on European bison carcasses
684 in Białowieża Primeval Forest (eastern Poland). *Ecoscience* 10: 303–311.

685 Selva N, Jędrzejewska B, Jędrzejewski W, Wajrak A (2005) Factors affecting carcass use by a guild of
686 scavengers in European temperate woodland. *Canadian Journal of Zoology* 83: 1590–1601.

687 Śmietana W, Klimek A (1993) Diet of wolves in the Bieszczady Mountains, Poland. *Acta Theriologica*
688 38: 245–251.

689 Sinclair ARE, Mduma S, Brashares JS (2003) Patterns of predation in a diverse predator-prey system.
690 *Nature* 425: 288–290.

691 Sinclair ARE, Arcese P (1995) Population consequences of predation-sensitive foraging. *Ecology* 76:
692 882–891.

693 Skinner JD, Chimimba CT (2005) *The Mammals of the Southern African Subregion*. Cambridge
694 University Press, Cambridge.

695 Smith DW, Peterson RO, Houston DB 2003. Yellowstone after wolves. *BioScience* 53: 330–340.

696 Stander PE (1992) Foraging dynamics of lions in a semi-arid environment. *Canadian Journal of Zoology*
697 70: 8–21.

698 Steinmetz J, Soluk DA, Kohler SL (2008) Facilitation between herons and smallmouth bass foraging on
699 common prey. *Environmental Biology of Fishes* 81: 51–61.

700 Swenson JE, Dahle B, Busk H, Opseth O, Johansen T, Söderberg, A, Walin K, Cederlund G (2007)
701 Predation on moose by European brown bears. *Journal of Wildlife Management* 71: 1993–1997.

702 Tveraa T, Fauchald P, Henaug C, Yoccoz NG (2003) An examination of a compensatory relationship
703 between food limitation and predation in semi-domestic reindeer. *Oecologia* 137: 370–376.

704 Valdmann H, Andersone-Lilley Z, Koppa O, Ozolins J, Bagrade G (2005) Winter diets of wolf *Canis*
705 *lupus* and lynx *Lynx lynx* in Estonia and Latvia. *Acta Theriologica* 50: 521–527.

706 Van Dijk J, Gustavsen L, Mysterud A, May R, Flagstad O, Broseth H, Andersen R, Anderson R, Steen
707 H, London A (2008) Diet shift of a facultative scavenger, the wolverine following recolonisation of
708 wolves. *Journal of Animal Ecology* 77: 1183–1190.

709 Watts PD, Jonkel C (1988) Energetic cost of winter dormancy in grizzly bear. *Journal of Wildlife*
710 *Management* 177: 193–208.

711 Webb SD (1977) A history of savanna vertebrates in the New World. Part II: South America and the
712 great interchange. *Annual Review of Ecology and Systematics* 9: 393–426.

713 Wilmers CC, Getz WM (2004) Simulating the effects of wolf-elk population dynamics on resource flow to
714 scavengers. *Ecological Modelling* 177: 193–208.

715 Wilmers CC, Getz, WM (2005) Gray wolves as climate change buffers in Yellowstone. *PLoS Biology* 3:
716 571–576.

717 Wilmers CC, Post E (2006) Predicting the influence of wolf-provided carrion on scavenger community
718 dynamics under climate change scenarios. *Global Change Biology* 12: 403–409.

719 Wilmers CC, Stahler DR, Crabtree RL, Smith D, Getz WM (2003) Resource dispersion and consumer
720 dominance: Scavenging at wolf and hunter-killed carcasses in Greater Yellowstone, USA. *Ecology*
721 *Letters* 6: 996–1003.

722 Wilson EE, Wolkovich EM (2011) Scavenging: how carnivore and carrion structure communities. *Trends*
723 *in Ecology and Evolution* 26: 129–135.