



City Research Online

City St George's, University of London

Citation: Pereira, L., Owen-Smith, N. & Moleon, M. (2013). Facultative predation and scavenging by mammalian carnivores: seasonal, regional and intra-guild comparisons. *Mammal review*, 44(1), pp. 44-55. doi: 10.1111/mam.12005

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/19463/>

Link to published version: <https://doi.org/10.1111/mam.12005>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Table 1. Relative contribution of hunting versus scavenging to the diets of lions and spotted hyenas in different places

Animal	Area	Estimated population	Regional density (no. 100 km ⁻²)	Ratio lion/hyena or hyena/lion densities	Contribution of hunting (%)	Contribution of scavenging (%)	Source
Lion	Serengeti ecosystem, Tanzania	3000	12	0.75	78 ¹	12 ¹	Schaller 1972, Bertram 1979
	Serengeti plains, Tanzania	200	7-10	0.08-0.12	47 ¹	33 ¹	Schaller 1972, Packer et al. 1993
	Ngorongoro crater, Tanzania	50-125	20-50	0.11-0.71	88 ¹	10 ¹	Hanby et al. 1995, Packer et al. 1993
	Kruger N. P., South Africa	1600	8	0.57	~100	?	Ferreira & Funston 2010
	Kruger N. P. (South-Central), South Africa	731	12	0.67	~100	?	Smuts 1978, Funston 1999
	Kalahari Gemsbok N. P., South Africa	150	3	1.5	95	5	Mills 1990
	Etosha N. P., Namibia	56	2.8	0.56	94.5	5.5	Stander 1992
Spotted hyena	Serengeti ecosystem, Tanzania	4000	16	1.33	43 ¹	33 ¹	Kruuk 1972, Borner et al. 1987
	Serengeti plains, Tanzania	2100 ²	84	8.4-12	?	?	Hofer & East 1995
	Ngorongoro crater, Tanzania	171-480	70-190	1.4-9.5	69-82	7-31 ¹	Kruuk 1972
	Masai Mara (Talek region), Kenya	65	110		85	15	Cooper et al. 1999
	Kruger N. P., South Africa	2800	14	1.75	50	?	Mills & Biggs 1993
	Kruger N. P. (– South-Central), South Africa	1000	18	1.5	?	?	Mills et al. 2001
	Kalahari Gemsbok N. P., South Africa	90	2	0.67	73	27	Mills 1990
	Etosha N. P., Namibia	100	5	1.79	75	?	Gasaway et al. 1989, 1991

¹Where percentages do not add up to 100%, the remainder is unknown

²Communting hyenas are down-weighted by 0.5