



City Research Online

City St George's, University of London

Citation: Linckelmann, M. (2018). A note on the depth of a source algebra over its defect group. *International Electronic Journal of Algebra*, 24, pp. 68-72. doi: 10.24330/ieja.440216

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/20246/>

Link to published version: <https://doi.org/10.24330/ieja.440216>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

A NOTE ON THE DEPTH OF A SOURCE ALGEBRA OVER ITS DEFECT GROUP

Markus Linckelmann

Received: 30 October 2017; Accepted: 24 April 2018

Communicated by Burcu Üngör

Dedicated to the memory of Professor John Clark

ABSTRACT. By results of Boltje and Külshammer, if a source algebra A of a principal p -block of a finite group with a defect group P with inertial quotient E is a depth two extension of the group algebra of P , then A is isomorphic to a twisted group algebra of the group $P \rtimes E$. We show in this note that this is true for arbitrary blocks. We observe further that the results of Boltje and Külshammer imply that A is a depth two extension of its hyperfocal subalgebra, with a criterion for when this is a depth one extension. By a result of Watanabe, this criterion is satisfied if the defect groups are abelian.

Mathematics Subject Classification (2010): 20C05, 20C20, 16D20

Keywords: Source algebra, depth

Let p be a prime and $\mathcal{O}$ a complete local principal ideal domain with an algebraically closed residue field k of characteristic p , allowing the case $\mathcal{O} = k$. We will make without further comment use of the fact that by [9, II, Prop. 8], the canonical group homomorphism $\mathcal{O}^\times \rightarrow k^\times$ splits canonically, and hence group cohomology with coefficients in $k^\times$ can be viewed as cohomology with coefficients in $\mathcal{O}^\times$. Following terminology in [4], a ring extension $B \rightarrow A$ is called *of depth one* if A is isomorphic, as a B - B -bimodule, to a direct summand of B^n for some positive integer n , and a ring extension $B \rightarrow A$ is called *of depth two* if $A \otimes_B A$ is isomorphic, as an A - B -bimodule, to a direct summand of A^n , for some positive integer n . Tensoring by $A \otimes_B -$ shows that a ring extension of depth one is also an extension of depth two.

Let A be a source algebra of a block algebra over $\mathcal{O}$ of a finite group, with a defect group P . Boltje and Külshammer showed in [2, 2.4] that if A is isomorphic to a twisted group algebra of the form $\mathcal{O}_\alpha(P \rtimes E)$ for some p' -subgroup E of $\text{Aut}(P)$ and some $\alpha \in H^2(E; k^\times)$, inflated trivially to $P \rtimes E$, then the canonical map $\mathcal{O}P \rightarrow A$ is an extension of depth two. Moreover, they showed that the converse holds for principal blocks. The following result shows that this converse holds for arbitrary

blocks. See for instance [10, §11, §38] and [5, §6, §7] for background material on the Brauer homomorphism Br_P and fusion in source algebras.

Theorem 1. *Let G be a finite group, b a block of $\mathcal{O}G$, P a defect group of b and $A = i\mathcal{O}Gi$ a source algebra of b , where i is a primitive idempotent in the P -fixed point algebra $(\mathcal{O}Gb)^P$ such that $\text{Br}_P(i) \neq 0$. The following are equivalent:*

- (i) *The ring extension $\mathcal{O}P \rightarrow A$ induced by the canonical map $P \rightarrow A^\times$ is of depth two.*
- (ii) *The ring extension $kP \rightarrow k \otimes_{\mathcal{O}} A$ induced by the canonical map $P \rightarrow A^\times$ is of depth two.*
- (iii) *There is an isomorphism of interior P -algebras $A \cong \mathcal{O}_\alpha(P \rtimes E)$ for some p' -subgroup E of $\text{Aut}(P)$ and some $\alpha \in H^2(E; k^\times)$ inflated trivially to $P \rtimes E$.*
- (iv) *There is an isomorphism of interior P -algebras $k \otimes_{\mathcal{O}} A \cong k_\alpha(P \rtimes E)$ for some p' -subgroup E of $\text{Aut}(P)$ and some $\alpha \in H^2(E; k^\times)$ inflated trivially to $P \rtimes E$.*

Proof. The equivalence of (iii) and (iv) is an immediate consequence of results of Puig (either apply [7, 14.6] over both $\mathcal{O}$ and k , or use the lifting property [6, 7.8] for source algebras). Statement (iv) implies (i) and (ii) by Boltje and Külshammer [2, 2.4]. The implication (i) $\Rightarrow$ (ii) is trivial. It suffices to show that (ii) implies (iv). We may therefore assume that $\mathcal{O} = k$. Suppose that (ii) holds but that (iv) does not hold. As an A - kP -bimodule, A is indecomposable since $1_A = i$ is primitive in A^P . Thus, if (ii) holds, then the Krull-Schmidt theorem implies that any indecomposable direct summand of $A \otimes_{kP} A$ as an A - kP -bimodule is isomorphic to A as an A - kP -bimodule. Now if (iv) does not hold, then by [7, 14.6], there is a proper subgroup Q of P and an injective group homomorphism φ from Q to P such that the indecomposable kP - kP -bimodule $kP \otimes_{kQ} (\varphi kP)$ is isomorphic to a direct summand of A as a kP - kP -bimodule. Thus $A \otimes_{kQ} (\varphi kP)$ is isomorphic to a direct summand of $A \otimes_{kP} A$ as an A - kP -bimodule, and hence so is $Aj \otimes_{kQ} (\varphi kP)$, where j is a primitive idempotent in A^Q . Since Aj is indecomposable as an A - kQ -bimodule, so is the $k(G \times Q)$ -module kGj . Green's indecomposability theorem implies that the $k(G \times P)$ -module $kGj \otimes_{kQ} (\varphi kP)$ is indecomposable. Using that multiplication by i yields a Morita equivalence between kGb and A it follows that the A - kP -bimodule $Aj \otimes_{kQ} (\varphi kP)$ is also indecomposable, hence isomorphic to A as an A - kP -bimodule, by the above. Since $\text{Br}_P(i) \neq 0$ this is, however, only possible if $Q = P$, a contradiction. $\square$

For the sake of completeness, we mention that the depth of an extension $D \rightarrow A$, where D is a hyperfocal subalgebra (cf. [8]) in a source algebra A of a block of a finite group, can be determined essentially as an application of the methods from [1] and [2]. The first statement of the following proposition is a special case of [1, 1.5].

Proposition 2. *Let A be a source algebra of a block of a finite group algebra over $\mathcal{O}$ with defect group P , and let D be a hyperfocal subalgebra of A . The following hold.*

- (i) *The extension $D \rightarrow A$ is of depth two.*
- (ii) *The extension $D \rightarrow A$ is of depth one if and only if P acts by inner automorphisms on D .*

Proof. As mentioned above, statement (i) is a special case of [1, 1.5], as A is P/Q -graded, with D as 1-component. Since the argument is short and some parts of the notation will be useful in the proof of (ii), we sketch this briefly. We identify P with its canonical image in $A^\times$. The following definitions and facts on the hyperfocal subalgebra D of A are from [8]. The subalgebra D is P -stable, and the group $Q = P \cap D^\times$ is the $\mathcal{F}$ -hyperfocal subgroup of P , where $\mathcal{F}$ is the fusion system of A on P . An immediate consequence of these properties is that D is indecomposable as an $\mathcal{O}$ -algebra. Indeed, we have $D^P \subseteq A^P$, which is local, and hence P permutes the blocks of D transitively. But we also have $\text{Br}_P(1_A) \neq 0$, and hence D has a unique block. By [8, Theorem 1.8] we have $A = \bigoplus_{u \in [P/Q]} Du$, where $[P/Q]$ is a set of representatives in P of P/Q . Since D is P -stable, this is a decomposition of A as a D - D -bimodule. Thus $A \otimes_D A = \bigoplus_{u \in [P/Q]} A \otimes_D Du$ is a decomposition of $A \otimes_D A$ as an A - D -bimodule. For $u \in P$, a trivial verification shows that the A - D -bimodule $A \otimes_D Du$ is isomorphic to A via the map sending $a \otimes du$ to adu , where $a \in A$ and $d \in D$. Thus any indecomposable direct summand of the A - D -bimodule $A \otimes_D A$ is isomorphic to a direct summand of A as an A - D -bimodule. This proves (i). The summands Du in the D - D -bimodule decomposition $A = \bigoplus_{u \in [P/Q]} Du$ are all indecomposable as D - D -bimodules. Indeed, D is indecomposable by the above, and Du is isomorphic to the image of D under the Morita equivalence on $\text{mod}(D \otimes_{\mathcal{O}} D^{\text{op}})$ obtained from twisting the right D -module structure by the automorphism induced by conjugation with u . Thus the extension $D \rightarrow A$ is of depth one if and only if $Du \cong D$ as D - D -bimodules, for all $u \in [P/Q]$, hence for all $u \in P$. By standard facts on automorphisms (cf. [3, §55A]) this is equivalent to the condition that u induces an inner automorphism of D , for all $u \in P$. This proves (ii). $\square$

In conjunction with a result of Watanabe [11], this yields the following consequence.

Corollary 3. *With the notation of Proposition 2, if P is abelian, then the extension $D \rightarrow A$ is of depth one.*

Proof. By [11, Theorem 2], if P is abelian, then P acts as inner automorphisms on D . Thus the result follows from Proposition 2 (ii). $\square$

Remark 4. *What we have called depth two in this note is called right $D2$ in [4, 3.1], with left $D2$ being the obvious analogue, requiring $A \otimes_B A$ to be a direct summand, as a B - A -bimodule, of A^n for some positive integer n . It is easy to see directly that left and right $D2$ are equivalent conditions for the extensions $\mathcal{O}P \rightarrow A$ and $D \rightarrow A$ considered in the results above; this follows also from a more general result in [4, 6.4]. See [2, §2.3] for a related discussion.*

References

- [1] R. Boltje and B. Külshammer, *On the depth 2 condition for group algebra and Hopf algebra extensions*, J. Algebra, 323(6) (2010), 1783-1796.
- [2] R. Boltje and B. Külshammer, *Group algebra extensions of depth one*, Algebra Number Theory, 5(1) (2011), 63-73.
- [3] C. W. Curtis and I. Reiner, *Methods of Representation Theory*, Vol. II, with applications to finite groups and orders, Pure and Applied Mathematics (New York), A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1987.
- [4] L. Kadison and K. Szlachányi, *Bialgebroid actions on depth two extensions*, Adv. Math., 179(1) (2003), 75-121.
- [5] M. Linckelmann, *On splendid derived and stable equivalences between blocks of finite groups*, J. Algebra, 242(2) (2001), 819-843.
- [6] L. Puig, *Nilpotent blocks and their source algebras*, Invent. Math., 93(1) (1988), 77-116.
- [7] L. Puig, *Pointed groups and construction of modules*, J. Algebra, 116(1) (1988), 7-129.
- [8] L. Puig, *The hyperfocal subalgebra of a block*, Invent. Math., 141(2) (2000), 365-397.
- [9] J.-P. Serre, *Corps Locaux*, Deuxième édition, Publications de l'Université de Nancago, No. VIII, Hermann, Paris, 1968.

- [10] J. Thévenaz, *G-Algebras and Modular Representation Theory*, Oxford Mathematical Monographs, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1995.
- [11] A. Watanabe, *Note on hyperfocal subalgebras of blocks of finite groups*, J. Algebra, 322(2) (2009), 449-452.

Markus Linckelmann

Department of Mathematics

City, University of London

London EC1V 0HB

United Kingdom

e-mail: Markus.Linckelmann.1@city.ac.uk