



City Research Online

City St George's, University of London

Citation: Kessar, R. & Linckelmann, M. (2018). Dade's ordinary conjecture implies the Alperin–McKay conjecture. *Archiv der Mathematik*, 112(1), pp. 19-25. doi: 10.1007/s00013-018-1230-9

This is the published version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/20446/>

Link to published version: <https://doi.org/10.1007/s00013-018-1230-9>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).



Dade's ordinary conjecture implies the Alperin–McKay conjecture

RADHA KESSAR AND MARKUS LINCKELMANN

Abstract. We show that Dade's ordinary conjecture implies the Alperin–McKay conjecture. We remark that some of the methods can be used to identify a canonical height zero character in a nilpotent block.

Mathematics Subject Classification. 20C20.

Keywords. Dade's ordinary conjecture, Alperin–McKay conjecture, Height zero characters.

Dade proved in [4] that his projective conjecture [4, 15.5] implies the Alperin–McKay conjecture. Navarro showed in [11, Theorem 9.27] that the group version of Dade's ordinary conjecture implies the McKay conjecture. We show here that Dade's ordinary conjecture [3, 6.3] implies the Alperin–McKay conjecture. Let p be a prime number.

Theorem 1. *If Dade's ordinary conjecture holds for all p -blocks of finite groups, then the Alperin–McKay conjecture holds for all p -blocks of finite groups.*

The proof combines arguments from Sambale [17] and formal properties of chains of subgroups in fusion systems from [7]. Let $(K, \mathcal{O}, k)$ be a p -modular system. We assume that k is algebraically closed, and let $\bar{K}$ be an algebraic closure of K . By a character of a finite group, we will mean a $\bar{K}$ -valued character. For a finite group G and a block B of $\mathcal{O}G$, let $\text{Irr}(B)$ denote the set of irreducible characters of G in the block B , and let $\text{Irr}_0(B)$ denote the set of irreducible height zero characters of G in B . For a central p -subgroup Z of G and a character η of Z , let $\text{Irr}_0(B|\eta)$ denote the subset of $\text{Irr}_0(B)$ consisting of

This material is based upon work supported by the National Science Foundation under Grant No. DMS-1440140 while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Spring 2018 semester. The second author acknowledges support from EPSRC Grant EP/M02525X/1.

those height zero characters which cover the character η . The following lemma is implicit in [17].

Lemma 2. *Let P be a finite p -group, let $\mathcal{F}$ be a saturated fusion system on P , and let $Z \leq Z(\mathcal{F})$. Suppose that η is a linear character of P . There exists a linear character $\hat{\eta}$ of P such that $\hat{\eta}|_Z = \eta|_Z$ and $\text{foc}(\mathcal{F}) \leq \text{Ker}(\hat{\eta})$.*

Proof. First consider the case that $\eta|_Z$ is faithful. Then $Z \cap [P, P] = 1$. Hence by [5, Lemma 4.3], $\text{foc}(\mathcal{F}) \cap Z = 1$. The result is now immediate. Now suppose $Z_0 = \text{Ker}(\eta|_Z)$ and let $\bar{\mathcal{F}} = \mathcal{F}/Z_0$. By the previous argument, applied to P/Z_0 and $\bar{\mathcal{F}}$, there exists a character $\hat{\eta}$ of P/Z_0 such that $\hat{\eta}|_{Z/Z_0} = \eta|_{Z/Z_0}$ and $\text{foc}(\bar{\mathcal{F}}) \leq \text{Ker}(\hat{\eta})$. Denote also by $\hat{\eta}$ the inflation of $\hat{\eta}$ to P . Then $\hat{\eta}$ has the required properties since $\text{foc}(\bar{\mathcal{F}}) = \text{foc}(\mathcal{F})Z_0/Z_0$. $\square$

The following result is a special case of a result due to Murai; we include a proof for convenience.

Lemma 3 (cf. [9, Theorem 4.4]). *Let G be a finite group, B be a block of $\mathcal{O}G$, and P a defect group of B . Let Z be a central p -subgroup of G and let η be an irreducible character of Z such that $\text{Irr}_0(B|\eta) \neq \emptyset$. Then η extends to P .*

Proof. By replacing K by a suitable finite extension, we may assume that K is a splitting field for all subgroups of G . Let $i \in B^P$ be a source idempotent of B and let V be a KG -module affording an element of $\text{Irr}_0(B|\eta)$. Then $n := \dim_K(iV)$ is prime to p (see [13]). Since i commutes with P , iV is a KP -module via $x \cdot iv = ixv$, where $x \in P, v \in V$. Let $\rho : P \rightarrow \text{GL}_n(K)$ be a corresponding representation and let $\delta : P \rightarrow K^\times$ be the determinantal character of ρ . Then $\delta|_Z = \eta^n$. The result follows since n is prime to p . $\square$

Lemma 4. *Let G be a finite group, let B be a block of $\mathcal{O}G$ with a defect group P , and let Z be a central p -subgroup of G . Then $|\text{Irr}_0(B)|$ equals the product of $|\text{Irr}_0(B|1_Z)|$ with the number of distinct linear characters η of Z which extend to P .*

Proof. Let $\mathcal{F} = \mathcal{F}_{(P, e_P)}(G, B)$ be the fusion system of B with respect to a maximal B -Brauer pair (P, e_P) , and let η be a linear character of Z which extends to P . Since $Z \leq Z(\mathcal{F})$, by Lemma 2 there exists a linear character $\hat{\eta}$ of P such that $\hat{\eta}|_Z = \eta$ and $\text{foc}(\mathcal{F}) \leq \text{Ker}(\hat{\eta})$. By the properties of the Broué-Puig $*$ -construction [1, 16] the map $\chi \mapsto \hat{\eta} * \chi$ is a bijection between $\text{Irr}_0(B|1_Z)$ and $\text{Irr}_0(B|\eta)$. The result follows by Lemma 3. $\square$

Slightly strengthening the terminology in [10], we say that a pair (G, B) consisting of a finite group G and a block B of $\mathcal{O}G$ is a *minimal counterexample to the Alperin–McKay conjecture* if B is a counterexample to the Alperin–McKay conjecture and if G is such that first $|G : Z(G)|$ is smallest possible and then $|G|$ is smallest possible.

Proposition 5. *Let (G, B) be a minimal counterexample to the Alperin–McKay conjecture. Then $O_p(G) = 1$.*

Proof. By a result of Murai [10], we have that $Z := O_p(G)$ is central in G . Let P be a defect group of B and let C be the block of $\mathcal{O}N_G(P)$ in Brauer correspondence with B . By Lemma 4, $|\text{Irr}_0(B)| = |\text{Irr}_0(C)|$ if and only if $|\text{Irr}_0(\bar{B})| = |\text{Irr}_0(\bar{C})|$ where $\bar{B}$ (resp. $\bar{C}$) is the block of $\mathcal{O}G/Z$ (resp. $\mathcal{O}N_G(P)/Z$) dominated by B (resp. C). The result follows since $N_{G/Z}(P/Z) = N_G(P)/Z$ and $\bar{B}$ and $\bar{C}$ are in Brauer correspondence. $\square$

Let $\mathcal{F}$ be a saturated fusion system on a finite p -group P , and let $\mathcal{C}$ be a full subcategory of $\mathcal{F}$ which is upwardly closed; that is, if Q, R are subgroups of P such that Q belongs to $\mathcal{C}$ and if $\text{Hom}_{\mathcal{F}}(Q, R)$ is nonempty, then also R belongs to $\mathcal{C}$. Drawing upon notation and facts from [7, §5], $S_{\triangleleft}(\mathcal{C})$ is the category having as objects nonempty chains $\sigma = Q_0 < Q_1 < \cdots < Q_m$ of subgroups Q_i of P belonging to $\mathcal{C}$ such that $m \geq 0$ and Q_i is normal in Q_m , for $0 \leq i \leq m$. Morphisms in $S_{\triangleleft}(\mathcal{C})$ are given by certain ‘obvious’ commutative diagrams of morphisms in $\mathcal{F}$; see [7, 2.1, 4.1] for details. With this notation, the *length* of a chain σ in $S_{\triangleleft}(\mathcal{C})$ is the integer $|\sigma| = m$. The chain σ is called *fully normalised* if Q_0 is fully $\mathcal{F}$ -normalised and if either $m = 0$ or the chain $\sigma_{\geq 1} = Q_1 < Q_2 < \cdots < Q_m$ is fully $N_{\mathcal{F}}(Q_0)$ -normalised. Every chain in $S_{\triangleleft}(\mathcal{C})$ is isomorphic (in the category $S_{\triangleleft}(\mathcal{C})$) to a fully normalised chain. There is an involution n on the set of fully normalised chains which fixes the chain of length zero P and which sends any other fully normalised chain σ to a fully normalised chain $n(\sigma)$ of length $|\sigma| \pm 1$. This involution is defined as follows. If $\sigma = P$, then set $n(\sigma) = \sigma$. If $\sigma = Q_0 < Q_1 < \cdots < Q_m$ is a fully normalised chain different from P such that $Q_m = N_P(\sigma)$, then define σ by removing the last term Q_m ; if $Q_m < N_P(\sigma)$, then define σ by adding $N_P(\sigma)$ as last term to the chain. Then $n(\sigma)$ is fully normalised, and $n(n(\sigma)) = \sigma$. Denote by $[S_{\triangleleft}(\mathcal{C})]$ the partially ordered set of isomorphism classes of chains in $S_{\triangleleft}(\mathcal{C})$, and for each chain σ by $[\sigma]$ its isomorphism class. We have a partition

$$[S_{\triangleleft}(\mathcal{C})] = \{[P]\} \cup \mathcal{B} \cup n(\mathcal{B}),$$

where $\mathcal{B}$ is the set of isomorphism classes of fully normalised chains σ satisfying $|n(\sigma)| = |\sigma| + 1$. The following Lemma is a very special case of a functor cohomological statement [7, Theorem 5.11].

Lemma 6. *With the notation above, let $f : [S_{\triangleleft}(\mathcal{C})] \rightarrow \mathbb{Z}$ be a function on the set of isomorphism classes of chains in $S_{\triangleleft}(\mathcal{C})$ satisfying $f([\sigma]) = f([n(\sigma)])$ for any fully normalised chain σ in $S_{\triangleleft}(\mathcal{C})$. Then*

$$\sum_{[\sigma] \in [S_{\triangleleft}(\mathcal{C})]} (-1)^{|\sigma|} f([\sigma]) = f([P]).$$

Proof. The hypothesis on f implies that the contributions from chains in $\mathcal{B}$ cancel those from chains in $n(\mathcal{B})$, whence the result. $\square$

Proposition 7. *Let G be a finite group such that $O_p(G) = 1$, and let B be a block of $\mathcal{O}G$ with nontrivial defect groups. Suppose that Dade’s ordinary conjecture holds for B and that the Alperin–McKay conjecture holds for any block of any proper subgroup of G . Then the Alperin–McKay conjecture holds for the block B .*

Proof. Let (P, e) be a maximal B -Brauer pair, and denote by $\mathcal{F}$ the associated fusion system on P . For d a positive integer, denote by $\mathbf{k}_d(G, B)$ the number of ordinary irreducible characters in B of defect d . If $p^d = |P|$, then $\mathbf{k}_d(G, B)$ is the number of height zero characters, and if $p^d > |P|$, then $\mathbf{k}_d(G, B) = 0$.

Let $\mathcal{C}$ be the full subcategory of $\mathcal{F}$ consisting of all nontrivial subgroups of P . We briefly describe the standard translation process between chains in a fusion system of a block and the associated chains of Brauer pairs. The map sending a chain $\sigma = Q_0 < Q_1 < \cdots < Q_m$ in $S_{\triangleleft}(\mathcal{C})$ to the unique chain of nontrivial B -Brauer pairs $\tau = (Q_0, e_0) < (Q_1, e_1) < \cdots < (Q_m, e_m)$ contained in (P, e) induces a bijection between isomorphism classes of chains in $S_{\triangleleft}(\mathcal{C})$ and the set of G -conjugacy classes of normal chains of nontrivial B -Brauer pairs (cf. [7, 2.5]). If σ is fully normalised, then the corresponding chain of Brauer pairs $\tau = (Q_0, e_0) < (Q_1, e_1) < \cdots < (Q_m, e_m)$ has the property that $e_\tau = e_m$ remains a block of $N_G(\tau)$, and by [7, 5.14], $N_P(\sigma) = N_P(\tau)$ is a defect group of e_τ as a block of $N_G(\tau)$. Denote by $n(\tau)$ the chain of Brauer pairs corresponding to $n(\sigma)$.

Let $d > 0$ such that $p^d = |P|$. Define a function f on $S_{\triangleleft}(\mathcal{C})$ by setting

$$f([\sigma]) = \mathbf{k}_d(N_G(\tau), e_\tau)$$

for any fully normalised chain σ and corresponding chain τ of Brauer pairs. If $N_P(\sigma)$ is a proper subgroup of P , then $f([\sigma]) = 0$, and if $N_P(\sigma) = P$, then $f([\sigma])$ is the number of height zero characters of the block e_τ of $N_G(\tau)$. Dade's ordinary conjecture for B , reformulated here in terms of chains of Brauer pairs, asserts that $\mathbf{k}_d(G, B)$ is equal to the alternating sum

$$\sum_{[\sigma] \in S_{\triangleleft}(\mathcal{C})} (-1)^{|\sigma|} f([\sigma]).$$

The passage between formulations in terms of normalisers of chains of Brauer pairs rather than normalisers of chains of p -subgroups is well known; see e.g. [6, 4.5], [15].

If $|n(\sigma)| = |\sigma| + 1$, then, setting $H = N_G(\tau)$, we have $N_G(n(\tau)) = N_H(N_P(\tau), e_{n(\tau)})$; that is, $(N_P(\tau), e_{n(\tau)})$ is a maximal (H, e_τ) -Brauer pair. By the assumptions, the Alperin–McKay conjecture holds for the block e_τ of H . This translates to the equality $f([\sigma]) = f([n(\sigma)])$. That is, the function f satisfies the hypotheses of Lemma 6. Thus the above alternating sum is equal to $f([P])$, which by definition is $\mathbf{k}_d(N_G(P, e), e)$, and thus the Alperin–McKay conjecture holds for B . $\square$

Theorem 1 follows now immediately from combining Propositions 5 and 7.

Remark 8. By work of Dade [2] and Okuyama and Wajima [12], the Alperin–McKay conjecture holds for blocks of finite p -solvable groups. G. R. Robinson pointed out that Proposition 5 yields another short proof of this fact.

Remark 9. Let G be a finite group, B a block algebra of $\mathcal{O}G$, (P, e_P) a maximal (G, B) -Brauer pair with associated fusion system $\mathcal{F}$ on P , and let Z be a central p -subgroup of G . Let η be a linear character of Z , and suppose that η extends to a linear character $\hat{\eta}$ of P satisfying $\text{foc}(\mathcal{F}) \leq \text{Ker}(\hat{\eta})$. The proof of

Lemma 4 is based on the fact that the $*$ -construction $\chi \mapsto \hat{\eta} * \chi$ yields a bijection $\text{Irr}(B|1_Z) \rightarrow \text{Irr}(B|\eta)$. There is some slightly more structural background to this. For $\chi \in \text{Irr}(B)$, denote by $e(\chi)$ the corresponding central primitive idempotent in $K \otimes_{\mathcal{O}} B$. Set

$$e_1 = \sum_{\chi \in \text{Irr}_0(B|1_Z)} e(\chi), \quad e_{\eta} = \sum_{\chi \in \text{Irr}_0(B|\eta)} e(\chi).$$

Identify B with its image in $K \otimes_{\mathcal{O}} B$. Multiplying B by the central idempotents e_1 and e_{η} in $K \otimes_{\mathcal{O}} B$ yields the two $\mathcal{O}$ -free $\mathcal{O}$ -algebra quotients Be_1 and Be_{η} of B . By [8, Theorem 1.1], there is an $\mathcal{O}$ -algebra automorphism α of B which induces the identity on $k \otimes_{\mathcal{O}} B$ and which acts on $\text{Irr}(B)$ as the map $\chi \rightarrow \hat{\eta} * \chi$. Thus the extension of α to $K \otimes_{\mathcal{O}} B$ sends e_1 to e_{η} , and hence induces an $\mathcal{O}$ -algebra isomorphism

$$Be_1 \cong Be_{\eta}.$$

We conclude this note with an observation regarding canonical height zero characters in nilpotent blocks, based in part on some of the above methods.

Let G be a finite group, B a block algebra of $\mathcal{O}G$, P a defect group of B , and $i \in B^P$ a source idempotent of B . Denote by $\mathcal{F}$ the fusion system of B on P determined by the choice of i . Suppose that K is a splitting field for all subgroups of G . For V a finitely generated $\mathcal{O}$ -free B -module, denote by

$$\Delta_{V,P,i} : P \rightarrow \mathcal{O}^{\times}$$

the map sending $u \in P$ to the determinant of the $\mathcal{O}$ -linear automorphism of iV induced by the action of u on V (this makes sense since all elements in P commute with i). By standard properties of determinants, this map depends only on the $(B^P)^{\times}$ -conjugacy class of i and the isomorphism class of the $K \otimes_{\mathcal{O}} B$ -module $K \otimes_{\mathcal{O}} V$. Thus if V affords a character $\chi \in \text{Irr}(B)$, we write $\Delta_{\chi,P,i}$ instead of $\Delta_{V,P,i}$.

Proposition 10. *With the notation above, let $\chi \in \text{Irr}(B)$ and $\eta \in \text{Irr}(P/\text{foc}(P))$. Regard η as a linear character of P . We have*

$$\Delta_{\eta * \chi, P, i} = \eta^{\chi(i)} \Delta_{\chi, P, i}.$$

Proof. The statement makes sense as the value of χ on an idempotent is a positive integer. Let V be an $\mathcal{O}$ -free $\mathcal{O}G$ -module affording χ . By [8, Theorem 1.1] there exists an $\mathcal{O}$ -algebra automorphism α of B such that the module V^{α} (obtained from twisting V by α) affords $\eta * \chi$ and such that $\alpha(ui) = \eta(u)ui$ for all $u \in P$. Since in particular $\alpha(i) = i$, it follows that

$$\Delta_{V^{\alpha}, P, i}(u) = \Delta_{V, P, i}(\eta(u)u)$$

for all $u \in P$. The result follows as $\text{rank}_{\mathcal{O}}(iV) = \chi(i)$. $\square$

Denote by $\text{Irr}'(B)$ the set of all $\chi \in \text{Irr}(B)$ such that $\Delta_{\chi, P, i}$ is the trivial map (sending all elements in P to 1). Set $\text{Irr}'_0(B) = \text{Irr}'(B) \cap \text{Irr}_0(B)$. The maximal local pointed groups on B are G -conjugate. Thus if P' is any other defect group of B and $i' \in B^{P'}$ a source idempotent, then there exist $g \in G$ and $c \in (B^{P'})^{\times}$ such that $P' = gPg^{-1}$ and $i' = c g i g^{-1} c^{-1}$. Therefore the map

$\Delta_{V,P,i}$ is trivial if and only if the map $\Delta_{V,P',i'}$ is trivial, and hence the sets $\text{Irr}'(B)$ and $\text{Irr}'_0(B)$ are independent of the choice of P and i . The following is immediate.

Proposition 11. *The sets $\text{Irr}'(B)$ and $\text{Irr}'_0(B)$ are invariant under any automorphism of G which stabilises B .*

The next result shows that if B is nilpotent, then $\text{Irr}'_0(B)$ consists of a single element.

Proposition 12. *Suppose that B is nilpotent. Then $|\text{Irr}'_0(B)| = 1$. Moreover, if p is odd, then the unique element of $\text{Irr}'_0(B)$ is the unique p -rational height zero character in B .*

Proof. Let $\chi \in \text{Irr}_0(B)$. Since i is a source idempotent of B , $\chi(i)$ is prime to p . Hence if η, ζ are linear characters of P , then $\eta^{\chi(i)} = \zeta^{\chi(i)}$ implies that $\eta = \zeta$. Since B is nilpotent, we have that $\text{foc}(\mathcal{F}) = [P, P]$ and $|\text{Irr}_0(B)| = |P : [P, P]|$. Thus, by Proposition 10, the map $\chi \mapsto \Delta_{\chi,P,i}$ is a bijection from $\text{Irr}_0(B)$ to $\text{Irr}(P/[P, P])$. This proves the first assertion.

Suppose that p is odd. Let χ_0 be the unique p -rational character in $\text{Irr}_0(B)$. Let $W(k)$ be the ring of Witt vectors in $\mathcal{O}$. By the structure theory of nilpotent blocks (see [14]), there exists a $W(k)G$ -module V affording χ_0 . Since the source idempotent i can be chosen to be in $W(k)G$, we have that $\Delta_{\chi,P,i}$ takes values in $W(k)$. Since p is odd, it follows that the trivial character of P is the unique linear character of P which takes values in $W(k)$. $\square$

Acknowledgements. We thank Gunter Malle for helpful comments on the first version of the paper.

Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- [1] Broué, M., Puig, L.: Characters and local structure in G -algebras. J. Algebra **63**, 306–317 (1980)
- [2] Dade, E.C.: A correspondence of characters, The Santa Cruz Conference on Finite Groups (Univ. California, Santa Cruz, Calif., 1979), In: Proceedings of Symposia in Pure Mathematics, vol. 37, pp. 401–403. American Mathematical Society, Providence (1980)
- [3] Dade, E.C.: Counting characters in blocks, I. Invent. Math. **109**, 187–210 (1992)
- [4] Dade, E.C.: Counting characters in blocks, II. J. Reine Angew. Math. **448**, 97–190 (1994)
- [5] Diaz, A., Glessner, A., Mazza, N., Park, S.: Control of transfer and weak closure in fusion systems. J. Algebra **323**, 382–392 (2010)

- [6] Linckelmann, M.: Alperin's weight conjecture in terms of equivariant Bredon cohomology. *Math. Z.* **250**, 495–513 (2005)
- [7] Linckelmann, M.: The orbit space of a fusion system is contractible. *Proc. Lond. Math. Soc.* **98**, 191–216 (2009)
- [8] Linckelmann, M.: On automorphisms and focal subgroups of blocks. Preprint (2016). [arXiv:1612.07739](https://arxiv.org/abs/1612.07739)
- [9] Murai, M.: Block induction, normal subgroups and characters of height zero. *Osaka J. Math.* **31**, 9–25 (1994)
- [10] Murai, M.: On a minimal counterexample to the Alperin–McKay conjecture. *Proc. Japan Acad.* **87**, 192–193 (2011)
- [11] Navarro, G.: Character theory and the McKay conjecture. *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge (2018). <https://doi.org/10.1017/9781108552790>
- [12] Okuyama, T., Wajima, M.: Character correspondence and p -blocks of p -solvable groups. *Osaka J. Math.* **17**, 801–806 (1980)
- [13] Picaronny, C., Puig, L.: Quelques remarques sur un thème de Knörr. *J. Algebra* **109**, 69–73 (1987)
- [14] Puig, L.: Nilpotent blocks and their source algebras. *Invent. Math.* **93**, 77–116 (1988)
- [15] Robinson, G.R.: Weight conjectures for ordinary characters. *J. Algebra* **276**, 761–775 (2004)
- [16] Robinson, G.R.: On the focal defect group of a block, characters of height zero, and lower defect group multiplicities. *J. Algebra* **320**(6), 2624–2628 (2008)
- [17] Sambale, B.: On the projective height zero conjecture. [arXiv:1801.04272](https://arxiv.org/abs/1801.04272)

RADHA KESSAR AND MARKUS LINCKELMANN

Department of Mathematics,

City, University of London,

London EC1V 0HB,

UK

e-mail: radha.kessar.1@city.ac.uk

MARKUS LINCKELMANN

e-mail: markus.linckelmann.1@city.ac.uk

Received: 15 May 2018