



City Research Online

City St George's, University of London

Citation: Lockett, R. D. (2009). Instabilities and soot formation in high pressure explosion flames. Paper presented at the The British-French Flame Days, 02 - 03 April 2009, Lille, France.

This is the unspecified version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/2068/>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Instabilities and soot formation in high pressure, rich, iso-octane-air explosion flames

R.D. Lockett, R. Morishima



Introduction

- Flame instabilities
- Schlieren and OH PLIF experiments in the Leeds Bomb
- Results
- LII experiments in the Shell Bomb
- Results
- Soot formation hypothesis
- Conclusion

Flame Instabilities

1. Rayleigh-Taylor Instability

This instability is a result of a cold fluid above a hot fluid. The hot fluid is less dense than the cold fluid. Therefore the hot fluid is buoyant relative to the cold fluid.



Therefore irregularities at the interface of the two fluids grow, in order to achieve convection.

Courtesy of R. Woolley, Univ. of Leeds


Flame Instabilities

3. Thermal-Diffusive Instability

$$Le = \frac{\alpha}{D} < Le^* \equiv O(1)$$



The Leeds Bomb




Schlieren Cinematography in the Leeds Bomb

Conditions:


Iso-octane-air mixture
 $\phi = 1.0$, $p = 5$ bar

Viewing diameter
 $D = 15$ cm

Courtesy of R. Woolley,
Univ. of Leeds, 2005




OH PLIF in the Leeds Bomb




OH PLIF in the Leeds Bomb

5.5 cm



Unprocessed OH PLIF Image of a $\phi = 1$
5 bar Iso-octane-Air Explosion Flame
obtained in the Leeds Bomb (Flame
radius ≈ 60 mm, $Pe \sim 600$)

This reveals the influence of the
Darrius-Landau (hydrodynamic)
instability.



Schlieren Cinematography in the Leeds Bomb

Conditions:


Iso-octane-air mixture

$\phi = 1.4$, $p = 5$ bar

Viewing diameter


$D = 15$ cm

Courtesy of R. Woolley,
Univ. of Leeds, 2005




Schlieren Images of Iso-octane-Air Explosion Flames (flame radius ~ 60 mm, $Pe \sim 600$)

$P = 5$ bar, $\phi = 1.0$





$P = 5$ bar, $\phi = 1.4$





Schlieren Images of Iso-octane-Air Explosion Flames (flame radius ~ 60 mm, $Pe \sim 600$)

$P = 5$ bar, $\phi = 1.0$




$P = 5$ bar, $\phi = 1.4$




OH PLIF in the Leeds Bomb

5.5 cm







Unprocessed OH PLIF Image of a $\phi = 1.4$, 5 bar Iso-octane-Air Explosion Flame obtained in the Leeds Bomb (Flame radius ≈ 60 mm, $Pe \sim 600$)

This reveals the influence of the Darrius-Landau (hydrodynamic) instability and the thermal-diffusive instability.




Cell Length Scale Analysis

5.5 cm






Soot Formation Measurements in the Leeds Bomb Simultaneous Rayleigh/OH PLIF Spectroscopy




Soot Formation in the Leeds Bomb

Soot formation in a 2 bar,
 $\phi = 2.0$ iso-octane-air flame.


Note the soot formed behind
deep cracks in the flame.



LII/Mie Scattering in the Shell Bomb




LII/Mie Scattering in the Shell Bomb




LII/Mie Scattering in the Shell Bomb

Processed LII image
obtained from a $\phi = 1.8$,
5 bar, iso-octane-air flame
(flame radius ~ 60 mm)


(relative soot volume
fraction)





Processed Mie scattering
image obtained from a
 $\phi = 1.8$, 5 bar, iso-octane-
air flame




Relative soot particle
size distribution d_{63}



Relative soot particle
number density



Soot Formation in Rich, High Pressure, Spherically Expanding Explosion Flames



Conclusions

1. Two distinct length scales associated with flame cracking have been observed from the Schlieren images and the OH PLIF images.
2. These length scales are identified with hydrodynamic effects and thermal-diffusive effects. The large length scale cracking (5 mm to 1 cm) is associated with the hydrodynamic instability, while the small length scale cracking is associated with the thermal-diffusive instability (~ 1 mm).
3. High pressure flames that are stable to thermal-diffusive cracking develop hydrodynamic cracks which do not develop into discrete cells, while high pressure flames unstable to thermal-diffusive cracking exhibit full cellular structure and hydrodynamic perturbations.
4. Flame reaction quenching has been observed in the regions between the smaller length scale flame cells.

Conclusions Cont.

5. In highly enriched, high pressure explosion flames ($\phi > 1.8$), soot was observed to be formed in a honeycomb-like structure behind the flame.
6. The soot cell size was observed to be of the order of 5 mm to 1 cm, which corresponded with the larger length scale cellularity, determined by the hydrodynamic instability.
7. A plausible hypothesis for the formation of soot in highly enriched, spherical explosion flames has been suggested.