



City Research Online

City St George's, University of London

Citation: Hadjichrysanthou, C. & Broom, M. (2012). When should animals share food? Game theory applied to kleptoparasitic populations with food sharing. Behavioral Ecology, 23(5), pp. 977-991. doi: 10.1093/beheco/ars061

This is the unspecified version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/2166/>

Link to published version: <https://doi.org/10.1093/beheco/ars061>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

1 When should animals share food? Game theory 2 applied to kleptoparasitic populations with food 3 sharing 4

5 Abstract

6 Animals adopt varied foraging tactics in order to survive. Kleptoparasitism,
7 where animals attempt to steal food already discovered by others, is very common
8 among animal species. In this situation, depending on the ecological conditions,
9 challenged animals might defend, share or even, retreat and leave their food to
10 the challenger. A key determinant of the likely behaviour is the nature of the
11 food itself. If food is discovered in divisible clumps, it can be divided between
12 animals in a number of ways. This is the general assumption in one type of game-
13 theoretical models of food stealing, producer-scrounger models. Alternatively, food
14 items may be essentially indivisible, so that sharing is impossible and either the
15 attacker or defender must retain control of all of the food. This is the assumption
16 of the alternative game-theoretical models of kleptoparasitism. In this paper, using
17 a game-theoretic approach, we relax this assumption of indivisibility and introduce
18 the possibility of limited food sharing behaviour between animals in kleptoparasitic
19 populations. Considering the conditions under which food sharing is likely to be
20 common, it is shown that food sharing should occur in a wide range of ecological
21 conditions. In particular, if food availability is limited, the sharing process does not
22 greatly reduce the short-term consumption rate of food and food defence has a high
23 cost and/or a low probability of success, then the use of the food sharing strategy is
24 beneficial. Thus, the assumption of the indivisibility of food items is an important
25 component of previous models.

26 *Key words:* social foraging, food stealing, evolutionary games, strategy, ESS.

27 Introduction

28 In many biological situations, animals may decide to share their food in order to avoid
29 any injuries or energetic and time costs of a possible conflict with an attacking foraging
30 animal, or to obtain other immediate or delayed benefits such as mating opportunities
31 and reciprocal altruism. Food sharing is commonly observed in animal populations in a
32 wide range of species, including social carnivores, insects, birds, cetaceans, vampire bats
33 and primates (see Feistner and McGrew, 1989; Stevens and Gilby, 2004, for reviews).
34 In the literature, food sharing is defined in many different ways and various theoretical
35 models have been developed to consider the different biological situations where food
36 sharing among animals occurs. In the rest of this paper, we consider food sharing in
37 kleptoparasitic populations, populations where foraging animals steal food discovered
38 by others. We define food sharing to be the situation where the resource owner shows
39 tolerance and allows a competitor animal to consume a part of its food although it has
40 the ability to fight and try to keep all of its food.

41 Kleptoparasitism is a common foraging strategy. Different forms of kleptoparasitic
42 behaviour are observed in many species of animals, for example species of spiders (e.g.,
43 Coyl et al., 1991), birds (e.g., Brockman and Barnard, 1979), snails (e.g., Iyengar, 2002),
44 lizards (e.g., Cooper and Perez-Mellado, 2003), fish (e.g., Hamilton and Dill, 2003), pri-
45 mates (e.g., Janson, 1985), carnivores (e.g., Carbone et al., 2005) and insects (e.g.,
46 Erlandsson, 1988). This behaviour of animals has been recently well documented in a
47 review paper (Iyengar, 2008). The biological phenomenon of kleptoparasitism has at-
48 tracted the interest of many researchers from different areas. There are a number of
49 theoretical models focused on the kleptoparasitic behaviour of animals using different
50 mathematical methods, in particular evolutionary game theory. Two of the fundamen-
51 tal game-theoretical models which consider kleptoparasitic behaviour are the producer-
52 scrounger model, originally introduced by Barnard and Sibly (1981), and the model
53 of Broom and Ruxton (1998). In its original form, the producer-scrounger game is a
54 frequency-dependent game where animals forage for food using two strategies. They
55 either search for food (producer's strategy) or search for opportunities to kleptopara-
56 sitize (scrounger's strategy). The scrounger strategy does better when scroungers are

57 rare and worse when they are common. When the frequency of the two strategies is
58 such that the payoff obtained by each strategy is the same, there is a stable equilibrium
59 where the two strategies coexist. Many variations of this model have followed in order
60 to consider different factors that might affect the foraging process (e.g., Caraco and
61 Giraldeau, 1991; Vickery et al., 1991; Dubois and Giraldeau, 2005). One key feature
62 of this type of models is that food is usually discovered in patches and can be easily
63 split over foraging animals. Hence, the concept of food sharing is central to these mod-
64 els. In addition, in these models costs from aggressive strategies are energetic, rather
65 than time, costs. Thus, the different strategies do not directly affect the distribution of
66 feeding and foraging animals and the main effect of population density is to reduce the
67 “finder’s share”, the portion of the food eaten by a finder before other foragers discover
68 it. The model of Broom and Ruxton (1998), based on the mechanistic model of Ruxton
69 and Moody (1997), follows a different approach. Food comes in single indivisible items,
70 which must be consumed completely by an individual. Thus, food can never be shared
71 and challenging animals attempt to steal the whole item from the owner (or not). In
72 particular, in this model it is assumed that each of the animals in the population ei-
73 ther searches for food, has already acquired and is handling a food item prior to its
74 consumption or fights with another animal over a food item. When foraging animals
75 encounter an animal in the handling state, they can either decide to attack in order to
76 steal the prey or ignore the handler animal and continue searching. Attacked animals
77 always defend their food and a fight takes place. The population density has a direct
78 effect in this model as fights take time, with this loss of time the cost of more aggressive
79 strategies, and the more potential kleptoparasites there are, the more time is wasted on
80 fighting. The model predicts the optimal strategy for a foraging animal (to attack or
81 not to attack) under varying food availability and fight time cost. Broom et al. (2004)
82 later reconstructed this model in a more general framework by introducing different
83 competitive abilities between the attacker and the attacked animal and allowing the
84 attacked animal to surrender its food to the attacker avoiding the time cost of a fight. A
85 series of publications has appeared developing the original model of Broom and Ruxton
86 (1998) in a number of ways (e.g., Broom and Ruxton, 2003; Broom and Rychtar, 2007;

87 Luther et al., 2007; Yates and Broom, 2007; Broom et al., 2008; Broom and Rychar,
88 2009; Broom and Rychtar, 2011). Crowe et al. (2009) provide a brief review on the main
89 theoretical work on kleptoparasitism prior to the investigation of a stochastic model of
90 kleptoparasitism in finite populations. A comparison between some main models of
91 kleptoparasitism following the two fundamental game-theoretic approaches is discussed
92 in Vahl (2006) (see Chapter 6) and an alternative model is presented. There is also a
93 series of related mechanistic, but not game-theoretic, models which investigate interfer-
94 ence competition where foraging animals engage in aggressive interactions in order for
95 example to defend their territory, resulting in negative effects on their foraging efficiency
96 (e.g., Beddington, 1975; Ruxton et al., 1992; Van der Meer and Ens, 1997; Vahl, 2006;
97 Smallegange and Van der Meer, 2009; Van der Meer and Smallegange, 2009).

98 There are many game-theoretical models which investigate sharing behaviour as an
99 alternative strategy of foraging animals. The Dove strategy in the famous and widely
100 used Hawk-Dove game (Maynard Smith and Price, 1973; Maynard Smith, 1982) can be
101 thought of as an example of this non-aggressive behaviour. However, the Hawk-Dove
102 game and a large number of variations of this game (see for example, Sirot, 2000; Dubois
103 et al., 2003) are unable to show why in many biological situations animals prefer to share
104 the acquired prey without any contest taking place. The non-aggressive behaviour of the
105 Dove is shown to never be a pure Evolutionarily Stable Strategy (ESS), i.e. a strategy
106 that if adopted by the population cannot be invaded by any alternative strategy, and
107 can only exist as a mixed ESS (with Hawks) in a proportion depending on the value of
108 the resource and the cost of a potential contest. This is mainly due to the fact that the
109 Hawk-Dove game considers just a single contest between the two strategies. Although
110 the reward of adopting the Hawk strategy against an animal playing Hawk might be
111 equal or lower than the reward of adopting the Dove strategy, in a contest between a
112 Hawk and a Dove, the Hawk is always the optimal strategy. However, in group foraging
113 populations, animals usually have repeated interactions over food items. In iterated
114 Hawk-Dove type games, it has been shown that if the attacked animal can adopt the
115 strategy of its opponent (for example play a Retaliator type strategy (Maynard Smith
116 and Price, 1973; Maynard Smith, 1982) or a tit-for-tat type strategy (Axelrod and

Hamilton, 1981)) then, under some circumstances, food sharing without any aggressive interactions might be an ESS (Dubois and Giraldeau, 2003, 2007). A different game-theoretical food sharing model is considered in Stevens and Stephens (2002) in a situation where the owner of the food might decide to share its food with a beggar due to the fitness costs of harassment or interference (e.g. screams, slapping of the ground, grabbing at the food). In this case, it is shown that food sharing might be the optimal choice for the food owner in situations where the fitness cost caused by the beggar's harassment, if the food is defended, exceeds the fitness cost of sharing.

In this paper, we extend the model of kleptoparasitism presented in Broom et al. (2004) by assuming divisible food items and allowing animals to share their prey with attacking foraging animals. A foraging animal, encountering an animal handling a food item has the possibility to either attack attempting to steal or share the food, or just ignore it and continue foraging. On the other hand, an attacked animal which owns a food item, has the possibility to defend its food, to share it or to retreat leaving all the food to the attacking animal. Through a game-theoretic approach we examine the optimal strategy for an animal under different ecological circumstances.

The model

In a population of foragers of density P , each animal might either be in a state of searching for food, or a state where it is handling a food item that it has acquired. Let S denote the density of searchers and H the density of handlers. Each handler consumes the food item and resumes searching in a time drawn from an exponential distribution with mean t_h , so equivalently following a Markov process at rate t_h^{-1} . There is a constant density of food items f available and searchers cover an area ν_f per unit time whilst searching for food, so that they find food at rate $\nu_f f$. As well as finding food themselves when foraging, searchers can acquire food by trying to steal it from a handler, and they can search an area ν_h per unit time for handlers. Once a searcher comes upon a handler, it attacks to either steal or share the food item with probability p_1 or ignores the handler with probability $1 - p_1$ and continues searching for food. If the searcher attacks, the handler might decide to resist and defend its food item. This happens with probability

146 which we label p_3 . In this case, the attacking searcher (A) and the defender (R) engage in
 147 a fight. The rate at which searchers encounter handlers and engage in a fight (become
 148 attackers, A) is equal to $p_1 p_3 \nu_h H$ while handlers are found by searchers and resist a
 149 possible attack (become defenders, R) with rate $p_1 p_3 \nu_h S$. The fight lasts for a time
 150 drawn from an exponential distribution with mean $t_a/2$. The attacker animal wins the
 151 fight and becomes a handler with probability α and thus, with the same probability,
 152 the defender loses its food and starts searching again; so this happens at rate $2\alpha/t_a$.
 153 Otherwise, the attacking searcher loses the fight and returns to the searching state with
 154 rate $2(1-\alpha)/t_a$ and thus, with the same rate, the defender wins and continues handling
 155 its food. Note that the winner of the fight might face other subsequent challenges. In
 156 general, the circumstances under which fights occur might give a high advantage to
 157 defender or attacker (the attacker might have to catch the defender in the air, but the
 158 defender may be hampered by a heavy food item) and so α may be significantly greater
 159 or less than 0.5.

160 So far, the model described is the same as the model investigated in Broom et al.
 161 (2004). In this paper, this model is extended by assuming that attacked animals can
 162 share a food item as follows. Assume that food items are divisible. The attacked handler
 163 might decide to share its food with an attacking searcher, with probability p_2 . In this
 164 case, searchers become sharers (C) with rate $p_1 p_2 \nu_h H$ and the attacked handlers with
 165 rate $p_1 p_2 \nu_h S$. If the handler decides to share its food with the searcher, both take a
 166 half of the food. It is assumed, for reasons of simplicity, that both the two sharers
 167 hold the food item and feed simultaneously on it. This discourages other animals from
 168 attempting to steal or share the food since this would be a difficult, risky and dangerous
 169 venture. So, food sharing results in the mutual protection of the two sharers from other
 170 predators. As a result, a sharer animal consumes its portion of the food item without
 171 any interruptions. Sharers eat their food unperturbed and again become searchers in a
 172 time drawn from an exponential distribution with mean t_c or equivalently with rate t_c^{-1} .
 173 Once the half of the food item is consumed, the sharer starts foraging again. Throughout
 174 the paper, it is assumed that $2t_c \geq t_h$, i.e. the decision of food sharing might either
 175 have no time cost or has some cost, but is never beneficial with respect to the handling

time.

The attacked handler, in order to avoid any time cost either from a fight or the sharing process, might decide neither to defend its food item nor to share it, but to leave it to the attacking animal and return to the searching state. This happens with probability $1 - p_2 - p_3$ for any challenge, and so occurs at rate $p_1(1 - p_2 - p_3)\nu_h H$ for each searcher and rate $p_1(1 - p_2 - p_3)\nu_h S$ for each handler.

The model parameters and notations are summarised in Table 1.

The differential equation based compartmental model that describes the dynamic of the different groups of the population in the above situation is the following

$$\frac{dS}{dt} = \frac{1}{t_h}H + \frac{1}{t_c}C + \frac{2}{t_a}(1 - \alpha)A + \frac{2}{t_a}\alpha R - \nu_f f S - p_1(p_2 + p_3)\nu_h SH, \quad (1)$$

$$\frac{dH}{dt} = \nu_f f S + \frac{2}{t_a}\alpha A + \frac{2}{t_a}(1 - \alpha)R - \frac{1}{t_h}H - p_1(p_2 + p_3)\nu_h SH, \quad (2)$$

$$\frac{dC}{dt} = 2p_1p_2\nu_h SH - \frac{1}{t_c}C, \quad (3)$$

$$\frac{dA}{dt} = p_1p_3\nu_h SH - \frac{2}{t_a}A, \quad (4)$$

$$\frac{dR}{dt} = p_1p_3\nu_h SH - \frac{2}{t_a}R. \quad (5)$$

The above system of equations is a closed system where the population density, P , remains constant, i.e.

$$P = S + H + C + A + R, \quad (6)$$

and one of the equations (1)–(5) is thus redundant.

We assume that the population rapidly converges to the equilibrium state (see Luther and Broom (2004) for a proof of this assumption for the original model of Broom and Ruxton (1998)). In the equilibrium conditions, the densities of the different groups of the population, S, H, C, A and R , are given by (see Appendix A)

$$(S, H, C, A, R) = \left(\frac{H}{t_h d(H, p_1, p_2)}, H, \frac{2p_1p_2t_c\nu_h H^2}{t_h d(H, p_1, p_2)}, \frac{1}{2} \frac{p_1p_3t_a\nu_h H^2}{t_h d(H, p_1, p_2)}, \frac{1}{2} \frac{p_1p_3t_a\nu_h H^2}{t_h d(H, p_1, p_2)} \right), \quad (7)$$

where $d(H, p_1, p_2) = \nu_f f - p_1p_2\nu_h H$, i.e. the difference between the rate at which searchers discover food items and the rate at which they become sharers. Note that this

term is clearly positive since every food item can be shared at most once (and some are not shared), and it must be discovered beforehand. By (6) and (7), H is given by the biologically relevant solution of the quadratic equation

$$p_1(p_2(2t_c - t_h) + p_3t_a)\nu_h H^2 + (p_1p_2t_h\nu_h P + t_h\nu_f f + 1)H - t_h\nu_f f P = 0, \quad (8)$$

i.e. the positive solution,

$$H = \frac{-(p_1p_2t_h\nu_h P + t_h\nu_f f + 1) + \sqrt{(p_1p_2t_h\nu_h P + t_h\nu_f f + 1)^2 + 4p_1t_h\nu_f f \nu_h P(p_2(2t_c - t_h) + p_3t_a)}}{2p_1\nu_h(p_2(2t_c - t_h) + p_3t_a)}, \quad (9)$$

given that $2p_1\nu_h(p_2(2t_c - t_h) + p_3t_a) > 0$.

Optimal strategies

We are interested in finding conditions under which animals playing strategy (p_1, p_2, p_3) , i.e. animals which attack handlers with probability p_1 and share or defend their food when they are attacked with probability p_2 and p_3 , respectively, have greater fitness than animals playing any other strategy (q_1, q_2, q_3) . We are ultimately looking for conditions when the overall strategy (p_1, p_2, p_3) is an ESS.

A strategy is considered to be the optimal strategy if it minimizes the average time needed to the consumption of a food item. This would result in the maximising of the long-term food intake rate of an animal playing this strategy and thus its fitness.

Average time for a single animal to consume a food item

Assume that a mutant animal playing strategy (q_1, q_2, q_3) invades into a population playing strategy (p_1, p_2, p_3) .

If the mutant is at the searching state and encounters a handler it has two options:

- It attacks in order to share or steal the food item with probability q_1 . Note that once it attacks, what will happen next depends on the handler's strategy.
- It ignores the handler animal and continues searching for a food item for itself with probability $1 - q_1$.

216 The time needed for the mutant searcher, who has just come upon a handler playing the
 217 population strategy, to consume a food item, T_{SA}^* , in the different scenarios is represented
 218 schematically in the diagram shown in Figure 1. The notation of food consumption times
 219 from the different foraging states is shown in Table 2.

220 If the mutant is at the handling state and is attacked by a searcher animal playing the
 221 population strategy it has three options:

- 222 - It shares the food item with probability q_2 .
- 223 - It defends its food and a fight takes place with probability q_3 .
- 224 - It leaves the food to the attacker and resumes searching with probability $1 - q_2 - q_3$.

The time required for the attacked mutant handler to consume a food item, T_{HA}^* , in the different scenarios is represented schematically in the diagram shown in Figure 2.

It is shown (see Appendix B) that T_{SA}^* and T_{HA}^* are given by the solution of the following system of equations

$$\begin{aligned} & \left(1 - \left(1 - q_1 + \frac{q_1 p_2}{2} + (1 - \alpha) q_1 p_3 \right) \frac{\nu_h H}{\nu_f f + \nu_h H} \right) T_{SA}^* = q_1 p_2 t_c + q_1 p_3 \frac{t_a}{2} + \\ & + \left(1 - q_1 + \frac{q_1 p_2}{2} + (1 - \alpha) q_1 p_3 \right) \frac{1}{\nu_f f + \nu_h H} + \\ & + \left(\left(1 - \frac{q_1 p_2}{2} \right) \nu_f f + q_1 (1 - p_2 - (1 - \alpha) p_3) \nu_h H \right) \frac{t_h (1 + \nu_h S T_{HA}^*)}{(1 + t_h \nu_h S)(\nu_f f + \nu_h H)}. \end{aligned} \quad (10)$$

$$\begin{aligned} & \left(1 - \frac{\left(\left(1 - \frac{p_1 q_2}{2} \right) \nu_f f + (1 - p_1 + (1 - \alpha) p_1 q_3) \nu_h H \right) t_h \nu_h S}{(1 + t_h \nu_h S)(\nu_f f + \nu_h H)} \right) T_{HA}^* = p_1 q_2 t_c + p_1 q_3 \frac{t_a}{2} + \\ & + \left(\left(1 - \frac{p_1 q_2}{2} \right) \nu_f f + (1 - p_1 + (1 - \alpha) p_1 q_3) \nu_h H \right) \frac{t_h}{(1 + t_h \nu_h S)(\nu_f f + \nu_h H)} + \\ & + p_1 \left(1 - \frac{q_2}{2} - (1 - \alpha) q_3 \right) \frac{1 + \nu_h H T_{SA}^*}{\nu_f f + \nu_h H}. \end{aligned} \quad (11)$$

225 The average required time to the consumption of a food item for a single searcher animal
 226 of a population playing strategy (p_1, p_2, p_3) , who has just met a handler animal of this
 227 population, T_{SA} , and the respective time of a single handler of the same population who
 228 has just met a searcher, T_{HA} , can be found by solving the system of equations (10) and
 229 (11) substituting (p_1, p_2, p_3) for (q_1, q_2, q_3) .

In the case where all the members of the population do not challenge, i.e. $p_1 = q_1 = 0$, the strategy used by an animal in the handling position may be thought irrelevant since none of the animals will ever be attacked and thus each searcher finds a food item for itself in an average time equal to $1/\nu_f f$ and each handler consumes a discovered food item in time t_h . However, we assume that occasionally a challenge occurs “by mistake” (this is a version of the classical trembling hand argument of Selten (1975)). Thus, a handler animal of a population where animals never challenge, at some point is faced by a foraging animal which attempts to steal or share the food. In this case, the average time needed for the attacked handler animal to consume a food item if it adopts a different from the population strategy, $(0, q_2, q_3)$, is given by (see Appendix B)

$$T_{\text{HA}}^* = q_2 \left(t_c - \frac{1}{2} \left(\frac{1}{\nu_f f} + t_h \right) \right) + q_3 \left(\frac{t_a}{2} - (1 - \alpha) \frac{1}{\nu_f f} \right) + \frac{1}{\nu_f f} + t_h. \quad (12)$$

230 If a mutant animal can invade a population then its strategy (q_1, q_2, q_3) is a better
 231 strategy than that of the population (p_1, p_2, p_3) at least at one of the two decision points,
 232 when a searcher and potentially making a challenge or when receiving a challenge as
 233 a handler. A mutant which follows a different strategy from that of the population at
 234 just one decision point and the strategy which is followed is better than that of the
 235 population, can obviously invade. When considering whether a particular strategy is an
 236 ESS or not, it is sufficient to investigate invasion by mutants which differ in strategy at
 237 one of the two decision points only. This is because if a mutant which differ in strategy
 238 at both of the decision points can invade, it must have a superior strategy at at least one
 239 of the decision points, and so an animal which shares the same strategy as the mutant
 240 at this decision point, and the same strategy as the population at the other, could also
 241 invade.

242 A mutant which uses a strategy different from that of the population at just the
 243 searching state is considered to use a better strategy, and thus be able to invade, if
 244 $T_{\text{SA}}^* \leq T_{\text{SA}}$, i.e. if the decision that it will make at the point when it will meet a handler,
 245 when searching for food, will lead to a smaller time until the consumption of a food item.
 246 Similarly, a mutant which plays differently from the population just at the handling state
 247 is considered to be able to invade if the decision it will make in an encounter with a

searcher, when handling a food item, will shorten the time to the consumption of a food item, i.e. if $T_{HA}^* \leq T_{HA}$. Note that it is possible that under certain parameters T_{SA}^* is independent of q_1 and all values $0 \leq q_1 \leq 1$ give identical times. Similarly, T_{HA}^* might be independent of q_2 and q_3 . In these circumstances, in such asymmetric games, the population can still be invaded by genetic drift.

In Appendix C, it is demonstrated through a combination of analytical and numerical investigation that the optimal strategy in either decision points is always pure (in some cases this is proved, in others it is not proved but an extensive numerical investigation has yielded consistent results with no mixed ESSs). Thus, if the population plays a non-pure strategy (p_1, p_2, p_3) , for an invading animal there will be a pure strategy that will do at least as well as playing the population strategy, and so (p_1, p_2, p_3) could not be an ESS since this pure strategy would invade the population. Hence, we need to consider only two strategies for a foraging animal (always or never attempt to steal or share the prey of the other animal when the opportunities arise) and three strategic choices for the attacked animal (either always surrender the food to the attacking animal, always share the food or always defend it) as the components of the potential optimal strategy in any given population. Therefore, there are six possible pure strategies that an animal can use and need to be considered:

- Strategy (0,0,0) (Dove, D): the animal does never challenge handlers and does never resist any challenges.
- Strategy (0,1,0) (Non-Attacking Sharer, NAS): the animal does never challenge handlers and always shares its food when it is challenged.
- Strategy (0,0,1) (Retaliator, R): the animal does never challenge handlers but always resists when it is challenged.
- Strategy (1,0,0) (Marauder, M): the animal challenges handlers at every opportunity but it does never resist any challenges.
- Strategy (1,1,0) (Attacking Sharer, AS): the animal challenges handlers at every opportunity and it always shares the food when it is challenged.

- Strategy (1,0,1) (Hawk, H): the animal challenges handlers at every opportunity and it always resists any challenges.

The optimal strategy for an animal at the searching state

Consider a population playing strategy (p_1, p_2, p_3) that is potentially invaded by a mutant animal playing a different strategy (q_1, q_2, q_3) . For reasons explained in the previous section, in order to study whether the mutant can evolve because it uses a better strategy at the searching state, we assume that the strategy which is used by all the animals when they are at the handling state is the same, i.e. $p_2 = q_2$ and $p_3 = q_3$. We consider the strategy used by a searcher animal of the population when coming across a handler, p_1 , to be advantageous over a mutant strategy, q_1 , (and thus the population cannot be invaded by the mutant) if the average time required for the searcher playing the population strategy to gain and consume a food item, T_{SA} , is less than that required for the mutant searcher, T_{SA}^* . Using the equations (10), (11) and (7)–(9) we find all the necessary conditions under which a mutant playing strategy $q_1 \in \{0, 1 : q_1 \neq p_1\}$ cannot invade a population playing strategy $p_1 \in \{0, 1 : p_1 \neq q_1\}$ for the cases where either $p_2 = q_2 = 0$ and $p_3 = q_3 = 1$, $p_2 = q_2 = 1$ and $p_3 = q_3 = 0$ or $p_2 = q_2 = 0$ and $p_3 = q_3 = 0$. These are summarised in Table 3 (conditions (C.3), (C.6), (C.9), (C.10), (C.13) and (C.16)).

The optimal strategy for an animal at the handling state

In the handling position an animal can use three strategies when it is challenged. Either it shares the food with the challenger, it defends its food or it retreats leaving the food to the attacking animal, and depending on the ecological conditions it obtains the highest benefit when it always takes one of these three actions. As before, assume that a population already at equilibrium conditions is invaded by a mutant, which now uses a different strategy as a handler but the same strategy as a searcher.

301 **Optimal strategies in an aggressive population**

302 Assume that all the members of the population behave aggressively when encountering
 303 a handler animal, i.e. $p_1 = q_1 = 1$. We consider the strategy of an attacked handler
 304 of the population to be advantageous over the strategy used by an attacked handler
 305 mutant (and thus the mutant cannot invade) if the average time required for the first to
 306 consume a food item, T_{HA} , is less than that required for the second, T_{HA}^* (in this case,
 307 this is equivalent to the comparison of T_{SA} with T_{SA}^* since the times needed for animals
 308 which always challenge, i.e. when $p_1 = q_1 = 1$, to acquire a food item and be discovered
 309 by a foraging animal are identical, independently of the strategies they use as handlers).
 310 Using again the equations (10), (11) and (7)–(9) we find the necessary conditions under
 311 which a mutant in this scenario cannot invade a population playing a different strategy
 312 at the handling state. These conditions are presented in Table 3 (conditions (C.11),
 313 (C.12), (C.14), (C.15), (C.17) and (C.18)).

314 **Optimal strategies in a non-aggressive population**

315 In the case where all the members of the population do not challenge, i.e. $p_1 = q_1 = 0$, an
 316 animal of the population playing $(0, p_2, p_3)$ does better than a mutant playing $(0, q_2, q_3)$,
 317 and thus the population cannot be invaded by this mutant, if $T_{HA} < T_{HA}^*$, where by
 318 (12) (T_{HA} in this case is similarly given by (12) substituting p_2 and p_3 for q_2 and q_3 ,
 319 respectively) we obtain the condition

$$(q_2 - p_2) \left(t_c - \frac{1}{2} \left(\frac{1}{\nu_{ff}} + t_h \right) \right) + (q_3 - p_3) \left(\frac{t_a}{2} - (1 - \alpha) \frac{1}{\nu_{ff}} \right) > 0. \quad (13)$$

320 The conditions under which a mutant playing strategy $(0, q_2, q_3)$ is unable to invade
 321 a population playing strategy $(0, p_2, p_3)$ are summarised in Table 3 (conditions (C.1),
 322 (C.2), (C.4), (C.5), (C.7), (C.8)).

323 **Evolutionarily Stable Strategies**

324 Table 3 shows all the appropriate conditions under which a population playing strategy
 325 (p_1, p_2, p_3) cannot be invaded by a mutant playing a different strategy at one of the

two decision points, (q_1, q_2, q_3) , for all the possible cases where all animals play a pure strategy.

According to the results shown in Table 3, strategies (0,0,0) and (0,1,0) can never resist all of the possible invading strategies and there are thus four possible ESSs:

- Strategy (0,0,1) is an ESS if the conditions (C.7), (C.8) and (C.9) are satisfied.
- Strategy (1,0,0) is an ESS if the conditions (C.11) and (C.12) are satisfied.
- Strategy (1,1,0) is an ESS if the conditions (C.13), (C.14) and (C.15) are satisfied.
- Strategy (1,0,1) is an ESS if the conditions (C.16), (C.17) and (C.18) are satisfied.

Figure 3 shows the regions in parameter space in which each of the four strategies, Retaliator, Marauder, Attacking Sharer and Hawk, is an ESS, for specific parameter values as the duration of the contest, $t_a/2$, and the handling time of a sharer, t_c , vary. Figure 4 shows how these regions vary as the density of the population, P , and the rate at which foragers find undiscovered food, $\nu_f f$, vary. Obviously, these regions in the t_a, t_c plane in Figure 3 and $P, \nu_f f$ plane in Figure 4 will vary, depending on the other parameter values. However, some general conclusions can be extracted. Figure 3 and Figure 4 suggests that between the regions where two strategies are unique ESSs, there can be a region where the two strategies are simultaneous ESSs and among the regions of three pairs of ESSs configured by three strategies, there might be a region where the three strategies might coexist as ESSs. This excludes the possibility of the Retaliator and the Hawk strategies being simultaneous ESSs, since this can never happen due to the contradiction of the conditions (C.9) and (C.16) (see Table 3). This gives eleven distinct regions as summarised in Figure 3 and Figure 4. It appears that every set of parameters yields one or more pure ESSs. Numerical examples on a wide range of parameter values indicate that there is no parameter set where this is not the case i.e. that there are not any mixtures of strategies or cases where there are no ESSs. Although we do not believe that there will be any parameter set where there will be such a polymorphic mixture or no ESS (in similar models such cases do not occur, and see Appendix C for an argument that actual mixed strategy ESSs are not possible), we cannot definitively rule out this possibility.

Predictions of the model

In the case where neither the members of the population nor any mutant share the food, i.e. in the case where $p_2 = q_2 = 0$, all the above results agree with the results obtained in previous work (Broom et al., 2004). Hence, here we concentrate on the cases where the members of the population or a mutant animal or both, always share their food when they are attacked, i.e. cases where either p_2 or q_2 or both are equal to 1. This provides both new potential ESSs and also new mutant strategies to invade other strategies, so that strategies that were ESSs in Broom et al. (2004) will no longer be in some cases.

In a non-attacking population, a sharer does better than a Dove when they are attacked if the average time needed for a sharer to consume a whole food item ($t_c + ((1/\nu_f f) + t_h)/2$) is less than the average time needed to find an undiscovered food item ($1/\nu_f f$) and consume it (t_h) (equivalently in this case, if the time the sharer needs to consume the half of the food item (t_c) is on average less than half of the time needed to find and consume a whole food item $((1/\nu_f f) + t_h)/2$). On the other hand, an Attacking Sharer mutant does better than a member of a population of Non-Attacking Sharers if $t_c \leq ((1/\nu_f f) + t_h)/2$ as well. Hence, as we see in Table 3, condition (C.4) contradicts condition (C.6) and thus a Non-Attacking Sharer is never an ESS. The food sharing strategy can be an ESS only if the sharer challenges a handler at every opportunity when it is at the searching state. A population of Attacking Sharers can potentially be invaded by Non-Attacking Sharers, Marauders and Hawks. The conditions under which a Non-Attacking Sharer and a Marauder can invade a population of Attacking Sharers are the same. This occurs because in such a population a Marauder can invade if it is better for any handler to give up a food item rather than share (so being a searcher is better than sharing a food item) and a Non-Attacking Sharer can invade if it is better not to challenge for a food item, which will be shared (so again searching is better than sharing). Increasing the rate at which foragers find food, $\nu_f f$, increases the parameter range where Non-Attacking Sharers and Marauders invade the population of Attacking Sharers. Depending on the values of the other parameters, the increase of $\nu_f f$ might favour the invasion of Hawks as well (usually when food is difficult to discover). Hence, increasing $\nu_f f$ decreases the range of the parameter values in which the Attacking

Sharer strategy is an ESS (see Figure 3 for an example). A similar situation appears by decreasing the area in which foragers search for handling sharers per unit time, ν_h . As it is observed in Figure 4 and the conditions (C.13)–(C.15), the decrease of the density of the population, P , might also create unpropitious circumstances for food sharing. For a given set of parameter values for which the Attacking Sharer strategy is an ESS, increasing the time cost of the sharing process which results in the increase of t_c , the area where the Attacking Sharer strategy is an ESS reduces, as one would expect. Depending on the other ecological conditions, this strategy might coexist as an ESS with either one of the other possible ESSs (Retaliator, Marauder or Hawk) or two of them (Retaliator and Marauder or Marauder and Hawk). At very high levels of t_c such that the time spent in sharing would be better spent in searching for another food item or in defending the food item, Attacking Sharer cannot be an ESS. In this case, the predictions of the model approach those of the model of Broom et al. (2004), where sharing was not possible. In conditions where the duration of aggressive interactions is high, the defending strategy is less profitable and thus the avoidance of any aggressive interaction is favoured. Hence, under these circumstances, it is observed that animals should decide either to surrender their food (use the Marauder strategy) or to share it (use the Attacking Sharer strategy) when they are challenged, even if they have a high probability of defending their food successfully. Therefore, at high fight durations each of Marauder and Attacking Sharer strategies might be the unique ESS or both might be ESSs simultaneously.

A special case

As a special case, we consider the case where $2t_c = t_h$, i.e. where sharing does not reduce the speed of food consumption. The results obtained in this case are shown in Table 4. It is observed that, as well as the Dove and Non-Attacking Sharer strategies which as we have seen in the previous section are never ESSs, in this case the Marauder strategy is also never an ESS since it can always be invaded by an Attacking Sharer animal. The Attacking Sharer strategy can only be invaded by the Hawk strategy. Moreover, this can happen just in few cases where the chance of a successful defence is relatively high, i.e. the probability α is relatively small, and the time spent in a contest, $t_a/2$, is small. For

$\alpha \geq 0.5$, the conditions (C.7) and (C.9) indicate that the Retaliator strategy can never be an ESS. In this case, the condition (C.18) also indicates that an Attacking Sharer can always invade a population playing Hawk and thus the Hawk strategy can never be an ESS as well. Hence, at least for $\alpha \geq 0.5$, Attacking Sharer is the only ESS no matter what the other parameter values are. The Hawk strategy is an ESS mainly when $t_a/2$ and α are small. As $t_a/2$ and/or α increase, depending on the other parameter values, there might be a range where Hawks and Attacking Sharers coexist as ESSs. When the defender is likely to succeed, i.e. α is small, defence of the food item might be the favoured strategy even if the fight time is relatively long, especially in cases where available food is scarce. Hence, there is a range where the Retaliator strategy is either the only ESS or coexists with the Attacking Sharer strategy. Figure 5 shows a region with all the possible ESSs in this specific case, as the probability α of the challenger winning and the duration of the content, $t_a/2$, vary.

Discussion

Food sharing is a very common tactic adopted by a broad group of animal species for their survival. Using a game theoretic approach, the present model investigates the ecological circumstances under which animals should share their food when they are challenged by other foraging animals. We have extended the game-theoretical model of Broom et al. (2004) by allowing animals to share their food. Hence, animals in this model can choose among two additional strategies: either to attempt to share or steal the food from a handler when foraging and share their food when they are challenged by a forager, or to ignore any opportunities to share or to steal food when foraging but share when another animal attacks. This model is likely to be an improvement if caught food items are at least partly divisible, for instance fruit species (e.g., White, 1994), as opposed to for example a nut or a fish (e.g., Iyengar, 2008) which are hard to divide, in which case the original modelling system will be more appropriate. At the opposite extreme, in situations where food items come in patches, for instance seed patches (e.g., Barnard and Sibly, 1981), which are easily divisible, then the producer-scrounger type models (e.g., Dubois and Giraldeau, 2003; Dubois et al., 2003; Dubois and Giraldeau,

2005, 2007) could be appropriate models.

Considering the time cost needed for a food item to be acquired and consumed, the model predicts that there is a wide range of ecological conditions in which attempting to share or steal the food at every opportunity and sharing the food when attacked is the optimal strategy that should be used by animals. The non-aggressive strategy where animals do not challenge other animals but share their food when challenged can never be an ESS since depending on the ecological parameters this strategy is always invaded either by the Dove or the Attacking Sharer strategy. This adds one possible ESS to the model of Broom et al. (2004). Investigation of the model suggests that under any ecological parameters, there is always at least one ESS that an animal can use. Every two ESSs can occur as ESSs simultaneously, apart from the Retaliator and the Hawk strategy where it is shown that they can never be ESSs simultaneously. It is also possible that under some conditions there are three simultaneous ESSs (Retaliator, Marauder and Attacking Sharer or Marauder, Attacking Sharer and Hawk).

Different ecological factors might influence the strategic choice of food sharing. Food availability is one of the crucial factors. In conditions of limited food, the use of the Attacking Sharer strategy is enhanced while at high food densities, food sharing becomes a less profitable strategy. A high time cost of food defence, a small probability of a successful food defence, a high rate at which searchers encounter handlers, a high population density and a low time cost of food sharing are also conditions which favour animals sharing their food. In the special case where food sharing has no additional time cost, foraging animals should almost always attempt to share food with a handler and handlers should almost always share their food. Defending the food might be the optimal strategy for the owner, especially when food is difficult to be discovered, and the success of this is likely. Moreover, attacking at every opportunity and defending when attacked is an ESS in just very few cases, where the time cost of the defence is small, but never attack and always defend might be an ESS even if the defence will result in a high time cost. Attack and always retreat when attacked never occurs in this case since sharing is always a better strategy.

Food sharing is a complicated mechanism. Different animal species share their food

473 for different reasons and under different ecological and biological conditions. In many
474 situations, food sharing is a voluntary process where animals choose to share their
475 food without any kind of menace from other foraging animals coming before. This
476 process might result to immediate benefits for animals, for example the creation of
477 cooperation for the increase of foraging success or predation avoidance, or to increase
478 mating opportunities (see Stevens and Gilby, 2004). It is also often the case that sharing
479 occurs between relatives or between animals with a social interaction, even if those
480 animals are not relatives e.g. between roost mates (Wilkinson, 1990). As a result, food
481 sharing might not be immediately beneficial but result in long term benefits such as
482 future reciprocal sharing, i.e. altruism (see Stevens and Gilby, 2004). In the present
483 model, food sharing is considered to be the process where a food owner shows tolerance
484 to an attacking animal and shares its food with it, although it would be better for
485 the owner not to be discovered by any other animal. This animal behaviour might
486 occur in cases where a beggar challenges a food owner, a situation which is observed
487 in monkeys and chimpanzees populations (see Stevens and Gilby (2004) for examples
488 of this behaviour). The particularity of this model compared to other models in the
489 literature, is that by sharing food, the two animals protect each other from potential
490 subsequent costly challenges that might extend the time until the consumption of a food
491 item. Hence, on average a half of the food item is consumed without the risk of other
492 delays apart from the time required for sharing. This, under certain conditions, might
493 be the least costly process with respect to the expected time needed for the consumption
494 of food and thus a process which maximises the food intake rate. Although there is no
495 empirical data to support precisely the above assumptions, there is evidence that in
496 nature, animals in many cases prefer to share food with other animals to reduce the risk
497 of losing the entire prey. For example, a lion instead of defending its prey against an
498 approaching member of the pride, it might share it in order to increase the efficiency of
499 defending the prey from invading hyenas (see for example, Cooper, 1991; Stevens and
500 Gilby, 2004).

501 In addition, our model assumes that the members of the population are of the same
502 type. However, real populations consist of individuals with biological and physiological

503 differences and the optimal strategic choices depend on the characteristics of the individ-
504 uals and those of their opponent. For example, recent observational and experimental
505 studies on the dung roller beetle *Canthon cyanellus cyanellus* have shown that males of
506 similar size are more likely to share the resource rather than to defend it (Chamorro-
507 Florescano et al., 2010). Fight duration may be correlated with the differences between
508 the opponents as well (e.g., Rovero et al., 2000). The size and the quality of the food
509 items or the estimation of the value of the resource might also affect significantly the
510 frequency of food sharing (see for example, White, 1994) as well as a contest duration
511 (see for example, Enquist and Leimar, 1987).

512 In our model all costs are expressed in terms of time used and we ignore other costs
513 which can be important, such as energy costs and possible injuries resulting from fights
514 (for a model which incorporates energy costs see Vahl, 2006). For simplicity we do
515 not impose extra time penalties on animals in contests. A resulting limitation is that
516 the winner and the loser of a contest face the same cost. Although this can be the
517 case in nature (e.g., Smith and Taylor, 1993), experimental studies have shown that
518 either the loser (e.g., Chellapa and Hungtingford, 1989; Neat et al., 1998) or the winner
519 (e.g., Hack, 1997) might suffer higher energetic or other cost, such as a high recovery
520 time cost (see also, Luther and Broom, 2004). For instance, if the handler uses more
521 energy (e.g. because it is carrying a food item during the contest) then it might need
522 a higher recovery time. This would decrease the food intake rate making the defending
523 strategy less attractive and the choice of alternative strategies more likely. In the same
524 way, although it is assumed that the cost from the sharing process is equal for the
525 two animals that share food, in reality the two animals might suffer a different cost.
526 Furthermore, it is assumed that once an animal loses a contest with another animal, it
527 does not initiate a new fight with the same animal but starts searching for alternative
528 food resources. This is generally reasonable, as often contests between animals can have
529 strong (at least short-term) effects on their relationship (winner and loser effects) which
530 reinforce the dominance of the winner (see for example, Dugatkin, 1997). Similarly, in
531 related contests between animals for territory acquisition animals that lose an agonistic
532 interaction often leave the areas in which they were defeated (see for example, Stamps

533 and Krishnan, 1994). However, we should note that in some cases an animal may attack
534 repeatedly the owner after iterated losing tries (e.g., Stamps, 1994). One way that
535 the model could be extended and relax this assumption, is to allow the loser animal
536 to attack repeatedly the winner. This could be done, for example, by introducing the
537 choice to the loser to attack again or not, following similar assumptions to those in some
538 owner-intruder type of games (e.g., Morrell and Kokko, 2003).

539 In natural systems, foraging animals might be faced with more than one foraging
540 option with different variance in food intake. For example, they might be faced with a
541 constant food resource versus a variable food resource, a food resource with fixed delay
542 versus the same food resource with variable delay, or an immediate gain of food versus
543 a delayed gain. There is strong empirical evidence that a forager's choice may depend
544 on many ecological factors, such as the energetic status of the animal, the type of food
545 variance, the energy requirements of the animal within a certain time interval and the
546 probability of delays due to different kind of unpredictable interruptions (for example
547 bad weather). The forager might be either risk-averse and choose the predictable option,
548 or risk-prone and choose a risky option, respectively (see for example, Kacelnik and
549 Bateson, 1996). For example, an animal with low food reserves might choose a safe
550 lower level of return, provided it was sufficient for survival. Food sharing might be
551 a way for animals to reduce such variances in food intake (e.g., Wenzel and Pickering,
552 1991). Although the present model does not consider any risk associated with alternative
553 food sources, it would be interesting to incorporate in future work such parameters that
554 might influence the foraging decisions.

555 Further research taking into consideration all these different factors will help us to
556 better understand the reasons why and the conditions under which animals prefer to
557 share their food.

Appendix A

The densities of the different groups of the population, S, H, C, A and R , in the equilibrium conditions

In the equilibrium conditions

$$\frac{dS}{dt} = \frac{dH}{dt} = \frac{dC}{dt} = \frac{dA}{dt} = \frac{dR}{dt} = 0. \quad (14)$$

From the equation

$$\frac{dC}{dt} = 2p_1p_2\nu_hSH - \frac{1}{t_c}C = 0, \quad (15)$$

it follows that in the equilibrium, the number of sharers is given by

$$C = 2p_1p_2t_c\nu_hSH. \quad (16)$$

Similarly, from the equations

$$\frac{dA}{dt} = \frac{dR}{dt} = 0, \quad (17)$$

it is derived that the number of attackers and defenders in the equilibrium is given by

$$A = R = \frac{p_1p_3t_a\nu_hSH}{2}. \quad (18)$$

Substituting equations (16) and (18) into the system of equations

$$\frac{dS}{dt} = \frac{dH}{dt} = 0, \quad (19)$$

and solving the system for S using equation (6), it is obtained that in the equilibrium, the densities of the different groups of the population, S, H, C, A and R , are given by (7).

Appendix B

Average time for a single animal to consume a food item

Average time for a single searcher animal to consume a food item when encountering a handler animal

Assume that a mutant searcher playing (q_1, q_2, q_3) has just come upon a handler playing the population strategy, (p_1, p_2, p_3) . If the mutant searcher ignores the handler, with probability $1 - q_1$, then it will need an average time T_S^* until the consumption of a food item. Otherwise, if the mutant attacks, with probability q_1 , the average time needed for the consumption of a food item depends on the action that the handler animal will take. If the handler decides to share the food, with probability p_2 , then the further expected time required to the consumption of a whole food item by the mutant is T_C^* . If the attacked handler decides to defend its food, with probability p_3 , then a fight takes place and the attacking mutant will need an average time T_A^* to acquire and consume a food item. Finally, if the attacked animal decides to leave its food to the attacking animal without taking any action, with probability $1 - p_2 - p_3$, the attacking searcher animal becomes a handler and it then requires an average time T_H^* until the consumption of a food item. T_{SA}^* is given by the following equation

$$T_{SA}^* = q_1(p_2T_C^* + p_3T_A^* + (1 - p_2 - p_3)T_H^*) + (1 - q_1)T_S^*. \quad (20)$$

Recall that we assume that two animals that share a food item do so equally. Each of the sharers needs a time t_c until consumption of the half of the food and once it consumes it, it returns to the searching state. From the searching state, the mutant needs a time on average equal to T_S^* in order to consume a whole food item. The average time needed for a mutant sharer to consume a whole food item, T_C^* , is given by

$$T_C^* = t_c + \frac{T_S^*}{2}. \quad (21)$$

The sharing process described above is, in terms of expected reward, entirely equivalent

to a process where if a searcher and a handler decide to share the food, at the end of the sharing period, with probability 0.5 one of the two animals obtains the food item while the other takes nothing. The loser then has to resume searching for a new food resource and thus spend an average time T_S^* until the consumption of a food item. Both animals suffer a time cost from the sharing process equal to t_c . Hence, the time that a sharer needs for the consumption of a food item is on average equal to $0.5t_c + 0.5(t_c + T_S^*)$, which leads to (21).

Substituting (21) into (20) we obtain

$$T_{SA}^* = q_1(p_2t_c + p_3T_A^* + (1 - p_2 - p_3)T_H^*) + \left(1 - q_1 + \frac{q_1p_2}{2}\right)T_S^*. \quad (22)$$

An attacker animal which has just been involved in a fight will have a cost of an average time $t_a/2$ spent in the contest. With probability $1 - \alpha$ the attacker loses the fight and starts searching again for food while with a complementary probability α , it beats the defender and acquires the food item. Thus, T_A^* is given by the following equation

$$T_A^* = \frac{t_a}{2} + (1 - \alpha)T_S^* + \alpha T_H^*. \quad (23)$$

A searcher animal is looking either for a food resource or a handler animal. At this stage, it spends an average time equal to $1/(\nu_f f + \nu_h H)$ before it finds either an unattended food item (this happens with probability $\nu_f f / (\nu_f f + \nu_h H)$) and becomes a handler or a handler animal (with probability $\nu_h H / (\nu_f f + \nu_h H)$). Thus, T_S^* is given by the following equation

$$T_S^* = \frac{\nu_h H}{\nu_f f + \nu_h H} T_{SA}^* + \frac{\nu_f f}{\nu_f f + \nu_h H} T_H^* + \frac{1}{\nu_f f + \nu_h H}. \quad (24)$$

Once the searcher animal acquires a food item, it either consumes it without being found by any searcher animal, with probability $(1/t_h)/((1/t_h) + \nu_h S)$, or it is discovered by a searcher, with probability $\nu_h S / ((1/t_h) + \nu_h S)$, resulting in an additional expected time cost T_{HA}^* until the consumption of a food item. The average time that the animal is at the handling state before it either consumes its food item or is discovered by a searcher

616 animal is equal to $1/((1/t_h) + \nu_h S)$. T_H^* is thus given by

$$T_H^* = \frac{1}{1 + t_h \nu_h S} 0 + \frac{t_h \nu_h S}{1 + t_h \nu_h S} T_{HA}^* + \frac{t_h}{1 + t_h \nu_h S}. \quad (25)$$

617 Substituting equations (23), (24) and (25) into (22), after some calculations we obtain
 618 equation (10).

619

620 **Average time for a single handler animal to consume a food item when**
 621 **encountering a searcher animal in an aggressive population**

622

623 If a mutant animal at the handling state is attacked by a searcher animal playing
 624 the population strategy, with a non-zero probability ($p_1 \neq 0$), then T_{HA}^* is given by the
 625 following equation

$$T_{HA}^* = p_1 (q_2 T_C^* + q_3 T_R^* + (1 - q_2 - q_3) T_S^*) + (1 - p_1) T_H^*, \quad (26)$$

626 where T_R^* is the average time cost of the decision of a mutant handler to defend its food
 627 against a challenge. Substituting (21) into (26) we obtain

$$T_{HA}^* = p_1 \left(q_2 t_c + q_3 T_R^* + \left(1 - \frac{q_2}{2} - q_3 \right) T_S^* \right) + (1 - p_1) T_H^*. \quad (27)$$

628 In a similar way as before, T_R^* is given by

$$T_R^* = \frac{t_a}{2} + \alpha T_S^* + (1 - \alpha) T_H^*. \quad (28)$$

629 Substituting equations (24), (25) and (28) into (27), we obtain (11).

630

631 **Average time for a single handler animal to consume a food item in a non-**
 632 **aggressive population**

633

634 In the case where all the members of the population do not challenge, i.e. $p_1 = q_1 =$
 635 0, but occasionally a challenge might occur, the average time needed for an attacked

handler mutant playing $(0, q_2, q_3)$ to consume a food item, T_{HA}^* , is given by

$$T_{\text{HA}}^* = q_2 \left(t_c + \frac{T_{\text{S}}^*}{2} \right) + q_3 T_{\text{R}}^* + (1 - q_2 - q_3) T_{\text{S}}^*, \quad (29)$$

where T_{R}^* is given by equation (28). Since the population is not making challenges,

$T_{\text{S}}^* = \frac{1}{\nu_{\text{ff}}} + t_h$ and $T_{\text{H}}^* = t_h$. Substituting into equation (29), we obtain (12).

Appendix C

The optimal strategy is always pure

In the present model, there are 21 possible groups of strategies that an animal can play, 6 of which consist of pure strategies and 15 of mixed strategies. These are summarised in Table 5.

Strategies denoted by (*) in Table 5, are strategies with $p_2 = 0$, that is, strategies where animals never share their food. In this case, the model reduces to the model considered in Broom et al. (2004). In this paper, the authors have shown that the mean time required for a searcher animal that has just encountered a handler to consume a food item is a strictly monotonic function (except with the possible exception of a non-generic parameter set, see below) of the probability with which the searcher attacks the handler, p_1 . Therefore, depending on the parameter values, the searcher animal minimises the time it needs for the consumption of a food item by playing either $p_1 = 0$ or $p_1 = 1$. Any other strategy $0 < p_1 < 1$ results in a higher expected time and thus cannot be evolutionarily stable. Similarly, it has been shown that the average time needed for a handler to consume a food item after being attacked by a searcher is either a strictly increasing or a strictly decreasing function of p_3 and therefore the optimal strategy is always either $p_3 = 0$ or $p_3 = 1$, depending on the parameter values.

Strategies denoted by (**) are the two additional to the Broom et al. (2004) model pure strategies where $p_2 = 1$, i.e. the strategies where animals always share their food when other animals attack. It has been shown in the present paper that under certain conditions one of these can be an ESS, the other not.

In the case where none of the animals of the population behave aggressively, i.e.

662 $p_1 = q_1 = 0$ (strategies denoted by (***) in Table 5), the average time required for an
 663 attacked mutant handler that plays strategy (q_1, q_2, q_3) to consume a food item, T_{HA}^* , is
 664 a function of the form (see equation (12))

$$T_{HA}^* = c_1 q_2 + c_2 q_3 + c_3, \quad (30)$$

665 where c_1 , c_2 and c_3 depend only on the parameters of the model $t_a, t_h, t_c, \nu_f f$ and α .
 666 Hence, if the values of the parameters are such that c_1 and c_2 are both greater than zero,
 667 then the optimal strategy for the mutant is $q_2 = q_3 = 0$. In any other case, if $c_1 < c_2$,
 668 the optimal strategy is $q_2 = 1$ and $q_3 = 0$ while if $c_1 > c_2$ the optimal strategy is $q_2 = 0$
 669 and $q_3 = 1$.

670 It remains to consider whether any of the strategies (S 1)–(S 7) is an ESS. Due to
 671 the complexity of the mathematical formulae, an analytic investigation is very difficult.
 672 Hence, we consider whether each of the remaining strategies is an ESS mainly through
 673 extensive numerical investigation.

674 Regarding strategies (S 1), from equation (22) we get that in a population which
 675 plays strategy $(0 < p_1 < 1, 1, 0)$,

$$T_{SA}^*(0, 1, 0) = T_S^*(0, 1, 0), \quad (31)$$

676 while

$$T_{SA}^*(1, 1, 0) = T_C^*(1, 1, 0) = t_c + \frac{T_S^*(1, 1, 0)}{2}. \quad (32)$$

677 If there is any equilibrium strategy $(p_1^*, 1, 0)$ in (S 1), then $T_{SA}(p_1^*, 1, 0)$ should be equal
 678 to $T_{SA}^*(0, 1, 0)$ and $T_{SA}^*(1, 1, 0)$. But when $T_{SA}^*(0, 1, 0) = T_{SA}^*(1, 1, 0)$ then $T_S^*(0, 1, 0) =$
 679 $T_S^*(1, 1, 0)$. Hence, equating equations (31) and (32) we get

$$T_S^*(0, 1, 0) = T_S^*(1, 1, 0) = T_S(p_1^*, 1, 0) = 2t_c. \quad (33)$$

On the other hand, if the strategy $(p_1^*, 1, 0)$ is an equilibrium strategy, then it cannot be
 invaded by the mutant strategy $(p_1^*, 0, 0)$, i.e. the average required time for the mutant
 handler that has just been attacked in a population which plays strategy $(p_1^*, 1, 0)$,

$T_{HA}^*(p_1^*, 0, 0)$, is higher than the average time required when playing the population strategy, $T_{HA}(p_1^*, 1, 0)$. Using equation (27) we find that

$$T_{HA}^*(p_1^*, 0, 0) > T_{HA}(p_1^*, 1, 0) \quad (34)$$

$$\Rightarrow T_S^*(p_1^*, 0, 0) > T_C(p_1^*, 1, 0) \Rightarrow T_S^*(p_1^*, 0, 0) = T_S(p_1^*, 1, 0) > 2t_c. \quad (35)$$

This contradicts (33). Consequently, there is no any equilibrium strategy ($0 < p_1 < 1, 1, 0$). This is also verified from the results of numerical examples for a wide range of parameter values (see Figure 6a for an example).

In a similar way it is proved that there is no equilibrium strategy in the class of strategies (S 6). If there was an equilibrium strategy ($0 < p_1^* < 1, 0 < p_2^* < 1, 0 < p_3^* < 1, p_2^* + p_3^* < 1$), then $T_{HA}^*(p_1^*, 0, 0)$, $T_{HA}^*(p_1^*, 1, 0)$ and $T_{HA}^*(p_1^*, 0, 1)$ should all be identical, otherwise one of the strategies $(p_1^*, 0, 0)$, $(p_1^*, 1, 0)$, $(p_1^*, 0, 1)$ could invade (p_1^*, p_2^*, p_3^*) . In this case, using equations (27) and (28) we find that

$$T_S(p_1^*, p_2^*, p_3^*) - T_H(p_1^*, p_2^*, p_3^*) = \frac{1}{1 - \alpha} \frac{t_a}{2}. \quad (36)$$

On the other hand, if (p_1^*, p_2^*, p_3^*) is an equilibrium strategy, then $T_{SA}^*(p_1^*, p_2^*, p_3^*) = T_{SA}^*(1, p_2^*, p_3^*) = T_{SA}^*(0, p_2^*, p_3^*)$ which yields that $T_{SA}^*(1, p_2^*, p_3^*) = T_S^*(0, p_2^*, p_3^*) = T_S^*(1, p_2^*, p_3^*) = T_S^*(p_1^*, p_2^*, p_3^*)$. Substituting into equation (24) we obtain that

$$T_S^*(p_1^*, p_2^*, p_3^*) - T_H^*(p_1^*, p_2^*, p_3^*) = \frac{1}{\nu_f f}. \quad (37)$$

Hence, if a strategy of the (S 6) class is an equilibrium strategy, then (36) and (37) must hold. This leads to

$$1 - \alpha = \nu_f f \frac{t_a}{2}. \quad (38)$$

i.e. that the probability of a challenger losing a fight is equal to the ratio of the expected duration of the fight and the mean time searching for food. These are all biologically-determined parameters, and we assume that the chance of their precise coincidence in this way is negligible (i.e. the case is non-generic). Thus, for example, such a case would correspond to a region of zero area in Figure 4, equivalent to the boundary lines.

698 Numerical investigation also indicates that mixed strategies are always invaded and
 699 so there are no mixed ESSs in the classes (S 4), (S 5) and (S 7).

700 Concerning strategies (S 2) and (S 3), numerical examples on a wide range of parame-
 701 ter values also imply that for every value of p_2 , $0 < p_2 < 1$, strategies $(0 < p_1 < 1, p_2, 0)$
 702 can always be invaded either by strategy $(0, p_2, 0)$ or by strategy $(1, p_2, 0)$. On the
 703 other hand, numerical examples indicate that for given p_1^* , $0 < p_1^* \leq 1$, there is a
 704 strategy p_2^* , $0 < p_2^* < 1$, such that for specific values of parameters all the invading
 705 strategies $(p_1^*, 0 \leq q_2 \leq 1, 0)$ do equally well in a population playing $(p_1^*, p_2^*, 0)$, i.e.
 706 $T_{\text{HA}}(p_1^*, p_2^*, 0) = T_{\text{HA}}^*(p_1^*, 0 \leq q_2 \leq 1, 0)$, while any other strategy does worse. Any other
 707 population playing a different strategy $(p_1^*, 0 < p_2 < 1, 0)$, $p_2 \neq p_2^*$, can be invaded
 708 either by the strategy $(p_1^*, 0, 0)$ or the strategy $(p_1^*, 1, 0)$ (see Figure 6b for an exam-
 709 ple). However, in a population which plays a strategy $(p_1, 0 \leq q_2 \leq 1, 0)$, the required
 710 time for an attacked handler playing the population strategy, $T_{\text{HA}}(p_1, 0 \leq q_2 \leq 1, 0)$,
 711 is less than that required by an attacked handler playing $(p_1^*, p_2^*, 0)$. In other words, if
 712 an infinitesimal portion of the population deviates from the equilibrium strategy, the
 713 evolution will drive away the population from that equilibrium. Thus, according to the
 714 second condition of Maynard Smith and Price (Maynard Smith and Price, 1973) for
 715 a strategy to be an ESS, the strategies $(p_1^*, p_2^*, 0)$ cannot be ESS. Hence, none of the
 716 strategies (S 2) and (S 3) can be evolutionarily stable.

References

- Axelrod R, Hamilton WD. 1981. The evolution of cooperation. *Science*. 211:1390–1396.
- Barnard CJ, Sibly RM. 1981. Producers and scroungers: a general model and its application to captive flocks of house sparrows. *Anim Behav*. 29:543–550.
- Beddington JR. 1975. Mutual interference between parasites or predators and its effect on searching efficiency. *J Anim Ecol*. 44:331–340.
- Brockman HJ, Barnard CJ. 1979. Kleptoparasitism in birds. *Anim Behav*. 27:487–514.
- Broom M, Luther RM, Ruxton GD. 2004. Resistance is useless? Extensions to the game theory of kleptoparasitism. *Bull Math Biol*. 66:1645–1658.
- Broom M, Luther RM, Ruxton GD, Rychtar J. 2008. A game-theoretic model of kleptoparasitic behavior in polymorphic populations. *J Theor Biol*. 255:81–91.
- Broom M, Ruxton GD. 1998. Evolutionarily stable stealing: game theory applied to kleptoparasitism. *Behav Ecol*. 9:397–403.
- Broom M, Ruxton GD. 2003. Evolutionarily stable kleptoparasitism: consequences of different prey types. *Behav Ecol*. 14:23–33.
- Broom M, Rychtar J. 2007. The evolution of a kleptoparasitic system under adaptive dynamics. *J Math Biol*. 54:151–177.
- Broom M, Rychtar J. 2009. A game theoretical model of kleptoparasitism with incomplete information. *J Math Biol*. 59:631–649.
- Broom M, Rychtar J. 2011. Kleptoparasitic melees - modelling food stealing featuring contests with multiple individuals. *Bull Math Biol*. 73:683–699.
- Caraco T, Giraldeau L-A. 1991. Social foraging: producing and scrounging in a stochastic environment. *J Theor Biol*. 153:559–583.
- Carbone C, Frame L, Frame G, Malcolm J, Fanshawe J, FitzGibbon C, Schaller G, Gordon IJ, Rowcliffe JM, Du Toit JT. 2005. Feeding success of African wild dogs

- (*Lycaon pictus*) in the Serengeti: the effects of group size and kleptoparasitism. *J Zool.* 266:153–161.
- Chamorro-Florescano IA, Favila ME, Macias-Ordonez R. 2010. Ownership, size and reproductive status affect the outcome of food ball contests in a dung roller beetle: when do enemies share? *Evol Ecol.* 25:277–289.
- Chellapa S, Hungtingford FA. 1989. Depletion of energy reserves during reproductive aggression in male three-spined stickleback, *Gasterosteus aculeatus* L. *J Fish Biol.* 35:315–316.
- Cooper SM. 1991. Optimal hunting groups size: the need for lions to defend their kills against loss to spotted hyaenas. *Afr J Ecol.* 29:130–136.
- Cooper WE, Perez-Mellado V. 2003. Kleptoparasitism in the Balearic lizard, *Podarcis lilfordi*. *Amphibia-Reptilia.* 24:219–224.
- Coyl FA, O'Shields TC, Perlmutter DG. 1991. Observations on the behaviour of the kleptoparasitic spider, *Mysmenopsis furtiva* (Araneae, Mysmenidae). *J Arachnol.* 19:62–66.
- Crowe M, Fitzgerald D, Remington D, Rychtar J. 2009. On deterministic and stochastic models of kleptoparasitism. *J Interdisciplinary Math.* 12:161–180.
- Dubois F, Giraldeau L-A. 2003. The forager's dilemma: food-sharing and food-defense as risk-sensitive foraging options. *Am Nat.* 162:768–779.
- Dubois F, Giraldeau L-A. 2005. Fighting for resources: the economics of defense and appropriation. *Ecology.* 86:3–11.
- Dubois F, Giraldeau L-A. 2007. Food sharing among retaliators: sequential arrivals and information asymmetries. *Behav Ecol Sociobiol.* 62:263–271.
- Dugatkin, LA. 1997. Winner and loser effects and the structure of dominance hierarchies. *Behav Ecol.* 8:583–587.
- Dubois F, Giraldeau L-A, Grant JWA. 2003. Resource defense in a group foraging context. *Behav Ecol.* 14:2–9.

- 769 Enquist M, Leimar O. 1987. Evolution of fighting behaviour: the effect of variation in
770 resource value. *J Theor Biol.* 127:187–205.
- 771 Erlandsson A. 1988. Food sharing vs. monopolising prey: a form of kleptoparasitism in
772 *Velia caprai* (Heteroptera). *Oikos.* 53:23–26.
- 773 Feistner ATC, McGrew WC. 1989. Food-sharing in primates: a critical review. In: Seth
774 PK, Seth S, editors. *Perspectives in primate biology*. Vol. 3. New Delhi: Today and
775 Tomorrow's Printers and Publishers. p. 21–36.
- 776 Hack MA. 1997. The energetic costs of fighting in the house cricket, *Acheta domesticus*
777 L. *Behav Ecol.* 8:28–36.
- 778 Hamilton WE, Dill LM. 2003. The use of territorial gardening versus kleptoparasitism,
779 Behavioral by a tropical reef fish (*Kyphosus cornelii*) is influenced by territory de-
780 pendability. *Behav Ecol.* 14:561–568.
- 781 Iyengar EV. 2002. Sneaky snails and wasted worms: kleptoparasitism by *Trichotropis*
782 cancellata (Mollusca, Gastropoda) on *Serpula columbiana* (Annelida, Polychaeta).
783 *Mar Ecol Prog Ser.* 244:153–162.
- 784 Iyengar EV. 2008. Kleptoparasitic interactions throughout the animal kingdom and a
785 re-evaluation, based on participant mobility, of the conditions promoting the evolution
786 of kleptoparasitism. *Biol J Linn Soc.* 93:745–762.
- 787 Janson CH. 1985. Aggressive competition and animal food consumption in wild brown
788 capuchin monkeys (*Cebus apella*). *Behav Ecol and Sociobiol.* 18:125–138.
- 789 Kacelnik A, Bateson M. 1996. Risky theories—The effects of variance on foraging deci-
790 sions. *Amer Zool.* 36:402–434.
- 791 Luther RM, Broom M. 2004. Rapid convergence to an equilibrium state in kleptopara-
792 sitic populations. *J Math Biol.* 48:325–339.
- 793 Luther RM, Broom M, Ruxton GD. 2007. Is food worth fighting for? ESS's in mixed
794 populations of kleptoparasites and foragers. *Bull Math Biol.* 69:1121–1146.

- 795 Maynard Smith J. 1982. Evolution and the theory of games. Cambridge, UK: Cambridge
796 University Press.
- 797 Maynard Smith J, Price GR. 1973. The logic of animal conflict. *Nature*. 246:15–18.
- 798 Morrell LJ, Kokko H. 2003. Adaptive strategies of territory formation. *Behav Ecol So-*
799 *ciobiol* 54:385–395.
- 800 Neat FC, Taylor AC, Huntingford FA. 1998. Proximate costs of fighting in male cichlid
801 fish: the role of injuries and energy metabolism. *Anim Behav*. 55:875–882.
- 802 Rovero F, Hughes RN, Whiteley NM, Chelazzi G. 2000. Estimating the energetic cost
803 of fighting in shore crabs by noninvasive monitoring of heartbeat rate. *Anim Behav*.
804 59:705–713.
- 805 Ruxton GD, Gurney WSC, De Roos AM. 1992. Interference and generation cycles.
806 *Theor Pop Biol*. 42:235–253.
- 807 Ruxton GD, Moody AL. 1997. The ideal free distribution with kleptoparasitism. *J Theor*
808 *Biol*. 186:449–458.
- 809 Selten R. 1975. Reexamination of the perfectness concept for equilibrium points in ex-
810 tensive games. *Int J Game Theory*. 4:25–55.
- 811 Sirot E. 2000. An evolutionarily stable strategy for aggressiveness in feeding groups.
812 *Behav Ecol*. 11:351–356.
- 813 Smallegange IM, Van der Meer J. 2009. The distribution of unequal predators across
814 food patches is not necessarily (semi)truncated. *Behav Ecol*. 20:525–534.
- 815 Smith IP, Taylor AC. 1993. The energetic cost of agnostic behaviour in the velvet swim-
816 ming crab, *Necora* (= *Liocarcinus*) *puber* (L.). *Anim Behav*. 45:375–391.
- 817 Stamps J. 1994. Territorial behavior: testing the assumptions. *Adv Study Behav*. 23:173–
818 232.
- 819 Stamps JA, Krishnan VV. 1994. Territory acquisition in lizards: II. Establishing social
820 and spatial relationships. *Anim Behav*. 47:1387–1400.

- 821 Stevens JR, Gilby IC. 2004. A conceptual framework for nonkin food sharing: timing
822 and currency of benefits. *Anim Behav.* 67:603–614.
- 823 Stevens JR, Stephens DW. 2002. Food sharing: a model of manipulation by harassment.
824 *Behav Ecol.* 13:393–400.
- 825 Vahl WK. 2006. Interference competition among foraging waders [Ph.D. Thesis]. Nether-
826 lands: University of Groningen.
- 827 Van der Meer J, Ens BJ. 1997. Models of interference and their consequences for the
828 spatial distribution of ideal and free predators. *J Anim Ecol.* 66:846–858.
- 829 Van der Meer J, Smallegange IM. 2009. A stochastic version of the Beddington-
830 DeAngelis functional response: modelling interference for a finite number of predators.
831 *J Anim Ecol.* 78:134–142.
- 832 Vickery WL, Giraldeau L-A, Templeton JJ, Kramer DL, Chapman CA. 1991. Producers,
833 scroungers and group foraging. *Am Nat.* 137:847–863.
- 834 Wenzel JW, Pickering J. 1991. Cooperative foraging, productivity, and the central limit
835 theorem. *Proc Natl Acad Sci USA.* 88:36–38.
- 836 White FJ. 1994. Food sharing in wild pygmy chimpanzees, *Pan paniscus*. In: Roeder
837 JJ, Thierry B, Anderson JR, Herrenschmidt N, editors. *Current Primatology: Social*
838 *Development, Learning and Behavior*. Vol. 2. Strasbourg: Universite Louis Pasteur.
839 p. 1–10.
- 840 Wilkinson GS. 1990. Food sharing in vampire bats. *Sci Am.* 262:76–82.
- 841 Yates GE, Broom M. 2007. Stochastic models of kleptoparasitism. *J Theor Biol.* 248:480–
842 489.

Table 1
The model notations

Population's densities	Meaning
P	Density of the population
S, H, C, A, R	Density of searchers, handlers, sharers, attackers and defenders
Model Parameters	Meaning
$\nu_f f$	Rate at which foragers find undiscovered food
$\nu_h H$	Rate at which foragers encounter handlers
t_h	Expected time for a handler to consume a food item if it is not attacked
t_c	Expected time for a sharer to consume the half of a food item
$t_a/2$	Expected duration of a fight
α	The probability that the attacker wins the fight
Strategies	Meaning
p_1	The probability that a searcher attacks a handler when they meet
p_2	The probability that an attacked handler shares its food item
p_3	The probability that an attacked handler defends its food item

Table 2

Notations of the required times to the consumption of a food item from the different foraging states

Notation	Meaning
T_{SA}	The average time needed for a searcher animal who has just encountered a handler to acquire and consume a food item
T_{HA}	The average time needed for a handler animal who has just encountered a searcher to consume a food item
T_S	The average time needed for an animal who has just become a searcher to acquire and consume a food item
T_H	The average time needed for an animal who has just become a handler to consume a food item
T_A	The average time needed for an attacker who has just engaged in a fight to acquire and consume a food item
T_R	The average time needed for a defender who has just engaged in a fight to consume a food item
T_C	The average time needed for a sharer to consume a food item

844

845

846

847

Table 3**Conditions under which a mutant playing strategy (q_1, q_2, q_3) cannot invade a population playing strategy (p_1, p_2, p_3)**

		Mutant's strategy, (q_1, q_2, q_3)					
		(0,0,0)	(0,1,0)	(0,0,1)	(1,0,0)	(1,1,0)	(1,0,1)
Population's strategy, (p_1, p_2, p_3)	(0,0,0)	/	$2t_c - t_h > \frac{1}{\nu_f f}$ (C.1)	$\nu_f f > \frac{2(1-\alpha)}{t_a}$ (C.2)	The mutant always invades (C.3)	—	—
	(0,1,0)	$2t_c - t_h < \frac{1}{\nu_f f}$ (C.4)	/	$2t_c - t_h < t_a - \frac{1-2\alpha}{\nu_f f}$ (C.5)	—	$2t_c - t_h > \frac{1}{\nu_f f}$ (C.6)	—
	(0,0,1)	$\nu_f f < \frac{2(1-\alpha)}{t_a}$ (C.7)	$2t_c - t_h > t_a - \frac{1-2\alpha}{\nu_f f}$ (C.8)	/	—	—	$\nu_f f > \frac{2\alpha}{t_a}$ (C.9)
	(1,0,0)	The mutant never invades (C.10)	—	—	/	$2t_c - t_h > \frac{1}{\nu_f f}$ (C.11)	$\nu_f f > \frac{2(1-\alpha)}{t_a} - \frac{t_h \nu_f f \nu_h P}{t_h \nu_f f + 1}$ (C.12)
	(1,1,0)	—	$2t_c - t_h < \frac{1}{\nu_f f - \nu_h H_a}$ * (C.13)	—	$2t_c - t_h < \frac{1}{\nu_f f - \nu_h H_a}$ * (C.14)	/	$(2t_c - t_h)(\nu_f f - \alpha \nu_h H_a) \nu_f f < (t_a \nu_f f + \alpha) \nu_f f + (1-\alpha)((t_h \nu_h P - 1) \nu_f f - (\nu_f f + \nu_h P) t_h \nu_h H_a)$ * (C.15)
	(1,0,1)	—	—	$\nu_f f < \frac{2\alpha}{t_a}$ (C.16)	$\nu_f f < \frac{2(1-\alpha)}{t_a} + (1-2\alpha) \nu_h H_b$ ** (C.17)	$(2t_c - t_h)(\nu_f f + \alpha \nu_h H_b) > t_a(\nu_f f + 2\alpha \nu_h H_b) + \alpha t_h(\nu_h P - \nu_h H_b) + 2\alpha - 1$ ** (C.18)	/

* H_a is given by the solution of the equation $(2t_c - t_h) \nu_h H_a^2 + (t_h \nu_f f + t_h \nu_h P + 1) H_a - t_h \nu_f f P = 0$.** H_b is given by the solution of the equation $t_a \nu_h H_b^2 + (t_h \nu_f f + 1) H_b - t_h \nu_f f P = 0$.

Table 4

Conditions under which a mutant playing strategy (q_1, q_2, q_3) cannot invade a population playing strategy (p_1, p_2, p_3) in the special case where $2t_c = t_h$

		Mutant's strategy, (q_1, q_2, q_3)					
		(0,0,0)	(0,1,0)	(0,0,1)	(1,0,0)	(1,1,0)	(1,0,1)
Population's strategy, (p_1, p_2, p_3)	(0,0,0)	/	The mutant always invades	$\nu_f f > \frac{2(1-\alpha)}{t_a}$	The mutant always invades	—	—
	(0,1,0)	The mutant never invades	/	$\nu_f f > \frac{1-2\alpha}{t_a}$	—	The mutant always invades	—
	(0,0,1)	$\nu_f f < \frac{2(1-\alpha)}{t_a}$	$\nu_f f < \frac{1-2\alpha}{t_a}$	/	—	—	$\nu_f f > \frac{2\alpha}{t_a}$
	(1,0,0)	The mutant never invades	—	—	/	The mutant always invades	$\nu_f f > \frac{2(1-\alpha)}{t_a} - \frac{t_h \nu_f f \nu_h P}{t_h \nu_f f + 1}$
	(1,1,0)	—	The mutant never invades	—	The mutant never invades	/	$(t_a \nu_f f + \alpha)(t_h \nu_f f + t_h \nu_h P + 1) - (1-\alpha)(t_h \nu_f f + 1) > 0$
	(1,0,1)	—	—	$\nu_f f < \frac{2\alpha}{t_a}$	$\nu_f f < \frac{2(1-\alpha)}{t_a} + (1-2\alpha)\nu_h H_b$ *	$t_a(\nu_f f + 2\alpha\nu_h H_b) + \alpha t_h(\nu_h P - \nu_h H_b) + 2\alpha - 1 < 0$ *	/

* H_b is given by the solution of the equation $t_a \nu_h H_b^2 + (t_h \nu_f f + 1)H_b - t_h \nu_f f P = 0$.

Table 5
Possible ESSs

		Strategy at the handling state, p_2, p_3						
		$p_2 = 1$	$p_2 = 0$		$p_2 = 0$	$0 < p_2 < 1$	$0 < p_2 < 1$	$0 < p_2 < 1$
		$p_3 = 0$	$p_3 = 1$	$p_2 = p_3 = 0$	$0 < p_3 < 1$	$p_3 = 0$	$0 < p_3 < 1$	$0 < p_3 < 1$
							$p_2 + p_3 = 1$	$p_2 + p_3 < 1$
Strategy at the searching state, p_1	$p_1 = 0$	**	*	*	*	***	***	***
	$0 < p_1 < 1$	S 1	*	*	*	S 2	S 4	S 6
	$p_1 = 1$	**	*	*	*	S 3	S 5	S 7

Figure legends

Figure 1: Schematic representation of all the possible events that might happen until the consumption of a food item by a mutant searcher playing strategy (q_1, q_2, q_3) who encounters a handler of a population playing strategy (p_1, p_2, p_3) . The transition probabilities and the expected times (in bold) to move from one state to another are shown.


Figure 2: Schematic representation of all the possible events that might happen until the consumption of a food item by a mutant handler playing strategy (q_1, q_2, q_3) who encounters a searcher of a population playing strategy (p_1, p_2, p_3) . The transition probabilities and the expected times (in bold) to move from one state to another are shown.

Figure 3: Graphs showing examples of the region where each of the four possible ESSs (Retaliator (R), Marauder (M), Attacking Sharer (AS) and Hawk (H)) is an ESS as the duration of the content, $t_a/2$, and the handling time of a sharer, t_c , vary. In each region, a single letter 'X' indicates that the strategy X is the unique ESS, 'X,Y' indicates that the strategies X and Y are simultaneous ESSs and 'X, Y, Z' that the three strategies X, Y and Z are simultaneous ESSs. (a) $t_h = 3, \nu_f f = 0.5, \nu_h = 1.5, \alpha = 0.7, P = 1$; (b) $t_h = 3, \nu_f f = 1, \nu_h = 2, \alpha = 0.2, P = 1$.


Figure 4: Graphs showing examples of the region where each of the four possible ESSs (Retaliator (R), Marauder (M), Attacking Sharer (AS) and Hawk (H)) is an ESS as the density of the population, P , and the rate at which foragers find undiscovered food, $\nu_f f$, vary. In each region, a single letter 'X' indicates that the strategy X is the unique ESS, 'X,Y' indicates that the strategies X and Y are simultaneous ESSs and 'X, Y, Z' that the three strategies X, Y and Z are simultaneous ESSs. (a) $t_a/2 = 0.5, t_h = 3, t_c = 4, \nu_h = 1.5, \alpha = 0.7$; (b) $t_a/2 = 0.5, t_h = 3, t_c = 2, \nu_h = 2, \alpha = 0.2$.

Figure 5: A graph showing an example of the region where each of the three possible ESSs (Retaliator (R), Attacking Sharer (AS) and Hawk (H)) can occur in the special case where $2t_c = t_h$, as the probability α of the challenger winning and the duration of the content, $t_a/2$, vary. In each region, a single letter 'X' indicates that the strategy X is the unique ESS, 'X,Y' indicates that the strategies X and Y are simultaneous ESSs and 'X, Y, Z' that the three strategies X, Y and Z are simultaneous ESSs. $2t_c = t_h = 3, \nu_f f = 0.5, \nu_h = 1.5, P = 1$.


Figure 6: (a) The expected time until the consumption of a food item of mutant searcher animals playing strategies $(0,1,0)$, $(1,1,0)$ and $(0 < p_1 < 1, 1, 0)$ in a population playing strategy $(0 < p_1 < 1, 1, 0)$ for the example considered in Figure 3b for $t_c = 2$. Numerical examples indicate that in every population which adopts a strategy $(0 < p_1 < 1, 1, 0)$, either a mutant animal that plays strategy $(0, 1, 0)$ or a mutant animal that plays strategy $(1, 1, 0)$ always does better than any other animal that uses the population strategy. Thus, such populations can be invaded by those mutant strategies and as a result, strategies $(0 < p_1 < 1, 1, 0)$ cannot be ESSs. (b) The expected time until the consumption of a food item of mutant handler animals playing strategies $(0.8,1,0)$, $(0.8,0,0)$ and $(0.8, 0 < p_2 < 1, 0)$ in a population playing strategy $(0.8, 0 < p_2 < 1, 0)$ for $t_h = 3, t_c = 2, \nu_f f = 1, \nu_h = 1.5, P = 1$. An equilibrium strategy $(0 < p_1 \leq 1, 0 < p_2 < 1, 0)$ cannot be evolutionarily stable.

**Figure 1**


Schematic representation of all the possible events that might happen until the consumption of a food item by a mutant searcher playing strategy (q_1, q_2, q_3) who encounters a handler of a population playing strategy (p_1, p_2, p_3) . The transition probabilities and the expected times (in bold) to move from one state to another are shown.

**Figure 2**


Schematic representation of all the possible events that might happen until the consumption of a food item by a mutant handler playing strategy (q_1, q_2, q_3) who encounters a searcher of a population playing strategy (p_1, p_2, p_3) . The transition probabilities and the expected times (in bold) to move from one state to another are shown.

**Figure 3**


Graphs showing examples of the region where each of the four possible ESSs (Retaliator (R), Marauder (M), Attacking Sharer (AS) and Hawk (H)) is an ESS as the duration of the content, $t_a/2$, and the handling time of a sharer, t_c , vary. In each region, a single letter 'X' indicates that the strategy X is the unique ESS, 'X,Y' indicates that the strategies X and Y are simultaneous ESSs and 'X, Y, Z' that the three strategies X, Y and Z are simultaneous ESSs. (a) $t_h = 3$, $\nu_f f = 0.5$, $\nu_h = 1.5$, $\alpha = 0.7$, $P = 1$; (b) $t_h = 3$, $\nu_f f = 1$, $\nu_h = 2$, $\alpha = 0.2$, $P = 1$.

**Figure 4**

Graphs showing examples of the region where each of the four possible ESSs (Retaliator (R), Marauder (M), Attacking Sharer (AS) and Hawk (H)) is an ESS as the density of the population, P , and the rate at which foragers find undiscovered food, $\nu_f f$, vary. In each region, a single letter 'X' indicates that the strategy X is the unique ESS, 'X,Y' indicates that the strategies X and Y are simultaneous ESSs and 'X, Y, Z' that the three strategies X, Y and Z are simultaneous ESSs. (a) $t_a/2 = 0.5$, $t_h = 3$, $t_c = 4$, $\nu_h = 1.5$, $\alpha = 0.7$; (b) $t_a/2 = 0.5$, $t_h = 3$, $t_c = 2$, $\nu_h = 2$, $\alpha = 0.2$.

**Figure 5**

A graph showing an example of the region where each of the three possible ESSs (Retaliator (R), Attacking Sharer (AS) and Hawk (H)) can occur in the special case where $2t_c = t_h$, as the probability α of the challenger winning and the duration of the content, $t_a/2$, vary. In each region, a single letter 'X' indicates that the strategy X is the unique ESS, 'X,Y' indicates that the strategies X and Y are simultaneous ESSs and 'X, Y, Z' that the three strategies X, Y and Z are simultaneous ESSs. $2t_c = t_h = 3, \nu_{ff} = 0.5, \nu_h = 1.5, P = 1$.

**Figure 6**

(a) The expected time until the consumption of a food item of mutant searcher animals playing strategies $(0,1,0)$, $(1,1,0)$ and $(0 < p_1 < 1, 1, 0)$ in a population playing strategy $(0 < p_1 < 1, 1, 0)$ for the example considered in Figure 3b for $t_c = 2$. Numerical examples indicate that in every population which adopts a strategy $(0 < p_1 < 1, 1, 0)$, either a mutant animal that plays strategy $(0, 1, 0)$ or a mutant animal that plays strategy $(1, 1, 0)$ always does better than any other animal that uses the population strategy. Thus, such populations can be invaded by those mutant strategies and as a result, strategies $(0 < p_1 < 1, 1, 0)$ cannot be ESSs. (b) The expected time until the consumption of a food item of mutant handler animals playing strategies $(0.8,1,0)$, $(0.8,0,0)$ and $(0.8, 0 < p_2 < 1, 0)$ in a population playing strategy $(0.8, 0 < p_2 < 1, 0)$ for $t_h = 3, t_c = 2, \nu_{ff} = 1, \nu_h = 1.5, P = 1$. An equilibrium strategy $(0 < p_1 \leq 1, 0 < p_2 < 1, 0)$ cannot be evolutionarily stable.