



# City Research Online

## City St George's, University of London

**Citation:** Hashemi, L., Ghasemi, M., Bartley, A., Fenaughty, J., Pirouzi, M. & Grant, C. (2024). Migration and infant immunization timeliness in New Zealand: Evidence from the Growing Up in New Zealand study. *Vaccine*, 42(9), pp. 2229-2238. doi: 10.1016/j.vaccine.2024.02.053

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

**Permanent repository link:** <https://city-ref-test.eprints-hosting.org/id/eprint/32505/>

**Link to published version:** <https://doi.org/10.1016/j.vaccine.2024.02.053>

**Copyright and Reuse:** Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

## **Migration and Infant Immunization Timeliness in New Zealand: Evidence from the Growing Up in New Zealand Study**

Immunization is widely recognized as one of the most cost-effective methods for preventing infectious diseases [1], [2], especially among young children. This age group is particularly vulnerable to severe diseases caused by vaccine-preventable infections and also plays a significant role in the spread of infections through households and communities.

Vaccine coverage is used as a measure to estimate the potential population protection offered by an immunization program. However, it is important to note that high coverage for individual vaccines may still be insufficient to fully prevent diseases [3]-[5]. In fact, when it comes to preventing diseases in young children, the *timeliness* of vaccine receipt is a key factor [6]-[8].

Outbreaks of vaccine-preventable diseases can occur if a sufficient proportion of the population does not receive *timely* vaccination, even if the overall population coverage rates are high [6], [8].

Timeliness of vaccination is particularly important for the primary infant vaccine series and serves as a key indicator of health care system performance and the accessibility of health services across different sectors of the community.

Studies conducted in developed countries have highlighted migration as an important determinant of child health outcomes, including childhood vaccination status [1], [9], [10]. Children from immigrant families are often at an elevated risk of not being adequately immunized [1], [11]-[13], leaving them more susceptible to vaccine-preventable diseases [14]-[16].

Barriers to immunization for migrant children encompass various factors, including differences in healthcare systems between the source and destination countries [17], language proficiency, cultural differences, and financial constraints [18], [19]. Recent immigrants encounter additional challenges as catch-up immunization may be overlooked upon arrival due to competing settlement

priorities. Additionally, during the early stage of settlement, there may be missed opportunities to complete immunization schedules on time [20]. These challenges place children of recent immigrants at an elevated risk of not being fully immunized or immunized on time [9], [20].

Lower rates of timely vaccination among children of foreign-born parents compared with children of native-born parents has been documented previously [9], [23]. However, the evidence remains inconclusive, as some studies have reported no significant association between parental migration and children's timely vaccination [21], [22], while others have found that children of foreign-born parents were more likely to be immunized than children of native-born parents [23]-[25].

These discrepancies in studies of immigrant children may arise from differences in how children's migration status is defined. While some studies classify foreign-born children as migrant children [26], [27], others include children born in the host country to migrant parents [26]-[29]. The variation in definitions can lead to differences in the composition of study populations and may influence the interpretation of findings.

An example of such research conducted in New Zealand (NZ) investigated differences in age-appropriate immunization coverage rates among children with and without migrant backgrounds. The study found that foreign-born children had lower recorded age-appropriate vaccination rates for all vaccines included in the NZ National Immunization Schedule compared with NZ-born children from migrant or non-migrant families [11]. Utilizing data from the NZ National Immunization Register (NIR), this study provided valuable insights into immunization inequities particularly for foreign-born children. However, it is important to note some limitations of the study. For example, less than half (45%) of foreign-born children had an immunization record in the NIR dataset compared to ~ 95% of NZ-born children with or without a migration background. This significant disparity highlights potential concerns about the completeness of vaccination data

for foreign-born children in the NIR. Moreover, difficulties in translating immunization schedules from the origin countries of migrants who vaccinated their children in their home country may contribute to inaccuracies in recording, potentially leading to an underestimation of migrant children's vaccination coverage. Such speculation is supported by another published research using the same sample/datasets [26], which found significantly lower hospitalization rates among foreign-born children compared with NZ-born non-migrant children for a wide range of vaccine-preventable diseases. Further research is warranted to explore and confirm these contradictory findings and to address the immunization inequalities among migrant children in New Zealand.

New Zealand has an estimated 25.2% of the resident population born overseas. Additionally, there has been a significant increase in the immigrant population, including children, over the last decade [30]. While vaccine coverage and timeliness disparities by ethnicity in NZ have been described [31], little is known about the role of parental migration status on child health outcomes, such as childhood immunization in NZ.

Understanding the health-related behaviors of this group is of paramount importance to both medical and public health professionals. This information is of particular importance as rates of immunizations for many infectious diseases, across various age groups, are declining in NZ. Additionally, some preventable diseases, such as measles and pertussis, continue to cause serious outbreaks, morbidity, and mortality [31]. Furthermore, immunization timeliness remains suboptimal, with delays in the receipt of scheduled immunizations continuing to limit the potential benefits that immunization could offer to population health in NZ [7], [20], [32].

Identifying groups with lower immunization uptake or untimely vaccination, as well as the factors associated with their immunization outcomes, is imperative for the success of immunization programs and the efficient allocation of public health resources [33].

To date, most studies in this field have primarily categorized children into two groups: migrant and non-migrant [11]. However, data from some studies [21], [34] suggest that a more refined stratification of the immigrant population into settled and recent immigrants could provide a more nuanced understanding of the impact of migration-related factors on the immunization status of immigrant children. By distinguishing between settled immigrants, who have resided in the host country for an extended period, and recent immigrants, who have arrived more recently, researchers can better assess the temporal aspects of migration and their influence on vaccination outcomes.

Furthermore, most of the research in this field has been predominantly focused on the role of mother's migration status on children's health outcomes. However, there is a paucity of research investigating how father's migration status might be influencing children's outcomes.

## **Current study**

Using data from a large, representative population-based cohort study in NZ, the present research aims to: a) explore disparities in vaccination timeliness among NZ-born children from immigrant and non-immigrant families; b) examine the role of residential duration of immigrant families on children's vaccination timeliness; c) investigate whether maternal versus paternal migration status has a different impact on child vaccination timeliness; and d) describe the differences in vaccination timeliness between children of one and two foreign-born parents versus children of two native-born parents.

## **Methods**

## ***Participants***

The data used in this study was taken from the Growing Up in New Zealand (GUiNZ) study, the largest contemporary longitudinal study of child development in NZ. The GUiNZ study has been designed and conducted by a team of researchers led by the University of Auckland. Detailed information regarding the study's design, conceptual framework, and recruitment procedures can be found elsewhere [35], [36]. A total of 6,822 pregnant women with an estimated delivery date between April 2009 and March 2010 were recruited from the Auckland, Counties Manukau, and Waikato District Health Board regions of NZ.

All participating women provided written informed consent. Partners were enrolled if the enrolled women provided consent for their participation. A total of 4,404 partners were enrolled in the study.

The child cohort (n=6853) comprised ~ 11% of births in NZ during the recruitment period. The characteristics of the cohort at birth generally aligned with the national birth cohort in NZ from 2007 to 2010 [36]. Ethical approval for the study was granted by the NZ Ministry of Health Northern Y Regional Ethics Committee (NTY/08106/055).

The current study utilizes data collected during the first data collection wave (antenatal) from the nine data collection waves conducted between 2009 and 2022. The antenatal data was linked with the National Immunization Register (NIR) dataset, a process carried out by the study team (the linkage between the child's unique identifier and NIR had been previously conducted by the GUiNZ team). Data collection waves used face-to-face and phone interviews to gather information. The antenatal interview was completed separately for mothers and partners. To avoid dependent observations, only one cohort child from each family was included in our analyses. As a result, the data for second-order children of 93 multiple births were deleted. Additionally, data

for a further 430 children were deleted due to not having any NIR data for linkage. The data for another 143 children were also deleted because their mothers or caregivers did not provide consent to collect data from external health agencies. This resulted in a final sample size of 6,156 children and their parents for this study.

## **Measures**

### **The main outcome of interest**

Each child's immunization record was obtained via each child's National Health Index (NHI) number linked with the NIR. The NHI number is a unique identifier assigned to every person having contact with health services in NZ [37]. Over the time the cohort was born, the infant immunization schedule included doses at 6-week, 3-month, and 5-month for two vaccines: a diphtheria/tetanus/acellular pertussis/*Haemophilus influenzae* type B/hepatitis B/poliovirus vaccine and a pneumococcal conjugate vaccine [37]. We refer to the three doses of these two vaccines as all six doses of vaccines in this paper.

Timely immunization was defined as vaccinations received within 30 days of their due date [38]. This definition has been previously used in the United States and Australia [39]. The estimate of immunization timeliness utilizing this method for the GUiNZ cohort (70%) closely mirrors the NZ's national immunization coverage at six months of age (71%), thereby confirming the validity of this measure [7].

### **Main exposure of interest**

#### **Immigration status**

All children included in the study cohort were born in NZ. Immigrant children were identified based on parental immigration status rather than the children's immigration status. Recent versus

settled immigrants were identified based on maternal residential duration. Recent immigrants were identified as mothers residing in NZ for less than five years at the time of antenatal data collection wave, while settled immigrants were identified as mothers who had been residing in the country for more than five years at the time of the antenatal data collection wave. This five-year residency benchmark was chosen following the guidelines of the NZ Ministry of Business, Innovation & Employment to differentiate between recent and settled immigrants [40]. Repeating the analyses with different criteria to define recent and settled immigrants (less than three years versus more than three years) yielded similar results.

#### **Sociodemographic characteristics**

Potential covariates for inclusion in multivariable analyses were selected based on prior research on factors associated with immunization timeliness in New Zealand [7], [8] and included: maternal age at pregnancy, maternal education (categorized into four groups: no secondary school, secondary school, diploma/trade, tertiary), household crowding index group [41] (classified into three categories: low [less than one person per bedroom], medium [one to less than two people per bedroom], and high [two or more people per bedroom] crowding score), language spoken at home (English, non-English), household annual income groups (categorized into three categories: < \$50K, \$50-\$70K, > \$70K), receipt of income tested benefit (yes/no), and area-level deprivation index (NZDep2006).

The NZDep2006 index of deprivation was used to measure neighborhood area deprivation [42]. This index is derived from the household geographical location and is measured at the meshblock level, which is the smallest census tract unit. It combines census data from 2006 relating to various factors including income, home ownership, employment, qualifications, family structure, housing,

and access to transport and communications [42]. The index scores are organized into deciles, with decile one representing the least deprived 10% of areas in NZ, and decile 10 indicating the most deprived 10% of areas. For the current analysis, deprivation scores were categorized into three groups: low deprivation (deciles 1 to 3), medium deprivation (deciles 4 to 7), and high deprivation areas (deciles 8 to 10).

### **Analytical procedures**

All analyses were conducted using Stata 14.2 [43]. Sociodemographic characteristics of the whole sample and stratified by maternal immigration status (immigrant versus non-immigrant), and maternal residential duration in NZ (recent versus settled immigrants) are presented in **Table 1**.

The prevalence rate of timely immunization of all studied vaccines (all doses of six-week, three-month, five-month vaccines, and all six doses of studied vaccines) for the overall sample and by maternal migration status and maternal residential duration in NZ are presented in **Table 3**. The same analyses were repeated for paternal immigration status (foreign-born father vs NZ-born father) and one and two foreign-born parents versus two NZ-born parents (**Table 3**).

Distribution of infant timely vaccination across sociodemographic factors are presented in **Supplementary Table 1**.

Chi-square tests were used to assess the associations between sociodemographic variables and maternal migration status and residential duration in NZ (**Table 1**). Similarly, chi-square tests were employed to examine the associations between sociodemographic variables and infant immunization timeliness (**Table 2**).

The bivariate associations between maternal immigration status, maternal residential duration, and infant vaccination timeliness at age six-week, three-month, and five-month were also examined

using univariate logistic regression models with results presented as unadjusted odds ratios (ORs) with 95% confidence intervals (CIs) (**Table 4**).

Then, to determine if the noted differences in the prevalence rates of timely infant vaccination between children of immigrant and non-immigrant families found in the univariate analyses remained significant after controlling for sociodemographic characteristics, the following steps were taken:

First, the bivariate association between each sociodemographic characteristic and infant vaccination status (timeliness) at age 6-weeks, three-months, and five-months (timely or not) was explored using univariate logistic regression models with results presented as unadjusted odds ratios (ORs) with 95% CIs (**Table 2**).

Second, multivariable logistic regression analyses were conducted, with maternal immigration status and sociodemographic characteristics included as exposures and infant vaccination status as the outcome variable. The results were presented as adjusted odds ratios (AOR) with 95% CIs. The same analyses were repeated with maternal residential duration as the main exposure variable (**Table 4**). Due to the high correlation between the “language spoken at home” variable and the main exposure variable (immigration status), “language spoken at home” was excluded from the multivariable regression analyses.

## **Results**

### **Sociodemographic characteristics of the study sample by maternal immigration status and maternal residential duration**

The study sample consisted of 6,156 mothers, of whom 36.4% (2,241) were born outside NZ. Among foreign-born mothers, 43.2% had attained tertiary education, compared to 35.6% of NZ-born mothers. There was a larger proportion of mothers aged less than 20 years in the NZ-born group compared to the foreign-born group (6.6% versus 1.8% respectively). Additionally, a larger proportion of foreign-born mothers (42.5%) resided in highly deprived areas compared to NZ-born mothers (36.7%). Over half of the foreign-born mothers (51.7%) lived in households with an income exceeding \$70k, in contrast to 64.6% of NZ-born mothers (**Table 1**).

Moreover, a higher proportion of foreign-born mothers lived in crowded households compared to NZ-born mothers (21.6% versus 10.1% respectively) and a larger proportion reported speaking a language other than English at home compared to NZ-born mothers (52% versus 2.4% respectively) (**Table 1**).

No significant differences were found between recent and settled immigrants regarding maternal education levels and household deprivation index. However, a higher proportion of recent immigrants were residing in crowded households compared to settled immigrants (24.6% versus 19.2% respectively). Conversely, a smaller proportion of recent immigrants lived in households with incomes exceeding \$70k, compared to settled immigrants (43.8% versus 57.5%, respectively), and a smaller proportion of recent immigrants reported receiving an income tested benefit compared to settled immigrants (7% versus 12.4%, respectively) (**Table 1**).

A smaller proportion of recent immigrants (37%) reported speaking a language other than English at home, compared to settled immigrants (55%) (**Table 1**).

**Insert Table 1 here**

**Association between sociodemographic factors and infant timely vaccination**

Maternal age at pregnancy was not found to be associated with timely infant vaccination at any time point. However, all indicators of higher socio-economic status among mothers, including attaining some level of education, having a household income exceeding \$70K, residing in less deprived areas, living in a household with low to medium crowding scores, or not receiving an income tested benefit, were significantly associated with increased odds of infant vaccinations being received on time. Additionally, speaking a language other than English was associated with an increased odd of timely infant vaccination (**Table 2**).

**Insert Table 2 here**

#### **Prevalence of timely completion of infant vaccinations by maternal immigration status**

A larger proportion of children born to foreign-born mothers had timely receipt of all six-week, three-month, and five-month vaccine series as well as all six doses of vaccines, with rates ranging from 95.9% to 85.8%, compared to children of NZ-born mothers, where rates ranged from 91.4% to 79.8% for the same vaccine series (**Table 3**).

A larger proportion of children born to recent immigrants achieved timely completion of all six-week, three-month, and five-month vaccine series, as well as all six doses of vaccines, compared to children of settled immigrants (**Table 3**).

For all child groups, the proportion of timely receipt of age-specific vaccines decreased with increasing infant age, from the six-week to the 5-month vaccinations. The largest decrease was observed for infants of foreign-born mothers who had resided in the country for more than five years (settled immigrants), with a difference of 12.2 percentage points between their timely completion rate for six-week and 5-month doses (95.5% and 83.3% respectively). However, their counterparts who resided in the country for less than five years (recent immigrants) experienced

the smallest decrease, with a difference of 7.5 percentage points from six weeks to five months (96.5% and 89.0% respectively) (**Table 3**).

**Insert Table 3 here**

#### **Maternal migration status and infant vaccination timeliness**

The results of bivariate logistic regressions (**Table 4**) were consistent with those presented in **Table 3**, indicating that children of foreign-born mothers had higher odds of being vaccinated on time compared to children of NZ-born mothers. Unadjusted odds ratios ranged from 2.22 to 1.46 for all doses of vaccines from six weeks to five months of age.

The results of multivariable logistic regressions indicated that after controlling for socio-demographic differences between immigrant and non-immigrant families, the odds of receiving all doses of vaccines on time increased and remained significant (adjusted odds ratios ranging from 2.55 to 1.54 for all doses of vaccines from age six weeks to 5 months). A steady decrease in the magnitude of both unadjusted and adjusted odds ratios with increasing infant age was observed, indicating that the gap in child vaccination timeliness narrowed between immigrant and non-immigrant groups as children aged. However, by the age of five months, differences were still statistically significant, with children of immigrants having higher odds of receiving the 5-month vaccine doses on time (**Table 4**).

**Insert table 4 here**

#### **Maternal residential duration and infant vaccination timeliness**

The children of recent immigrants had increased odds of being vaccinated on time compared with children of settled immigrants. The only exception was for the 6-week vaccine series where odds

ratios did not reach significance. After adjusting for socio-demographic characteristics, children of recent immigrants had increased odds of receiving the three-month and five-month vaccine doses on time compared to those of settled immigrants (AOR=1.52, 95%CI 1.05-2.22 and AOR=1.72, 95%CI 1.28-2.33 respectively) (**Table 4**).

### **The impact of paternal immigration status and having both foreign-born parents on infant vaccination timeliness**

Having a foreign-born father was associated with identical timeliness rates of all vaccination series compared to those with a foreign-born mother (83 versus 83%). Additionally, having a foreign-born father was positively and significantly associated with timely completion of all six doses of vaccines (83% versus 80% among children with a NZ-born father,  $p<0.001$ ). Moreover, children with both parents born overseas had the highest rates of timely vaccination for all six doses of vaccines (85.2%), followed by those with one foreign-born parent (82.1%). Conversely, children with both parents born in NZ had the lowest rates of timely vaccination for all six doses of vaccines (79.3%). (**Table 3**).

### **Discussion**

The study utilized data from a large representative population-based study in NZ to investigate the impact of parental migration status and residential duration on the timely vaccination of infants. The findings revealed that even after accounting for sociodemographic variations, infants born to foreign-born parents were more likely to receive vaccination on schedule compared to those born to native-born parents. Moreover, the findings showed that infants of recent immigrants, who had lived in NZ for less than five years, had higher odds of receiving timely vaccinations compared to

infants of settled immigrants who had been residing in the country for five years or longer. These findings underscore the importance of considering migration status and residential duration when designing interventions to promote timely vaccination among infants in diverse populations.

#### **Timely infant vaccination among children of immigrant and non-immigrant families**

Our findings indicated that regardless of residential duration, infants with foreign-born mothers were more likely to have timely receipt of the six week, three-month, and five-month vaccinations as well as for all six doses of studied vaccines, compared to infants with native-born mothers. The same pattern was observed for infants with foreign-born fathers. Furthermore, infants with two immigrant parents demonstrated higher odds of receiving timely vaccinations for all scheduled vaccines compared to children with two native-born parents.

These findings align with several US-based studies [23]-[25] which similarly found that children of foreign-born parents, particularly those originating from African countries (excluding Somalia), Central and South America and the Caribbean, Mexico, or India, exhibited higher rates of immunization compared to children of US-born parents. However, it is worth noting that our findings contrast with certain other US studies [9], [23] that reported children of foreign-born parents were less likely to be vaccinated on time compared to children of native-born parents.

Differences in the composition of immigrant samples, such as inclusion of foreign-born children who may have been vaccinated overseas, into immigrant categories in some studies but not in others might have contributed to the noted inconsistencies between studies. Inclusion of such children within the immigrant category might result in underestimation of their true vaccination records due to factors such as insufficient documentation of vaccines or memory effects [11], [25], [44]. Such nuances highlight the importance of carefully considering sample composition when interpreting and comparing findings across studies.

Moreover, even after accounting for socio-demographic factors linked to children's timely vaccination, our findings indicate that children from immigrant families had higher odds of receiving vaccinations on schedule, despite the economic disadvantages often faced by immigrant families compared to non-immigrant families. This suggests the existence of potentially significant factors beyond economic barriers that contribute to the timely vaccination of children from immigrant backgrounds. To gain deeper insights into these dynamics, qualitative and in-depth research is warranted. Such research could shed light on the specific mechanisms and strategies employed by immigrant families to overcome economic barriers and ensure their children receive vaccinations on time.

The findings from this study challenge the common notion that immigration universally disrupts immigrants' access to health resources and influences their health seeking behaviors negatively. However, many demographers agree that individuals and families utilize migration as a strategy to enhance their economic [45], social [46], and cultural [47] circumstances in ways that will benefit their own and their children's well-being [48]. In this context, higher rates of timely vaccination observed among children of immigrants in the current study seems plausible, as timely immunization is recognized as a means of enhancing health conditions. This suggests that immigrant families may prioritize health-related behaviors, such as timely vaccination, as part of their efforts to improve their overall quality of life following migration.

A selection effect could also have contributed to the favorable outcomes for immigrant subgroups. It is possible that individuals who migrate are predisposed to good health or are able to demonstrate that they "have an acceptable standard of health", which may increase their likelihood of migrating [49]. Consequently, such individuals may be more inclined to seek and secure vaccinations for their children [50]. In line with this notion, research has indicated that immigrant women are often

characterized by greater motivation, higher aspirations, and stronger intentions compared to their non-immigrant counterparts—factors that could potentially enhance their children’s likelihood of timely vaccination [51].

Another plausible explanation for the higher rates of timely vaccination among immigrants could be attributed to disparities between health systems, immunization policies, and attitudes towards children’s vaccination in the country of origin versus the host country. Immigrants originating from countries where child vaccination is compulsory or culturally emphasized, may carry over these practices and attitudes when they migrate, even if vaccination is not compulsory in the host country, as is the case in NZ.

#### **Recent versus settled immigrants**

A finer stratification of immigrants into two exclusive sub-groups, based on their length of residential tenure, revealed that after adjustment for socio-demographic factors, children of recent immigrants who had resided in the country for less than five years exhibited a higher likelihood of receiving timely vaccination compared to children of settled immigrants who had been in the country for five or more years at the time of data collection. This finding holds significance as recent immigrant families tended to have a lower socio-economic status compared to settled immigrants. Despite these socio-economic disadvantages, their children displayed a higher likelihood of timely vaccination. This finding aligns with a study conducted in NZ [26] which revealed that hospitalization rates for a wide range of vaccine preventable disease were significantly lower among children from recent immigrant families compared to children from families without recent migration backgrounds.

Immigrants are known to be less integrated residents and to have competing needs, particularly during the early years following immigration, which might hinder their health seeking behaviors, consequently putting their children at elevated risk of not being fully or timely vaccinated. Indeed, some previous studies indicate that children born to mothers who recently migrated to the USA were at higher risk of not being adequately immunized compared with more settled immigrants [9].

However, recent immigrants are also generally known to be highly motivated, receptive to change, and possess greater social and cultural capital, factors that can to somewhat counteract the lack of social integration in the new country [51] and enhance their health outcomes and health seeking behaviors. This well-documented phenomena is known as the “immigrant paradox” [52], [53]. By analyzing data on recent and established immigrants, as well as data on first- and subsequent-generation immigrants, researchers have found that recent immigrants often outperform more established immigrants and non-immigrants on various health, education, and conduct or crime-related outcomes, despite the numerous barriers they face to successful social integration [54], [55].

Length of stay, which along with language proficiency has been used as a measure of acculturation in some studies [56], could potentially facilitate immigrants’ access to and uptake of primary care and preventive services such as vaccination. However, there is evidence suggesting that acculturation could come with a cost. For example, immigrants often experience a decline in health status as their years in the host community increase [57]. In the context of the current research, lower timely vaccination rates among children of settled immigrants could be due to the fact that the longer immigrant families live in NZ, the more likely they are to come into contact with vaccine-refusing parents. Although the majority of New Zealanders consistently exhibit strong

vaccine confidence, a considerable proportion shows steadily decreasing confidence over time [58].

It is also possible that as the English fluency of immigrant families improves, they become more exposed to anti-vaccine rhetoric through English-speaking media. In this scenario, increased acculturation, indicated by longer residential duration and better language proficiency, could lead to more extensive or prolonged exposure to discouraging information about vaccination. Consequently, this heightened exposure may contribute to lower rates of timely vaccination among settled immigrants.

Similarly, the higher likelihood of timely vaccination among children of recent immigrants may stem from their shorter exposure to vaccine-discouraging information received from peers or media, as well as the ambiguity created by such information. This explanation aligns with the findings of a study conducted in the USA, which revealed that childhood vaccination coverage was lower in more acculturated Latino families compared to less acculturated Latino families [59], [60]. Additionally, another US-based study found that Hispanic girls residing in predominantly Hispanic neighborhoods were more likely to be vaccinated against human papillomavirus compared to Hispanic girls residing in predominantly white neighborhoods [61].

## **Strengths and Limitations**

One of the strengths of this study is its novelty in the New Zealand context. To date, few NZ studies have explored the impact of parental migration status on children's vaccination outcomes [11]. This study builds upon previous work by utilizing a large, nationally representative sample of children and their families to examine timely vaccination disparities among children of immigrant and non-immigrant families in NZ. Additionally, the study investigates the impact of

residential duration on children's immunization timeliness, contributing to a deeper understanding of vaccination patterns within immigrant populations.

This is unique internationally, as most previous research has examined the effects of parental migration on children's immunization coverage, but little research has been conducted on immunization timeliness among immigrant sub-populations. Another important strength of this study is the utilization of administrative data from the NIR, which was linked with large, representative survey data. This integration allowed for a comprehensive analysis of vaccination patterns and their timeliness among immigrant families, providing valuable insights into the dynamics of immunization within these populations.

Another study strength is the inclusion of the migration status of the mother, father and both parents in the analyses as it provides more nuanced understanding of the impact of parental nativity on children's outcomes. Future studies could extend this work by exploring factors that enable immigrant families, and particularly recent immigrant families, to overcome barriers (such as economic disadvantages) and vaccinate their children on time.

Future studies could also extend this work by exploring disparities in parental vaccination intention during pregnancy among immigrant and non-immigrant families to see if the observed differences in children's vaccination outcomes are due to the initial parental vaccination intentions or are driven by barriers in accessing vaccination services.

Moreover, in this paper, we deliberately refrain from exploring the characteristics of the most common countries of origin from which people migrate to NZ as a factor influencing immunization outcomes. While acknowledging potential differences in health systems, immunization policies, and attitudes towards children's vaccination between countries of origin and the host country, our focus remained on understanding how migration status itself impacts timely vaccination rates. This

deliberate choice enabled us to offer a thorough examination of the influence of migration on immunization outcomes, without confounding factors related to specific countries of origin. Future research may explore country-level factors in greater detail to enrich our understanding of immunization disparities among migrant sub-populations.

This study is subject to several limitations. Due to the study design (observational) and the correlational nature of the analyses, a causal link between the parental migration status and children's timely immunization cannot be inferred. Other limitations include that we did not investigate the relationship of the immigrant families with primary health care in NZ nor with their exposure to discrimination.

Also, we were unable to identify asylum seekers and refugees due to the unavailability of data on visa types in the GUINZ dataset. This is important given that previous studies have shown that refugees are at a particular disadvantage when it comes to successful resettlement in the host society, which could be translated into poorer vaccination outcomes for their children [11], [62].

## **Conclusion**

Our study findings revealed that children of immigrant parents, identified based on both maternal and paternal migration status, in NZ demonstrated a higher likelihood of receiving timely immunization compared to children of non-immigrant parents. This suggests that immigrant groups could play a vital role in promoting vaccination and informing public health programs on strategies to ensure timely vaccine administration for all children.

However, prolonged residency in the country was associated with decreased likelihood of children receiving timely vaccination, highlighting the necessity of providing support to settled immigrants to maintain vaccination rates.

455 Additionally, when studying vaccination among immigrant children, it will be crucial to  
456 differentiate between children born overseas and children of overseas-born parents. This  
457 differentiation is necessary to understand the unique challenges and factors influencing vaccination  
458 uptake and timeliness in these distinct groups and to maximize the population health benefits  
459 derived from immunization programs.

460  
461 **Acknowledgements**

462 First and foremost, we extend our heartfelt gratitude to the children and families who generously  
463 participated in the *Growing Up in New Zealand* (GUiNZ) study. Their invaluable contributions  
464 have been instrumental in advancing our understanding of childhood vaccination behaviors.

465 We also wish to acknowledge the initial funders of GUiNZ study, notably the New Zealand  
466 Ministry of Social Development, with support from the Health Research Council. Furthermore,  
467 we express our appreciation for the ongoing support provided by Auckland UniServices and The  
468 University of Auckland, which has been crucial in sustaining the continuation of GUiNZ study.  
469 Lastly, we extend our thanks to all the members of the GUiNZ team whose dedication and hard  
470 work have been essential in the success of this project.

471 **Insert Supplementary Table 1 here**

472  
473 **References**

- 474 [1] I. D. Pavlopoulou *et al*, "Immunization coverage and predictive factors for complete and age-  
475 appropriate vaccination among preschoolers in Athens, Greece: a cross--sectional study," *BMC*  
476 *Public Health*, vol. 13, pp. 908-908, 2013. . DOI: 10.1186/1471-2458-13-908 [doi].
- 477 [2] N. Turner, "The challenge of improving immunization coverage: the New Zealand example,"  
478 *Expert Rev. Vaccines*, vol. 11, (1), pp. 9-11, 2012. . DOI: 10.1586/erv.11.157 [doi].

- 479 [3] J. P. Buttery and S. M. Graham, "Immunisation timing: the protective layer in vaccine  
480 coverage," *The Lancet (British Edition)*, vol. 373, (9674), pp. 1499-1500, 2009. Available:  
481 <https://www.clinicalkey.es/playcontent/1-s2.0-S0140673609603408>. DOI: 10.1016/S0140-  
482 6736(09)60340-8.
- 483 [4] S. C. Redd *et al*, "Comparison of vaccination with measles-mumps-rubella vaccine at 9, 12,  
484 and 15 months of age," *J. Infect. Dis.*, vol. 189 Suppl 1, pp. 116, 2004. . DOI: JID20796 [pii].
- 485 [5] H. Saffar *et al*, "Measles and rubella serosusceptibility among population vaccinated with  
486 different schedules: the potential impact on measles-rubella elimination in Iran," *BMC Infectious*  
487 *Diseases*, vol. 21, (1), pp. 305, 2021. Available:  
488 <https://www.ncbi.nlm.nih.gov/pubmed/33765968>. DOI: 10.1186/s12879-021-05970-7.
- 489 [6] C. C. Grant *et al*, "Delayed immunisation and risk of pertussis in infants: unmatched case-  
490 control study," *Bmj*, vol. 326, (7394), pp. 852-853, 2003. . DOI: 326/7394/852 [pii].
- 491 [7] C. C. Grant *et al*, "Antenatal immunisation intentions of expectant parents: Relationship to  
492 immunisation timeliness during infancy," *Vaccine*, vol. 34, (11), pp. 1379-1388, 2016. . DOI:  
493 S0264-410X(16)00100-6 [pii].
- 494 [8] H. Petousis-Harris *et al*, "What contributes to delays? The primary care determinants of  
495 immunisation timeliness in New Zealand," *J. Prim. Health. Care.*, vol. 4, (1), pp. 12-20, 2012.
- 496 [9] V. H. Buelow and J. Van Hook, "Timely immunization series completion among children of  
497 immigrants," *J. Immigr Minor. Health.*, vol. 10, (1), pp. 37-44, 2008. . DOI: 10.1007/s10903-  
498 007-9048-8 [doi].
- 499 [10] S. Karki *et al*, "Comparison of influenza vaccination coverage between immigrant and  
500 Australian-born adults," *Vaccine*, vol. 34, (50), pp. 6388-6395, 2016. . DOI: S0264-  
501 410X(16)30910-0 [pii].
- 502 [11] N. A. Charania *et al*, "Exploring immunisation inequities among migrant and refugee  
503 children in New Zealand," *Hum. Vaccin Immunother.*, vol. 14, (12), pp. 3026-3033, 2018. . DOI:  
504 10.1080/21645515.2018.1496769 [doi].
- 505 [12] D. Mipatrini *et al*, "Vaccinations in migrants and refugees: a challenge for European health  
506 systems. A systematic review of current scientific evidence," *Pathog. Glob. Health.*, vol. 111,  
507 (2), pp. 59-68, 2017. . DOI: 10.1080/20477724.2017.1281374 [doi].
- 508 [13] G. A. Paxton *et al*, "East African immigrant children in Australia have poor immunisation  
509 coverage," *J. Paediatr. Child Health*, vol. 47, (12), pp. 888-892, 2011. . DOI: 10.1111/j.1440-  
510 1754.2011.02099.x [doi].

- 511 [14] R. C. Clark and J. Mytton, "Estimating infectious disease in UK asylum seekers and  
512 refugees: a systematic review of prevalence studies," *J. Public. Health. (Oxf)*, vol. 29, (4), pp.  
513 420-428, 2007. . DOI: fdm063 [pii].
- 514 [15] N. Coppola *et al*, "Hepatitis B virus, hepatitis C virus and human immunodeficiency virus  
515 infection in undocumented migrants and refugees in southern Italy, January 2012 to June 2013,"  
516 *Euro Surveill.*, vol. 20, (35), pp. 30009-7917.ES.2015.20.35.30009, 2015. . DOI: 10.2807/1560-  
517 7917.ES.2015.20.35.30009 [doi].
- 518 [16] B. Monge-Maillo *et al*, "Screening of imported infectious diseases among asymptomatic  
519 sub-Saharan African and Latin American immigrants: a public health challenge," *Am. J. Trop.*  
520 *Med. Hyg.*, vol. 92, (4), pp. 848-856, 2015. . DOI: 10.4269/ajtmh.14-0520 [doi].
- 521 [17] G. A. Williams *et al*, "Measles among migrants in the European Union and the European  
522 Economic Area," *Scand. J. Public Health*, vol. 44, (1), pp. 6-13, 2016. . DOI:  
523 10.1177/1403494815610182 [doi].
- 524 [18] E. T. Luman *et al*, "Maternal characteristics associated with vaccination of young children,"  
525 *Pediatrics*, vol. 111, (5 Pt 2), pp. 1215-1218, 2003.
- 526 [19] L. Mollema *et al*, "An exploratory qualitative assessment of factors influencing childhood  
527 vaccine providers' intention to recommend immunization in the Netherlands," *BMC Public*  
528 *Health*, vol. 12, pp. 128-128, 2012. . DOI: 10.1186/1471-2458-12-128 [doi].
- 529 [20] E. Kpozehouen *et al*, "Improving access to immunisation for migrants and refugees:  
530 recommendations from a stakeholder workshop," *Aust. N. Z. J. Public Health*, vol. 41, (2), pp.  
531 118-120, 2017. . DOI: 10.1111/1753-6405.12602 [doi].
- 532 [21] R. Chen *et al*, "Hepatitis B birth dose vaccination rates among children in Beijing: A  
533 comparison of local residents and first and second generation migrants," *Hum. Vaccin*  
534 *Immunother.*, vol. 12, (5), pp. 1141-1148, 2016. . DOI: 10.1080/21645515.2015.1131370 [doi].
- 535 [22] J. M. Porth *et al*, "The influence of maternal migration on child vaccination in Kenya: An  
536 inverse probability of treatment-weighted analysis," *Int. J. Infect. Dis.*, vol. 106, pp. 105-114,  
537 2021. . DOI: S1201-9712(21)00285-X [pii].
- 538 [23] M. Leeds and M. H. Muscoplat, "Timeliness of Receipt of Early Childhood Vaccinations  
539 Among Children of Immigrants - Minnesota, 2016," *MMWR Morb. Mortal. Wkly. Rep.*, vol. 66,  
540 (42), pp. 1125-1129, 2017. . DOI: 10.15585/mmwr.mm6642a1 [doi].
- 541 [24] E. Wolf *et al*, "Parental Country of Birth and Childhood Vaccination Uptake in Washington  
542 State," *Pediatrics*, vol. 138, (1), pp. e20154544. doi: 10.1542/peds.2015-4544, 2016. . DOI:  
543 10.1542/peds.2015-4544 [doi].

544 [25] V. Yassenov *et al*, "Association between Parents' Nativity Status and Influenza Vaccination  
545 Rates Among Children," *SSRN Electronic Journal*, 2021. . DOI: 10.2139/ssrn.3934763.

546 [26] N. A. Charania *et al*, "Vaccine-Preventable Disease-Associated Hospitalisations Among  
547 Migrant and Non-migrant Children in New Zealand," *J. Immigr Minor. Health.*, vol. 22, (2), pp.  
548 223-231, 2020. . DOI: 10.1007/s10903-019-00888-4 [doi].

549 [27] A. K. Varan *et al*, "Vaccination Coverage Disparities Between Foreign-Born and U.S.-Born  
550 Children Aged 19-35 Months, United States, 2010-2012," *J. Immigr Minor. Health.*, vol. 19, (4),  
551 pp. 779-789, 2017. . DOI: 10.1007/s10903-016-0465-4 [doi].

552 [28] A. J. Armitage *et al*, "Health outcomes in international migrant children: protocol for a  
553 systematic review," *BMJ Open*, vol. 11, (5), pp. e041173-041173, 2021. . DOI:  
554 10.1136/bmjopen-2020-041173 [doi].

555 [29] V. Ceri *et al*, "Mental health problems of second generation children and adolescents with  
556 migration background," *Int. J. Psychiatry Clin. Pract.*, vol. 21, (2), pp. 142-147, 2017. . DOI:  
557 10.1080/13651501.2017.1291820 [doi].

558 [30] (). *2018 Census*  
559 . Available: <https://www.stats.govt.nz/2018-census/>.

560 [31] Walker *et al*, "Improving new zealand's childhood immunisation rates evidence review,"  
561 New Zealand Ministry of Health, Wellington, -07-05. 2019.

562 [32] A. Ganeshalingham *et al*, "Hospital costs of Bordetella pertussis in New Zealand children,"  
563 *N. Z. Med. J.*, vol. 129, (1445), pp. 75-82, 2016.

564 [33] S. Bonu, M. Rani and T. D. Baker, "The impact of the national polio immunization  
565 campaign on levels and equity in immunization coverage: evidence from rural North India,"  
566 *Social Science & Medicine* (1982), vol. 57, (10), pp. 1807-1819, 2003. Available:  
567 [https://dx.doi.org/10.1016/S0277-9536\(03\)00056-X](https://dx.doi.org/10.1016/S0277-9536(03)00056-X). DOI: 10.1016/S0277-9536(03)00056-X.

568 [34] J. Whelan *et al*, "Incidence of acute hepatitis B in different ethnic groups in a low-endemic  
569 country, 1992-2009: increased risk in second generation migrants," *Vaccine*, vol. 30, (38), pp.  
570 5651-5655, 2012. . DOI: 10.1016/j.vaccine.2012.06.080 [doi].

571 [35] S. M. B. Morton *et al*, "Cohort profile: Growing Up in New Zealand," *International Journal*  
572 *of Epidemiology*, vol. 42, (1), pp. 65-75, 2013. Available:  
573 <https://www.ncbi.nlm.nih.gov/pubmed/22253317>. DOI: 10.1093/ije/dyr206.

574 [36] S. M. B. Morton *et al*, "Growing Up in New Zealand cohort alignment with all New  
575 Zealand births," *Australian and New Zealand Journal of Public Health*, vol. 39, (1), pp. 82-87,  
576 2015. Available: <https://onlinelibrary.wiley.com/doi/abs/10.1111/1753-6405.12220>. DOI:  
577 10.1111/1753-6405.12220.

578 [37] Ministry of Health, *Immunisation Handbook*. Wellington: Ministry of Health, 2020.

579 [38] V. J. Dietz, E. R. Zell and J. Stevenson, "Defining delayed immunization," *Pediatr. Infect.*  
580 *Dis. J.*, vol. 12, (4), pp. 353-354, 1993. . DOI: 10.1097/00006454-199304000-00022 [doi].

581 [39] C. Wallace, J. Leask and L. J. Trevena, "Effects of a web based decision aid on parental  
582 attitudes to MMR vaccination: a before and after study," *Bmj*, vol. 332, (7534), pp. 146-148,  
583 2006. Available: <http://dx.doi.org/10.1136/bmj.38678.681840.68>. DOI:  
584 10.1136/bmj.38678.681840.68.

585 [40] New Zealand Ministry of Business, Innovation, and Employment, "Migrant youth statistical  
586 profile," y Ministry of Business, Innovation and Employment, Wellington, Sep. 2017.

587 [41] S. M. B. Morton *et al*, "Growing up in new zealand: A longitudinal study of new zealand  
588 children and their families. report 2: Now we are born ," Auckland: Growing Up in New  
589 Zealand, March. 2012.

590 [42] C. Salmond, P. Crampton and J. Atkinson, "NZDep2006 index of deprivation," Department  
591 of Public Health, University of Otago, Wellington, Wellington, New Zealand, 2007.

592 [43] StataCorp, "Stata Statistical Software: Release 14," 2015.

593 [44] S. Stokley, L. E. Rodewald and E. F. Maes, "The impact of record scattering on the  
594 measurement of immunization coverage," *Pediatrics*, vol. 107, (1), pp. 91-96, 2001. . DOI:  
595 10.1542/peds.107.1.91 [doi].

596 [45] M. P. Todaro, "Income expectations, rural-urban migration and employment in Africa," *Int.*  
597 *Labour Rev.*, vol. 135, (3-4), pp. 421-444, 1996.

598 [46] D. S. Massey, "Social structure, household strategies, and the cumulative causation of  
599 migration," *Popul. Index*, vol. 56, (1), pp. 3-26, 1990.

600 [47] W. Kandel and D. S. Massey, "The culture of Mexican migration: A theoretical and  
601 empirical analysis," *Social Forces*, vol. 80, (3), pp. 981-1004, 2002. . DOI:  
602 10.1353/sof.2002.0009.

603 [48] G. F. DE Jong, "Expectations, gender, and norms in migration decision-making," *Popul.*  
604 *Stud. (Camb)*, vol. 54, (3), pp. 307-319, 2000. . DOI: 10.1080/713779089 [doi].

605 [49] (Dec 8). *Meeting the criteria*. Available: <https://www.immigration.govt.nz/new-zealand-visas/apply-for-a-visa/criteria/skilled-migrant-category-resident-visa?nationality=nationality-CHE&country=residence-CHE&applying=co>.  
606  
607

- 608 [50] J. Kraśnicka *et al*, "The impact of parents' health behaviours on their preferences regarding  
609 vaccinations in Białystok, Poland," *BMC Pediatr.*, vol. 20, (1), pp. 354-1, 2020. . DOI:  
610 10.1186/s12887-020-02235-1 [doi].
- 611 [51] G. E. Kiros and M. J. White, "Migration, community context, and child immunization in  
612 Ethiopia," *Soc. Sci. Med.*, vol. 59, (12), pp. 2603-2616, 2004. . DOI: S0277953604001881 [pii].
- 613 [52] S. L. Hofferth and U. J. Moon, "How do they do it? The immigrant paradox in the transition  
614 to adulthood," *Soc. Sci. Res.*, vol. 57, pp. 177-194, 2016. . DOI: S0049-089X(16)00012-0 [pii].
- 615 [53] L. Zhang, A. Bo and W. Lu, "To Unfold the Immigrant Paradox: Maltreatment Risk and  
616 Mental Health of Racial-Ethnic Minority Children," *Frontiers in Public Health*, vol. 9, pp.  
617 619164, 2021. Available: <https://www.ncbi.nlm.nih.gov/pubmed/33681132>. DOI:  
618 10.3389/fpubh.2021.619164.
- 619 [54] C. P. Salas-Wright *et al*, "Substance use disorders among first- and second- generation  
620 immigrant adults in the United States: evidence of an immigrant paradox?" *J. Stud. Alcohol.*  
621 *Drugs*, vol. 75, (6), pp. 958-967, 2014. . DOI: 958 [pii].
- 622 [55] A. E. Vaughn *et al*, "Fundamental constructs in food parenting practices: a content map to  
623 guide future research," *Nutr. Rev.*, vol. 74, (2), pp. 98-117, 2016. . DOI: 10.1093/nutrit/nuv061  
624 [doi].
- 625 [56] A. G. Yong *et al*, "Acculturation in preventive health for immigrants: A systematic review  
626 on influenza vaccination programs in a socio-ecological framework," *Canadian*  
627 *Psychology/Psychologie Canadienne*, vol. 57, (4), pp. 340-355, 2016. . DOI:  
628 10.1037/cap0000075.
- 629 [57] Canada Health, "Migration health: Embracing a determinants of health approach," Health  
630 Policy Research Bulletin, Ottawa, Tech. Rep. H12-36/17–2010, 2010.
- 631 [58] C. H. J. Lee and C. G. Sibley, "Attitudes toward vaccinations are becoming more polarized  
632 in New Zealand: Findings from a longitudinal survey," *EClinicalMedicine*, vol. 23, pp. 100387,  
633 2020. . DOI: 10.1016/j.eclinm.2020.100387 [doi].
- 634 [59] L. M. Anderson, D. L. Wood and C. D. Sherbourne, "Maternal acculturation and childhood  
635 immunization levels among children in Latino families in Los Angeles," *Am. J. Public Health*,  
636 vol. 87, (12), pp. 2018-2021, 1997. . DOI: 10.2105/ajph.87.12.2018 [doi].
- 637 [60] S. E. Panameno, "Survey of Vaccine Views in Hispanic Parents (SOVVIHP)," 2021.  
638 Available: <https://escholarship.org/uc/item/7j68d6v9>.
- 639 [61] K. A. Henry *et al*, "Geographic Factors and Human Papillomavirus (HPV) Vaccination  
640 Initiation among Adolescent Girls in the United States," *Cancer Epidemiol.Biomarkers Prev.*,  
641 vol. 25, (2), pp. 309-317, 2016. . DOI: 10.1158/1055-9965.EPI-15-0658 [doi].

642 [62] P. J. Masters *et al*, "Health issues of refugees attending an infectious disease refugee health  
643 clinic in a regional Australian hospital," *Aust.J.Gen.Pract.*, vol. 47, (5), pp. 305-310, 2018. .  
644 DOI: 10.31128/AFP-10-17-4355 [doi].