



City Research Online

City St George's, University of London

Citation: Press, C., Berlot, E., Bird, G., Ivry, R. & Cook, R. (2014). Moving time: The influence of action on duration perception. JOURNAL OF EXPERIMENTAL PSYCHOLOGY-GENERAL, 143(5), pp. 1787-1793. doi: 10.1037/a0037650

This is the accepted version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/4541/>

Link to published version: <https://doi.org/10.1037/a0037650>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

In press at *Journal of Experimental Psychology: General*

Moving time: The influence of action on duration perception

Clare Press^{1*}, Eva Berlot², Geoff Bird³, Richard Ivry⁴, and Richard Cook⁵

* Corresponding author

Phone: (+44) 20 3073 8007

Email: c.press@bbk.ac.uk

¹Department of Psychological Sciences, Birkbeck, University of London,
Malet Street, London, UK

²Division of Psychology and Language Sciences, UCL, Bedford Way, London, UK

³MRC Social, Genetic, and Developmental Psychiatry Centre, Institute of Psychiatry, De
Crespigny Park, London, UK

⁴Department of Psychology, UC Berkeley, Tolman Hall, Berkeley, US.

⁵Department of Psychology, City University London,
Whiskin Street, London, UK

Word count (excl. title, references, affiliations, acknowledgments, figures, incl. abstract):

2997

Abstract

Perceiving the sensory consequences of action accurately is essential for appropriate interaction with our physical and social environments. Prediction mechanisms are considered necessary for fine-tuned sensory control of action, yet paradoxically may distort perception. Here we examine this paradox by addressing how movement influences the perceived duration of sensory outcomes congruent with action. Experiment 1 required participants to make judgments about the duration of vibrations applied to a moving or stationary finger. In Experiments 2 and 3, participants judged observed finger movements, congruent or incongruent with their own actions. In all experiments, target events were perceived as longer when congruent with movement. Interestingly, this temporal dilation did not differ as a function of stimulus perspective (first or third person) or spatial location. We propose that this bias may reflect the operation of an adaptive mechanism for sensorimotor selection and control that pre-activates anticipated outcomes of action. The bias itself may have surprising implications both for action control and perception of others - we may be in contact with grasped objects for less time than we realize and others' reactions to us may be briefer than we believe.

Keywords: Motor Processes, Perceptual Motor Coordination, Time Perception, Social Perception

Introduction

To interact appropriately with physical and social environments, actors must predict and evaluate the sensory consequences of their actions. We select actions based on their predicted outcomes (Greenwald, 1970; Hommel, Müsseler, Aschersleben, & Prinz, 2001), and when the experienced sensory information deviates from our prediction, corrective adjustments can be made ensuring successful execution. For example, when picking up a cup of tea, the motor commands generated result in both visual (e.g., sight of grasping and lifting) and tactile (e.g., pressure on the fingertips) sensory consequences. If the actual feedback differs from the anticipated sensory outcomes, rapid corrective actions can be executed to avoid spillage. Similarly, when interacting with others, rapid response prediction and error correction may facilitate smooth social interactions (Wolpert, Doya, & Kawato, 2003).

Successful interaction with the environment requires perception not only of the nature of our action outcomes (e.g., somatosensation on the fingertips during grasping), but also crucially, the onset and duration of those outcomes. We are sensitive both to the ‘what’ and ‘when’ of sensory predictions (Blakemore, Frith, & Wolpert, 1999; Blakemore, Wolpert, & Frith, 1998; Christensen, Ilg, & Giese, 2011; Fagioli, Hommel, & Schubotz, 2007; Hommel, 2010; Lee, Young, Reddish, Lough, & Clayton, 1983; Schubotz, 2007). For example, lifting the teacup from a saucer requires an anticipatory response to maintain postural stability (Diedrichsen, Verstynen, Hon, Lehman, & Ivry, 2003; Dufossé, Hugon, & Massion, 1985). Similarly, anticipating the duration of the lift phase is essential for coordinating hand and mouth gestures, and when shaking someone’s hand, contact must be made for an appropriate length of time, neither too long, nor too short, to convey the intended social message.

While the preceding examples underscore the importance of temporal information in the generation and perception of sensory expectancies, duration perception is frequently distorted. For example, perceived motion of upright point light walkers is temporally dilated relative to inverted walkers (Wang & Jiang, 2012; see also Gavazzi, Bisio, & Pozzo, 2013). The present experiments examine how movement influences the perceived duration of sensory outcomes of action. Sensory prediction mechanisms essential for action selection and fine-tuned control may, paradoxically, distort the

perceived duration of outcomes, with consequences for action-control and perception in a variety of contexts. In Experiment 1, participants were required to perform a lifting movement with either their index or middle finger. A short target vibratory tactile stimulus was presented to the moving or stationary finger, followed by a second reference vibration. Participants judged which was of longer duration, allowing us to determine how action influences duration perception. Experiments 2-4 asked whether action influences the perceived duration of visual events in a comparable manner.

Experiment 1

Sixteen right-handed, healthy adults (12 male) with a mean age of 29.3 years (SEM = 2.5) participated in the experiment. Three were replacements for participants who could not perform the perceptual discrimination (psychometric functions could not be modeled or their point of subjective equivalence (PSE) fell outside the range of presented stimuli). All experiments were undertaken in accordance with the 1964 Declaration of Helsinki.

The experiment was conducted in MATLAB using the Cogent toolboxⁱ. Two 5V solenoids, each driving a metal rod with a blunt conical tip (diameter $\approx$ 1.5 mm, skin indentation $\approx$ 1 mm), were attached to the distal phalange (ventral surface) of the index and middle fingers on the right hand. Participants held down two keys on the keyboard until an imperative cue instructed them to lift either their index ('I') or middle ('M') finger. They were instructed to make large, rapid, single-movement lifts. Their response hand was visually occluded. Approximately 10 ms after the cued finger was lifted, a target vibration lasting for one of seven durations (104 – 296 ms, 32 ms steps) was applied to the moving ('congruent') or stationary ('incongruent') finger (see Fig. 1). After an inter-stimulus interval (ISI; 300 – 500 ms), a 200 ms reference vibration was applied to the same finger. Both vibratory stimuli were presented at 62.5 Hz.

Participants judged whether the target or reference vibration was longer, responding with a button press made with their left hand. Following this response, they returned the lifted finger to the start position. The next trial started after 2000 ms. There were 280 trials; 140 in which stimuli were applied to the congruent finger and 140 where

they were applied to the incongruent finger. Trial type was randomized and participants completed 8 practice trials.

To estimate psychometric functions, the responses for each individual were modeled by fitting cumulative Gaussians, and associated pDev statistics were calculated to establish the goodness-of-fit of each function (Palamedes toolbox, Kingdom & Prins, 2010). This procedure was performed separately for congruent and incongruent response data. In each condition, bias was inferred from the PSE and precision from the difference threshold (Fig. 2).

The participants were more precise in their judgments when the vibratory stimuli were applied to the congruent relative to incongruent finger ($t(15) = 2.3$, $p < 0.05$, $\eta^2 = 0.26$; Table 1). There was also an effect on PSE: Target events were judged longer when the stimulus was applied to the congruent relative to incongruent finger ($t(15) = 2.6$, $p < 0.02$, $\eta^2 = 0.32$; Fig. 2 & Supplementary Fig. 1). In sum, tactile events presented to a moving effector are perceived to be longer and are judged more consistently than when that effector is stationary.

Experiment 2

If prediction mechanisms operate in social contexts, we may predict and evaluate sensation not only related to our own actions, but also actions produced by interactants (Wolpert et al., 2003). As such, we should observe comparable action-related predictive modulation with visual action stimulus events. Additionally, such mechanisms should operate across perspectives given the range of viewpoints from which others' actions are observed. Experiment 2 therefore examined duration perception of congruent and incongruent visual events during action, across stimuli presented from first and third person perspectivesⁱⁱ.

Sixteen right-handed, healthy adults (12 male) with a mean age of 25.9 years (SEM = 1.9) participated in the experiment. Five were replacements for participants who could not perform the discrimination. Unless otherwise stated, procedural information already outlined in Experiment 1 is identical in this, and all subsequent, experiments. Participants compared the duration of two finger movements simulated visually by

gestures of an avatar hand. At the start of the trial, the avatar hand was presented in a neutral position on the monitor (Fig. 1; screen refresh rate = 85 Hz). An imperative cue ('1' or '2') was presented between the index and middle fingers. When participants lifted the cued finger, the neutral hand image was immediately replaced (within the constraints of the refresh rate) by one depicting the avatar hand performing either an index or middle finger lift for 120 - 480 ms (7 levels; 60 ms steps). This resulted in apparent motion of the avatar's finger approximately synchronized with the participant's action. At the offset of the target event, the avatar hand resumed the neutral position for an ISI of 300 - 500 ms, followed by a second image of the same lifted finger for a reference duration of 300 ms, and then the neutral image again (300 ms). Participants judged which lift lasted longer. The range of durations was chosen to match discrimination performance in Experiment 1.

There were four block types. In spatially aligned first person perspective (1PP) blocks, participants viewed a right avatar hand with fingers aligned in the horizontal plane with their own right hand (Fig. 1). In spatially aligned third person perspective (3PP) blocks, the avatar hand was rotated about the horizontal axis (therefore presenting a left hand). The remaining blocks consisted of these stimuli flipped on a vertical axis, such that corresponding finger movements did not match in spatial location (necessitating left hand for 1PP and right hand for 3PP). These blocks thereby controlled for the spatial location of finger movement (Press, Gherri, Heyes, & Eimer, 2010). The four blocks each comprised 140 trials and were completed in a counterbalanced order.

The precision and PSE distributions were analyzed using separate three-way ANOVAs, with factors of movement congruency (avatar and participant moved the congruent/incongruent finger), location (avatar and participant finger movements made in aligned/misaligned horizontal locations), and perspective (1PP or 3PP). No precision effects were observed (all F s < 2.1, all p s > 0.17; Table 1). However, as in Experiment 1, target events were perceived as longer when the avatar and participant moved the same finger ($F(1,15) = 5.3$, $p < 0.04$, $\eta^2 = 0.26$). There were no other main effects or interactions (all F s < 2.5, all p s > 0.14). These results indicate a bias to judge target events as longer when observed actions are congruent with self-generated actions, regardless of whether stimuli are observed from first or third person

perspectives. Notably these effects reflect congruency between effectors (same finger) rather than spatial location.

Experiment 3

Experiment 2 suggests that action performance influences the perceived duration of effector-congruent visual events. However, it is possible that, despite informing participants that the reference event was always presented for the same length of time, participants' actions might have influenced the perceived duration of the reference rather than the target event. To control for this possibility, the reference event was modified in Experiment 3. Rather than define the reference duration by a second avatar movement, this interval was defined by the stimulus duration of a rectangle, a neutral stimulus selected because it exhibited no congruency relationship with the fingers.

Sixteen right-handed, healthy adults (11 male) with a mean age of 28.3 years ($SEM = 2.2$) participated in the experiment. Three were replacements for participants who could not perform the discrimination. The imperative cue ('I' or 'M') was presented between the index and middle fingers of the observed hand. When participants lifted the cued finger, the neutral hand image was immediately replaced by an image of a hand with a lifted index or middle finger for 150 – 900 ms (7 levels; 125 ms steps). Following an ISI of 300 - 500 ms, a rectangle was presented for a reference interval of 525 ms. The color, luminance, and aspect ratio of the rectangle were identical to that of the avatar hand. The test stimulus range was selected based on piloting to yield comparable performance to that observed in Experiments 1 and 2. Participants again judged which of the two intervals was longer. Given that spatial location had no impact on the effect in Experiment 2, only aligned blocks were included. Participants completed 1PP and 3PP blocks, each comprising 140 trials, in a counterbalanced order.

The precision analysis yielded no main effects or interactions (all $F_s < 1.4$, all $p_s > 0.25$; see Table 1). However, the PSE phenomenon observed in Experiments 1 and 2 was replicated: Target events were perceived as longer when the observed event was congruent with the participant's action ($F(1,15) = 6.5, p < 0.03, \eta^2 = 0.30$; see Fig. 2). As in Experiment 2, this effect did not interact with perspective ($F(1,15) = 0.05, p = 0.8, \eta^2$

= 0.02). These findings demonstrate that action biases perception of the temporally contiguous target event, rather than reference events presented after a delay.

Experiment 4

It is possible that the temporal dilation effects in Experiments 2 and 3 result from attentional orienting towards the location of the congruent effector. Increased attention may influence the perceived duration of events at this location irrespective of action-stimulus congruency. A final experiment was conducted to test this possibility, identical to Experiment 3, except that target durations were defined by the presentation of a rectangle over the fingertip of the index or middle finger rather than by a finger movement (see Fig. 1). If attentional orienting generates temporal dilation effects irrespective of the nature of the target event, similar influences of congruency will be observed.

Sixteen right-handed, healthy adults (7 male) with a mean age of 27.7 years (SEM = 1.7) participated in the experiment. One was a replacement for a participant who could not perform the discrimination. The precision analysis yielded no main effects or interactions (all F s < 0.7, all p s > 0.41; see Table 1). Unlike Experiments 1-3, there was also no PSE effect ($F(1,15) = 0.7$, $p = 0.42$, see Fig. 2). ANOVA conducted on the PSE data from both Experiments 3 and 4, with experiment as a between-subjects factor, revealed a congruency $\times$ experiment interaction ($F(1,30) = 6.8$, $p < 0.02$, $\eta^2 = 0.2$). These results argue against this attentional orienting account of the congruency-induced temporal dilation.

Discussion

The present findings demonstrate a bias to judge sensory events as longer when congruent with a concurrently performed action. This effect was found when participants judged the duration of tactile vibrations applied to a moving finger, as well as when assessing the duration of observed finger movementsⁱⁱⁱ. These results indicate that subjective action-time can be subject to temporal dilation: Events effector-congruent with performed actions are perceived as longer than events incongruent with those actions.

These effects may be a consequence of pre-activated action expectancies during selection and preparation (Greenwald, 1970; Hommel et al., 2001), whereby congruent sensory events are perceived to begin before action onset. Imperfect distinctions between anticipated and actual sensory consequences would cause congruent sensation to be perceived as longer. In contrast, when action consequences are unexpected, pre-activated outcomes differ from the actual sensory consequences and can thus be discriminated. The hypothesis that duration biases result from imperfect distinctions between predicted and stimulus-driven percepts is consistent with the finding that imagined and perceived visual events activate common occipital representations (Kosslyn et al., 1993; Albers, Kok, Toni, Dijkerman, & de Lange, 2013; see also Buetti & Macaluso, 2010), and that action preparation activates representations of the anticipated effects (Müsseler & Hommel, 1997; Kühn, Keizer, Rombouts, & Hommel, 2011). Furthermore, the idea that the perceived onset of anticipated events is shifted in time is consistent with a number of temporal distortions in the action control literature. For example, it has long been recognized that, when tapping to a metronome, movements show a phase lead to the pacing signals (Dunlap, 1910; Bartlett & Bartlett, 1959). Moreover, effects resulting from action but at delay are perceived to occur earlier than in reality (Haggard, 2005).

Temporal biases resulting from the prediction of congruent action consequences might be expected to detract from effective action control. However, illusory biases often result from the operation of adaptive mechanisms. For example, visual aftereffects, defined by significant sensory distortion, are believed to be the products of ongoing perceptual recalibration to ambient sensory inputs (Clifford & Rhodes, 2005; Thompson & Burr, 2009; see also Yarrow, Haggard, Heal, Brown, & Rothwell, 2001). Similarly, stimulus-general temporal dilation during action planning may maximize information acquisition prior to movement (Hagura, Kanai, Orgs, & Haggard, 2012). Following this line of reasoning, we suggest that the dilation of subjective action-time observed for anticipated sensory outcomes may be indicative of an adaptive mechanism optimized for online action control. Anticipation of the sensory consequences of action is essential for action selection and subsequent error correction. Imperfect distinction between anticipated and actual sensory outcomes may reflect exploitation of mechanisms adapted for perception during action planning. While these mechanisms broadly benefit

actors, there may be surprising consequences for tightly time-locked action control and social perception. For example, we may be in contact with grasped objects for less time than we realize and handshakes may be briefer than we believe.

Equivalent effects when observing sensory events from first and third person perspectives suggests that common mechanisms anticipate the consequences of our own actions as well as the imitative reactions of others. Wolpert et al. (2003) proposed that sensory prediction mechanisms for action control may also operate when interacting with others, but this possibility has received little empirical investigation. The present study provides support for this hypothesis, suggesting that we overestimate the duration, not only of our own actions, but also others' imitative reactions. Future investigations must establish whether these effects are found when other individuals react in a non-imitative, but predictable, manner; for example, when dominant body postures result in complementary submissive postures of an interactant (Tiedens & Fragale, 2003).

Neuropsychological and neuroimaging studies have implicated motor structures in duration perception, even when action is not required. For example, the cerebellum and basal ganglia are thought to play key roles in a range of temporal judgments (Harrington, Haaland, & Hermanowicz, 1998; Ivry & Keele, 1989; Ivry, Spencer, Zelaznik, & Diedrichsen, 2002; Koch et al., 2007). Additionally, greater activation has been observed in cortical motor areas, including the supplementary motor area (SMA) and dorsal premotor cortex, when judging the duration of visual events (Coull, Nazarian, & Vidal, 2008; Ferrandez et al., 2003), than when making intensity or color judgments about the same stimuli. These duration judgments may recruit the motor system to exploit mechanisms adapted, either phylogenetically or ontogenetically (Heyes, 2003), for action control.

302 **Acknowledgments**

303 This research was funded by a Wellcome Trust Biomedical Studentship awarded to EB
304 and CP, a Birkbeck School of Science grant awarded to CP, and an ESRC Future Research
305 Leaders grant (ES/K008226/1) awarded to RC. We are grateful to Martin Eimer for the
306 loan of solenoids.

307

References

- Albers, A.M., Kok, P., Toni, I., Dijkerman, H.C., & de Lange, F.P. (2013). Shared representations for working memory and mental imagery in early visual cortex. *Current Biology*, 23, 1-5. doi: 10.1016/j.cub.2013.05.06
- Bartlett, N.R., & Bartlett, S.C. (1959). Synchronization of a motor response with an anticipated sensory event. *Psychological Review* 66(4), 203-218.
- Blakemore, S. J., Frith, C. D., & Wolpert, D. M. (1999). Spatio-temporal prediction modulates the perception of self-produced stimuli. *Journal of Cognitive Neuroscience*, 11(5), 551-559.
- Blakemore, S. J., Wolpert, D. M., & Frith, C. D. (1998). Central cancellation of self-produced tickle sensation. *Nature Neuroscience*, 1(7), 635-640. doi:10.1038/2870
- Bueti, D., & Macaluso, E. (2010). Auditory temporal expectations modulate activity in visual cortex. *NeuroImage*, 51(3), 1168-1183. doi: 10.1016/j.neuroimage.2010.03.023
- Christensen, A., Ilg, W., & Giese, M. A. (2011). Spatiotemporal Tuning of the Facilitation of Biological Motion Perception by Concurrent Motor Execution. *Journal of Neuroscience*, 31(9), 3493-3499. doi:10.1523/JNEUROSCI.4277-10.2011
- Clifford, C. W. G., & Rhodes, G. (2005). *Fitting the mind to the world: Adaptation and after-effects in high-level vision*. Oxford University Press.
- Coull, J. T., Nazarian, B., & Vidal, F. (2008). Timing, storage, and comparison of stimulus duration engage discrete anatomical components of a perceptual timing network. *Journal of Cognitive Neuroscience*, 20(12), 2185-2197. doi:10.1162/jocn.2008.20153
- Diedrichsen, J., Verstynen, T., Hon, A., Lehman, S. L., & Ivry, R. B. (2003). Anticipatory adjustments in the unloading task: Is an efference copy necessary for learning? *Experimental Brain Research*, 148(2), 272-276. doi:10.1007/s00221-002-1318-z
- Dufossé, M., Hugon, M., & Massion, J. (1985). Postural forearm changes induced by predictable in time or voluntary triggered unloading in man. *Experimental Brain Research*, 60(2), 330-334. doi:10.1007/BF00235928
- Dunlap, K. (1910). Reaction to rhythmic stimuli with attempt to synchronize. *Psychological Review*, 17(6), 399-416.
- Fagioli, S., Hommel, B., & Schubotz, R.I. (2007). Intentional control of attention: Action planning primes action-related stimulus dimensions. *Psychological Research*, 71, 22-29. doi: 10.1007/s00426-005-0033-3.
- Ferrandez, A. ., Hugueville, L., Lehericy, S., Poline, J. ., Marsault, C., & Pouthas, V. (2003). Basal ganglia and supplementary motor area subtend duration perception: an fMRI study. *NeuroImage*, 19(4), 1532-1544. doi:10.1016/S1053-8119(03)00159-9
- Gavazzi, G., Bisio, A., & Pozzo, T. (2013). Time perception of visual motion is tuned by the motor representation of human actions. *Scientific Reports*, 3, 1168. doi: 10.1038/srep01168
- Greenwald, A. G. (1970). Sensory feedback mechanisms in performance control: With special reference to the ideo-motor mechanism. *Psychological Review*, 77(2), 73-99. doi:10.1037/h0028689

- 348 Haggard, P. (2005). Conscious intention and motor cognition. *Trends in Cognitive Sciences*, 9,
349 290-295. doi: 10.1016/j.tics.2005.04.012
- 350 Hagura, N., Kanai, R., Orgs, G., & Haggard, P. (2012) Ready steady slow: action preparation slows
351 the subjective passage of time. *Proceedings of the Royal Society of London, B*. 279(1746),
352 4399-4406. doi: 10.1098/rspb.2012.1339
- 353 Harrington, D. L., Haaland, K. Y., & Hermanowicz, N. (1998). Temporal processing in the basal
354 ganglia. *Neuropsychology*, 12(1), 3-12.
- 355 Heyes, C. (2003). Four routes of cognitive evolution. *Psychological Review*, 110(4), 713-727.
356 doi:10.1037/0033-295X.110.4.713
- 357 Hommel, B., Müsseler, J., Aschersleben, G., & Prinz, W. (2001). The Theory of Event Coding
358 (TEC): A framework for perception and action planning. *Behavioral and Brain Sciences*,
359 24(05), 849-878. doi:10.1017/S0140525X01000103
- 360 Hommel, B. (2010). Grounding attention in action control: The intentional control of selection.
361 In B.J Bruya (ed.), *Effortless attention: A new perspective in the cognitive science of attention*
362 *and action* (pp.121-140). Boston, MA: MIT Press.
- 363 Ivry, R. B., & Keele, S. W. (1989). Timing functions of the cerebellum. *Journal of Cognitive*
364 *Neuroscience*, 1(2), 136-152. doi:10.1162/jocn.1989.1.2.136
- 365 Ivry, R. B., Spencer, R. M., Zelaznik, H. N., & Diedrichsen, J. (2002). The cerebellum and event
366 timing. *Annals of the New York Academy of Sciences*, 978(1), 302-317. doi:10.1111/j.1749-
367 6632.2002.tb07576.x
- 368 Kingdom, F.A.A., & Prins, N. (2010). *Psychophysics: A practical introduction*. London, UK:
369 Elsevier.
- 370 Koch, G., Oliveri, M., Torriero, S., Salerno, S., Lo Gerfo, E., & Caltagirone, C. (2007). Repetitive TMS
371 of cerebellum interferes with millisecond time processing. *Experimental Brain Research*,
372 179(2), 291-299. doi:10.1007/s00221-006-0791-1
- 373 Kosslyn, S.M., Alpert, N.M., Thompson, W.L., Maljkovic, V., Weise, S.B., Chabris, C.F., et al. (1993).
374 Visual mental imagery activates topographically organized visual cortex: PET investigations.
375 *Journal of Cognitive Neuroscience*, 5(3), 263-287. doi:10.1162/jocn.1993.5.3.263
- 376 Kühn, S., Keizer, A., Rombouts, S.A.R.B., & Hommel, B. (2011). The functional and neural
377 mechanism of action preparation: Roles of EBA and FFA in voluntary action control. *Journal*
378 *of Cognitive Neuroscience*, 23, 214-220. doi: 10.1162/jocn.2010.21418
- 379 Lee, D. N., Young, D. S., Reddish, P. E., Lough, S., & Clayton, T. M. H. (1983). Visual timing in
380 hitting an accelerating ball. *The Quarterly Journal of Experimental Psychology Section A*,
381 35(2), 333-346. doi:10.1080/14640748308402138
- 382 Müsseler, J., & Hommel, B. (1997). Blindness to response-compatible stimuli. *Journal of*
383 *Experimental Psychology: Human Perception and Performance*, 23, 861-872. doi:
384 10.1037/0096-1523.23.3.861
- 385 Press, C., Gherri, E., Heyes, C., & Eimer, M. (2010). Action preparation helps and hinders
386 perception of action. *Journal of Cognitive Neuroscience*, 22(10), 2198-2211.
387 doi:10.1162/jocn.2009.21409

388 Schubotz, R.I. (2007). Prediction of external events with our motor system: towards a new
389 framework. *Trends in Cognitive Sciences*, 11(5), 211-218. doi: 10.1016/j.tics.2007.02.006

390 Thompson, P., & Burr, D. (2009). Visual aftereffects. *Current Biology*, 19(1), R11-R14.
391 doi:10.1016/j.cub.2008.10.014

392 Tiedens, L.Z., & Fragale, A.R. (2003). Power moves: Complementarity in dominant and
393 submissive nonverbal behaviour. *Journal of Personality and Social Psychology*, 84(3),
394 558-568. doi:10.1037/0022-3514.84.3.558

395 Wang, L. & Jiang, Y. (2012). Life motion signals lengthen perceived temporal duration.
396 *Proceedings of the National Academy of Sciences*, 109(11), 673-677. doi:
397 10.1073/pnas.1115515109

398 Wolpert, D. M., Doya, K., & Kawato, M. (2003). A unifying computational framework for motor
399 control and social interaction. *Philosophical Transactions of the Royal Society of London.*
400 *Series B: Biological Sciences*, 358(1431), 593-602. doi:10.1098/rstb.2002.1238

401 Yarrow, K., Haggard, P., Heal, R., Brown, P., & Rothwell, J. C. (2001). Illusory perceptions of space
402 and time preserve cross-saccadic perceptual continuity. *Nature*, 414(6861), 302-305.
403 doi:10.1038/35104551

404




Figure 1. The visual stimuli (created using Smith Micro Software's Poser 7.0) and timecourse for the action-related events in each of the four experiments. Timecourse stimuli depict the avatar hand in first person perspective.




Figure 2. Top panel: Demonstration of how the PSE was calculated with psychometric functions for an example participant, with stimuli congruent and incongruent with moving fingers. The PSE describes the point where participants judge the target and reference events as having equal duration. Judgment precision was inferred from the standard deviation of the Gaussian distribution which best fits the data; it pertains to the inverse of the slope, and lower thresholds reflect more consistent categorizations, thereby indicating better performance. Other panels: Mean PSEs for stimuli congruent and incongruent with moving fingers, for all experiments and perspectives. 1PP = first person perspective, 3PP = third person perspective. Error bars represent the standard error of the mean.

420

	Congruent	Incongruent
Experiment 1: Tactile	107.5 (52.3)	129.4 (68.5)
Experiment 2: Visual – 1PP	100.6 (13.5)	100.2 (12.1)
Experiment 2: Visual – 3PP	100.5 (14.5)	111.3 (14.8)
Experiment 3: Visual – 1PP	379.9 (67.1)	346.8 (48.2)
Experiment 3: Visual – 3PP	330.3 (40.6)	287.3 (35.2)
Experiment 4: 1PP	294.2 (21.9)	283.1 (28.6)
Experiment 4: 3PP	318.1 (36.5)	319.5 (37.7)

421 **Table 1. Mean precision estimates for stimuli congruent and incongruent with moving fingers, shown**
422 **separately for each experiment and perspective. Standard error of the mean is displayed in brackets in each**
423 **condition. 1PP = first person perspective, 3PP = third person perspective.**
424

425

426 **FOOTNOTES**

427 ⁱ Developed by the Cogent 2000 team at the FIL and the ICN and Cogent Graphics
428 developed by John Romaya at the LON at the Wellcome Trust Centre for Neuroimaging.

429 ⁱⁱ Examining congruency-induced temporal dilation in the visual modality also permits
430 better isolation of perceptual effects from the direct effects of action performance.
431 Visually-defined congruency is eliminated when the hands are occluded. It is not
432 possible to eliminate tactile-defined congruency without some form of sensory
433 deafferentation.

434 ⁱⁱⁱ This similarity was observed across experiments despite changes in the range of
435 durations presented. It is worth noting that piloting indicated these shifts in duration to
436 be necessary for two reasons. First, the apparent motion in Experiments 2 and 3 did not
437 appear natural with short durations. Second, the duration judgments became more
438 difficult across experiments, moving from punctate touch to apparent motion in vision
439 in Experiment 2, and changing the nature of the reference relative to the target in
440 Experiment 3. Given these changes to the durations presented in Experiments 1-3, it is
441 difficult to draw conclusions concerning the presence of a precision effect in Experiment
442 1 and its absence in Experiments 2 and 3.

443