



City Research Online

City St George's, University of London

Citation: Bagchi, B. & Fring, A. (2009). Comment on "Non-Hermitian Quantum Mechanics with Minimal Length Uncertainty". Symmetry, Integrability and Geometry: Methods and Applications (SIGMA), 5(089), 089. doi: 10.3842/sigma.2009.089

This is the unspecified version of the paper.

This version of the publication may differ from the final published version. To cite this item please consult the publisher's version.

Permanent repository link: <https://city-ref-test.eprints-hosting.org/id/eprint/748/>

Link to published version: <https://doi.org/10.3842/sigma.2009.089>

Copyright and Reuse: Copyright and Moral Rights remain with the author(s) and/or copyright holders. Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge, unless otherwise indicated, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way. For full details of reuse please refer to [City Research Online policy](#).

Comment on “Non-Hermitian Quantum Mechanics with Minimal Length Uncertainty”^{*}

Bijan BAGCHI[†] and Andreas FRING[‡]

[†] Department of Applied Mathematics, University of Calcutta,
92 Acharya Prafulla Chandra Road, Kolkata 700 009, India
E-mail: BBagchi123@rediffmail.com

[‡] Centre for Mathematical Science, City University London,
Northampton Square, London EC1V 0HB, UK
E-mail: A.Fring@city.ac.uk

Received August 18, 2009; Published online September 17, 2009

doi:10.3842/SIGMA.2009.089

Abstract. We demonstrate that the recent paper by Jana and Roy entitled “Non-Hermitian quantum mechanics with minimal length uncertainty” [*SIGMA* 5 (2009), 083, 7 pages, arXiv:0908.1755] contains various misconceptions. We compare with an analysis on the same topic carried out previously in our manuscript [arXiv:0907.5354]. In particular, we show that the metric operators computed for the deformed non-Hermitian Swanson models differs in both cases and is inconsistent in the former.

Key words: non-Hermitian Hamiltonians; deformed canonical commutation relations; minimal length

2000 Mathematics Subject Classification: 81Q10; 46C15; 81Q12

It is known for some time that the deformations of the standard canonical commutation relations between the position operator P and the momentum operator X will inevitably lead to a minimal length, that is a bound beyond which the localization of space-time events are no longer possible. In a recent manuscript [1] we investigated various limits of the q -deformation relations

$$[X, P] = i\hbar q^{f(N)}(\alpha\delta + \beta\gamma) + \frac{i\hbar(q^2 - 1)}{\alpha\delta + \beta\gamma}(\delta\gamma X^2 + \alpha\beta P^2 + i\alpha\delta XP - i\beta\gamma PX),$$

in conjunction with the constraint $4\alpha\gamma = (q^2 + 1)$, with $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ and f being an arbitrary function of the number operator N . One may consider various types of Hamiltonian systems, either Hermitian or non-Hermitian, and replace the original standard canonical variables (x_0, p_0) , obeying $[x_0, p_0] = i\hbar$, by (X, P) . It is crucial to note that even when the undeformed Hamiltonian is Hermitian $H(x_0, p_0) = H^\dagger(x_0, p_0)$ the deformed Hamiltonian is inevitably non-Hermitian $H(X, P) \neq H^\dagger(X, P)$ as a consequence of the fact that X and/or P are no longer Hermitian. Of course one may also deform Hamiltonians, which are already non-Hermitian when undeformed $H(x_0, p_0) \neq H^\dagger(x_0, p_0)$. In both cases a proper quantum mechanical description requires the re-definition of the metric to compensate for the introduction of non-Hermitian variables and in the latter an additional change due to the fact that the Hamiltonian was non-Hermitian in the first place.

In a certain limit, as specified in [1], X and P allow for a well-known representation of the form $X = (1 + \tau p_0^2)x_0$ and $P = p_0$, which in momentum space, i.e. $x_0 = i\hbar\partial_{p_0}$, corresponds to the one used by Jana and Roy [2], up to an irrelevant additional term $i\hbar\tilde{\gamma}P$. (Whenever constants with the same name but different meanings occur in [2] and [1] we dress the former with a tilde.)

^{*}This paper is a contribution to the Proceedings of the 5-th Microconference “Analytic and Algebraic Methods V”. The full collection is available at <http://www.emis.de/journals/SIGMA/Prague2009.html>

The additional term can simply be gauged away and has no physical significance. Jana and Roy have studied the non-Hermitian displaced harmonic oscillator and the Swanson model. As we have previously also investigated the latter in [1], we shall comment on the differences. The conventions in [2] are

$$H_{\text{JR}}(a, a^\dagger) = \omega a^\dagger a + \lambda a^2 + \tilde{\delta}(a^\dagger)^2 + \frac{\omega}{2}$$

with $\lambda \neq \tilde{\delta} \in \mathbb{R}$ and $a = (P - i\omega X)/\sqrt{2m\hbar\omega}$, $a^\dagger = (P + i\omega X)/\sqrt{2m\hbar\omega}$, whereas in [1] we used

$$H_{\text{BF}}(X, P) = \frac{P^2}{2m} + \frac{m\omega^2}{2}X^2 + i\mu\{X, P\}$$

with $\mu \in \mathbb{R}$ as a starting point. Setting $\hbar = m = 1$ it is easy to see that the models coincide when $\lambda = -\tilde{\delta}$ and $\mu = \tilde{\delta} - \lambda$. The Hamiltonians exhibit a “twofold” non-Hermiticity, one resulting from the fact that when $\lambda \neq \tilde{\delta}$ even the undeformed Hamiltonian is non-Hermitian and the other resulting from the replacement of the Hermitian variables (x_0, p_0) by (X, P) . The factor of the metric operator to compensate for the non-Hermiticity of X coincides in both cases, but the factor which is required due to the non-Hermitian nature of the undeformed case differs in both cases

$$\rho_{\text{BF}} = e^{2\mu P^2} \quad \text{and} \quad \rho_{\text{JR}} = (1 + \tau P^2)^{\frac{\mu}{\omega^2 \tau}}.$$

We have made the above identifications such that $H_{\text{JR}}(a, a^\dagger) = H_{\text{BF}}(X, P)$ and replaced the deformation parameter β used in [2] by τ employed in [1]. It is well known that when given only a non-Hermitian Hamiltonian, the metric operator can not be uniquely determined. However, as argued in [1] with the specification of the observable X , which coincides in [2] and [1], the outcome is unique and we can therefore directly compare ρ_{BF} and ρ_{JR} . The limit $\tau \rightarrow 0$ reduces the deformed Hamiltonian $H_{\text{JR}} = H_{\text{BF}}$ to the standard Swanson Hamiltonian, such that ρ_{JR} and ρ_{BF} should acquire the form of a previously constructed metric operator. This is indeed the case for ρ_{BF} , but not for ρ_{JR} . In fact it is unclear how to carry out this limit for ρ_{JR} and we therefore conclude that the metric ρ_{JR} is incorrect.

References

- [1] Bagchi B., Fring A., Minimal length in quantum mechanics and non-Hermitian Hamiltonian systems, arXiv:0907.5354.
- [2] Jana T.K., Roy P., Non-Hermitian quantum mechanics with minimal length uncertainty, *SIGMA* **5** (2009), 083, 7 pages, arXiv:0908.1755.